



# e-Interventions for Alcohol Misuse

September 2014

## Prepared for:

Department of Veterans Affairs  
Veterans Health Administration  
Quality Enhancement Research Initiative  
Health Services Research and Development Service  
Washington, DC, 20420

## Prepared by:

Evidence-based Synthesis Program (ESP) Center  
Durham Veterans Affairs Healthcare System  
Durham, NC  
John W. Williams, Jr., MD, MHSc, Director

## Investigators:

Eric Dedert, PhD  
John W. Williams, Jr., MD, MHSc

## Co-Investigators:

Roy Stein, MD  
J. Murray McNeil, PhD  
Jennifer McDuffie, PhD  
Isabel Ross, MD  
Caroline Feiermuth, MD  
Adam Hemminger, BA  
Andrzej Kosinski, PhD

## Research Associates:

Avishek Nagi, MS



Defining  
**EXCELLENCE**  
in the 21st Century

## PREFACE

Quality Enhancement Research Initiative's (QUERI) Evidence-based Synthesis Program (ESP) was established to provide timely and accurate syntheses of targeted healthcare topics of particular importance to Veterans Affairs (VA) clinicians, managers and policymakers as they work to improve the health and healthcare of Veterans. The ESP disseminates these reports throughout the VA, and some evidence syntheses inform the clinical guidelines of large professional organizations.

QUERI provides funding for four ESP Centers and each Center has an active university affiliation. The ESP Centers generate evidence syntheses on important clinical practice topics, and these reports help:

- develop clinical policies informed by evidence;
- guide the implementation of effective services to improve patient outcomes and to support VA clinical practice guidelines and performance measures; and
- set the direction for future research to address gaps in clinical knowledge.

In, 2009, the ESP Coordinating Center was created to expand the capacity of HSR&D Central Office and the four ESP sites by developing and maintaining program processes. In addition, the Center established a Steering Committee comprised of QUERI field-based investigators, VA Patient Care Services, Office of Quality and Performance, and Veterans Integrated Service Networks (VISN) Clinical Management Officers. The Steering Committee provides program oversight, guides strategic planning, coordinates dissemination activities, and develops collaborations with VA leadership to identify new ESP topics of importance to Veterans and the VA healthcare system.

Comments on this evidence report are welcome and can be sent to Nicole Floyd, ESP Coordinating Center Program Manager, at [Nicole.Floyd@va.gov](mailto:Nicole.Floyd@va.gov).

**Recommended citation:** Dedert E, Williams JW Jr., Stein R, McNeil JM, McDuffie J, Ross I, Feiermuth C, Hemminger A, Kosinski A, Nagi A. Evidence Report: E-Interventions for Alcohol Misuse. VA ESP Project #09-010; 2014.

This report is based on research conducted by the Evidence-based Synthesis Program (ESP) Center located at the Durham VA Medical Center, Durham, NC, funded by the Department of Veterans Affairs, Veterans Health Administration, Office of Research and Development, Quality Enhancement Research Initiative. The findings and conclusions in this document are those of the author(s) who are responsible for its contents; the findings and conclusions do not necessarily represent the views of the Department of Veterans Affairs or the United States government. Therefore, no statement in this article should be construed as an official position of the Department of Veterans Affairs. No investigators have any affiliations or financial involvement (eg, employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties) that conflict with material presented in the report.



# TABLE OF CONTENTS

<a href="#">EXECUTIVE SUMMARY</a> .....	1
<a href="#">Introduction</a> .....	1
<a href="#">Methods</a> .....	2
<a href="#">Data Sources and Searches</a> .....	2
<a href="#">Study Selection</a> .....	2
<a href="#">Data Abstraction and Quality Assessment</a> .....	2
<a href="#">Data Synthesis and Analysis</a> .....	3
<a href="#">Results</a> .....	3
<a href="#">Results of Literature Search</a> .....	3
<a href="#">Summary of Results for Key Questions (KQs)</a> .....	4
<a href="#">Discussion</a> .....	7
<a href="#">Key Findings and Strength of Evidence</a> .....	7
<a href="#">Clinical and Policy Implications</a> .....	8
<a href="#">Strengths and Limitations</a> .....	9
<a href="#">Applicability</a> .....	9
<a href="#">Research Gaps/Future Research</a> .....	10
<a href="#">Conclusions</a> .....	10
<a href="#">Abbreviations Table</a> .....	10
<a href="#">INTRODUCTION</a> .....	11
<a href="#">METHODS</a> .....	14
<a href="#">Topic Development</a> .....	14
<a href="#">Search Strategy</a> .....	14
<a href="#">Study Selection</a> .....	15
<a href="#">Data Abstraction</a> .....	16
<a href="#">Quality Assessment</a> .....	18
<a href="#">Data Synthesis</a> .....	18
<a href="#">Rating the Body of Evidence</a> .....	19
<a href="#">Peer Review</a> .....	20
<a href="#">RESULTS</a> .....	21
<a href="#">Literature Flow</a> .....	22
<a href="#">Description of Included Studies</a> .....	23
<a href="#">Key Question 1: For e-interventions targeting adults who misuse alcohol or who have a diagnosis of AUD, what level, type, and modality of user support is provided (eg, daily telephone calls, weekly email correspondence), by whom (eg, professional counselor, technical support staff), and in what clinical context (adjunct to therapy or primary intervention)?</a> .....	23
<a href="#">Key Points</a> .....	23
<a href="#">General Description</a> .....	24
<a href="#">Level of Support</a> .....	25



Intervention Intensity: Number of Planned Sessions and Mean Duration of Planned Sessions.....	26
Delivery Mode and Delivery Location: Online, Desktop, IVR, and On- or Off-Site.....	27
E-Intervention Content: Theoretical Basis and Most Common Features .....	27
Summary of Findings.....	29
Key Question 2: For adults who misuse alcohol but do not meet diagnostic criteria for AUD, what are the effects of e-interventions compared with inactive controls? .....	29
Key Points.....	29
Results from Studies of Adults .....	31
Results from Studies of College Students.....	34
Summary of Findings.....	38
Quality of Evidence for KQ 2.....	38
Key Question 3: For adults at high risk of AUD (eg, AUDIT-C $\geq$ 8), or who have a diagnosis of AUD, what are the effects of e-interventions compared with inactive controls?.....	38
Key Points.....	38
Study Characteristics and Treatment Effects .....	39
Summary of Findings.....	42
Quality of Evidence for KQ 3.....	42
Key Question 4: For adults who misuse alcohol, are at high risk of AUD, or have a diagnosis of AUD, what are the effects of e-interventions alone or used in combination with face-to-face therapy compared with face-to-face therapy alone?.....	42
Key Points.....	42
Study Characteristics .....	43
E-Intervention Characteristics .....	45
Alcohol Consumption.....	45
Other Outcomes .....	47
Summary of Findings.....	47
Quality of Evidence for KQ 4.....	47
<b>SUMMARY AND DISCUSSION</b> .....	49
Summary of Evidence by Key Question .....	52
KQ 1. Characteristics of and User Support for E-Interventions .....	52
KQ 2. Effects of E-Interventions Compared with Inactive Controls in Adults who Misuse Alcohol.....	52
KQ 3. Effects of E-Interventions Compared with Inactive Controls in Adults at High Risk of AUD (eg, AUDIT-C $\geq$ 8) or with a Diagnosis of AUD .....	52
KQ 4. Effects of E-Interventions Alone or Used in Combination with Face-to-Face Therapy Compared with Face-to-Face Therapy Alone in Adults who Misuse Alcohol .....	53
Clinical and Policy Implications .....	53
Limitations.....	55
Publication Bias .....	56
Study Quality .....	56
Heterogeneity.....	56
Applicability of Findings to the VA Population.....	56
Research Gaps/Future Research .....	57
Conclusions .....	57



<a href="#">REFERENCES</a> .....	<a href="#">59</a>
----------------------------------	--------------------

## [TABLES](#)

<a href="#">Table ES-1. Summary SOE Ratings</a> .....	<a href="#">8</a>
<a href="#">Table 1. Definitions of the Spectrum of Alcohol Misuse</a> .....	<a href="#">11</a>
<a href="#">Table 2. Inclusion and Exclusion Criteria</a> .....	<a href="#">15</a>
<a href="#">Table 3. Characteristics of E-Interventions (KQ 1)</a> .....	<a href="#">24</a>
<a href="#">Table 4. Examples of Level 2 and Level 3 Interventions (KQ 1)</a> .....	<a href="#">28</a>
<a href="#">Table 5. Characteristics of E-Interventions (KQ 2)</a> .....	<a href="#">30</a>
<a href="#">Table 6. Study Characteristics (KQ 3)</a> .....	<a href="#">40</a>
<a href="#">Table 7. Study Characteristics (KQ 4)</a> .....	<a href="#">44</a>
<a href="#">Table 8. Characteristics of E-Interventions (KQ 4)</a> .....	<a href="#">45</a>
<a href="#">Table 9. Detailed SOE Ratings</a> .....	<a href="#">50</a>
<a href="#">Table 10. Highest Priority Evidence Gaps</a> .....	<a href="#">57</a>

## [FIGURES](#)

<a href="#">Figure ES-1. Alcohol Consumption at 6 Months in Studies of Adults</a> .....	<a href="#">5</a>
<a href="#">Figure ES-2. Alcohol Consumption at 6 Months in Studies of College Students</a> .....	<a href="#">6</a>
<a href="#">Figure 1. Literature Flow Chart</a> .....	<a href="#">22</a>
<a href="#">Figure 2. Strategies Used in E-Interventions</a> .....	<a href="#">28</a>
<a href="#">Figure 3. Alcohol Consumption at 6 Months in Studies of Adults</a> .....	<a href="#">32</a>
<a href="#">Figure 4. Alcohol Consumption at 12 Months in Studies of Adults</a> .....	<a href="#">32</a>
<a href="#">Figure 5. Alcohol Reduction to Meet Drinking Limit Guidelines at 6 Months in Studies of Adults</a> .....	<a href="#">33</a>
<a href="#">Figure 6. Alcohol Consumption at 6 Months in Studies of College Students</a> .....	<a href="#">35</a>
<a href="#">Figure 7. Alcohol Consumption at 12 Months in Studies of College Students</a> .....	<a href="#">35</a>
<a href="#">Figure 8. Episodes of Binge Drinking at 6 Months in Studies of College Students</a> .....	<a href="#">36</a>
<a href="#">Figure 9. Negative Social Consequences of Drinking at 6 Months in Studies of College Students</a> .....	<a href="#">37</a>
<a href="#">Figure 10. Negative Social Consequences of Drinking at 12 Months in Studies of College Students</a> .....	<a href="#">37</a>

<a href="#">APPENDIX A. SEARCH STRATEGIES</a> .....	<a href="#">65</a>
---	--------------------

<a href="#">APPENDIX B. CRITERIA USED IN QUALITY (RISK OF BIAS) ASSESSMENT OF RCTs</a> .....	<a href="#">68</a>
--	--------------------

<a href="#">APPENDIX C. QUALITY OF INCLUDED STUDIES</a> .....	<a href="#">70</a>
---	--------------------

<a href="#">APPENDIX D. PEER REVIEW COMMENTS/AUTHOR RESPONSES</a> .....	<a href="#">73</a>
---	--------------------

<a href="#">APPENDIX E. STUDY CHARACTERISTICS</a> .....	<a href="#">86</a>
---	--------------------

<a href="#">APPENDIX F. E-INTERVENTION CHARACTERISTICS</a> .....	<a href="#">92</a>
--	--------------------



## EVIDENCE REPORT: E-INTERVENTIONS FOR ALCOHOL MISUSE

### INTRODUCTION

The economic, social, and health burden of alcohol misuse is widely recognized,<sup>1</sup> as is the need for effective interventions to reduce this burden.<sup>2,3</sup> Alcohol misuse constitutes the third leading cause of morbidity and mortality worldwide<sup>4-6</sup> and is the third leading cause of preventable death in the United States, after tobacco use and obesity.<sup>7,8</sup> Alcohol misuse contributes to a multitude of medical and psychiatric illnesses and presents across all medical specialties. The associated costs amount to more than 1% of the gross national product in high- and middle-income countries.<sup>6,9</sup> Veterans, who account for approximately 21.8 million people in the United States,<sup>10</sup> are among those significantly affected by addiction.<sup>11</sup> Substance use disorders, including alcohol use disorder (AUD), are among the most common and costly conditions among Veterans presenting for treatment in the Veterans Health Administration (VHA) system.<sup>12,13</sup> Among Veterans with at least one primary care visit in VHA within the past year, 589,094 (14.5%) screened positive for alcohol misuse, and 396,374 had a diagnosis of AUD with or without another substance use disorder (Daniel Kivlahan, PhD, e-mail communication, July 2014).

Studies on alcohol misuse conceptualize the target problem differently, and the recently introduced *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition (DSM-5),<sup>14</sup> criteria include important revisions to the nosology. The key change for DSM-5 was to integrate “alcohol abuse” and “alcohol dependence” diagnoses into a single disorder, “alcohol use disorder” (AUD), with mild, moderate, and severe subclassifications. To provide clarity on the use of terminology for alcohol consumption patterns and related problems, Table 1 provides operational definitions for the terms used in this report.

**Table 1. Definitions of the Spectrum of Alcohol Misuse\***

Term	Definition (Reference Time Period: 1 Year)
Risky Use or Hazardous Use	Excess daily consumption (>4 drinks/day in men, >3 drinks/day in women and men over age 65) or excess total consumption (>14 drinks/week in men, >7 drinks/week in women and men over age 65) associated with increased risk of health problems.
Harmful use	A pattern of drinking that is already causing damage to health. The damage may be either physical (eg, liver damage) or mental (eg, depressive episodes).
Alcohol abuse (DSM-IV criteria)†	A maladaptive pattern of alcohol use leading to clinically significant impairment or distress (eg, failure to fulfill major obligations). Continued use despite persistent or recurrent social or interpersonal problems caused or exacerbated by alcohol.

Term	Definition (Reference Time Period: 1 Year)
Alcohol dependence (DSM-IV criteria)†	A maladaptive pattern of alcohol use leading to clinically significant impairment or distress, which may include symptoms associated with alcoholism or addiction: tolerance, withdrawal, excessive amounts consumed or time spent drinking, unsuccessful attempts to decrease use, pattern continues despite persistent problems caused by or associated with alcohol.
Alcohol use disorder (AUD) (DSM-5 criteria)†	This new category integrates the 2 DSM-IV disorders “alcohol abuse” and “alcohol dependence” into a single disorder for DSM-5: An individual continues pattern of alcohol use despite significant substance-related problems in one or more of the following areas: impaired control over use (eg, inability to decrease consumption), social impairment (eg, failing an obligation or foregoing a favorite activity), health consequences (physical or mental), and physiological dependence (eg, cravings). The disorder is classified as mild, moderate, or severe depending on number of symptoms.

\*The term “alcohol misuse” (sometimes termed “unhealthy alcohol use”) is an umbrella term for a spectrum of potentially problematic patterns of alcohol use. This table is adapted with permission from Table 1 in Jonas et al, 2012,<sup>8</sup> and uses terminology from the DSM-IV<sup>15</sup> for alcohol abuse and alcohol dependence, and from the DSM-5<sup>14</sup> for AUD. The source table was abbreviated and updated to reflect the DSM-5 terminology for this report in collaboration with Dr. Jonas.

†Not all exact criteria are listed.

Abbreviations: AUD=alcohol use disorder; DSM-5=Diagnostic and Statistical Manual of Mental Disorders, 5th edition; DSM-IV=Diagnostic and Statistical Manual of Mental Disorders, 4th edition

Traditional treatment approaches for AUD, which usually consist of initial intensive and time-limited interventions followed by less intensive follow-up care,<sup>12</sup> can be prohibitive for individuals seeking treatment for a variety of reasons. Even the use of screening and brief interventions for less severe alcohol misuse, which have been shown to be effective, are constrained by barriers such as adequate funding, time, and adequately trained personnel.<sup>16-18</sup> Thus, electronic interventions (e-interventions) may prove to be a useful way to extend the reach of interventions for alcohol misuse. With 87% of the U.S. population using the Internet,<sup>19</sup> e-interventions have the potential to reach individuals with drinking problems who wish to remain anonymous, have little time for traditional therapy, need therapy to be available during non-standard business hours due to shift work, live at great distance from traditional therapy, or cannot afford such therapy.<sup>20,21</sup> Given that Veterans can encounter most, if not all, of the barriers to accessing care for alcohol misuse, e-interventions may prove a promising avenue, especially for the younger, more Internet-savvy Veterans returning from recent deployments in places such as Iraq and Afghanistan.

Prior reviews focus on student or young adult populations,<sup>22-24</sup> do not include the most recent trials,<sup>21,25,26</sup> include studies with only short-term outcomes<sup>27</sup> or studies restricted to web-based interventions,<sup>22,25</sup> or combine alcohol and other substance use in the same analyses.<sup>28</sup> To inform

policy for VHA, we focused on studies that reported longer term, clinically important outcomes and evaluated effects separately for student and non-student populations. We offer a systematic review of randomized controlled trials (RCTs) assessing CD-ROM-based, web-based, interactive voice response (IVR), or mobile applications of e-interventions for alcohol misuse in order to assess for changes in alcohol consumption, alcohol-related health problems, alcohol-related social or legal problems, health-related quality of life (HRQOL), functional status, medical utilization, and adverse effects. We intend the evidence synthesis to be used to inform the decision whether to disseminate e-interventions for alcohol misuse in VHA, and how best to implement programs.



## METHODS

### TOPIC DEVELOPMENT

We followed a standard protocol for this review. The topic was nominated after a process that included a preliminary review of published peer-reviewed literature and consultation with investigators, Veterans Affairs (VA) and non-VA experts, and key stakeholders (Mental Health Web Services, Mental Health Services, and Mental Health Quality Enhancement Research Initiative [QUERI]).

The key questions (KQs) are:

- KQ 1: For e-interventions targeting adults who misuse alcohol or who have a diagnosis of AUD, what level, type, and modality of user support is provided (*eg*, daily telephone calls, weekly email correspondence), by whom (*eg*, professional counselor, technical support staff), and in what clinical context (adjunct to therapy or primary intervention)?
- KQ 2: For adults who misuse alcohol but do not meet diagnostic criteria for AUD, what are the effects of e-interventions compared with inactive controls?
- KQ 3: For adults at high risk of AUD (*eg*, AUDIT-C  $\geq 8$ ), or who have a diagnosis of AUD, what are the effects of e-interventions compared with inactive controls?
- KQ 4: For adults who misuse alcohol, are at high risk of AUD, or have a diagnosis of AUD, what are the effects of e-interventions alone or used in combination with face-to-face therapy compared with face-to-face therapy alone?

### SEARCH STRATEGY

We conducted a primary review of the literature by systematically searching, reviewing, and analyzing the scientific evidence as it pertains to the KQs. To identify relevant articles, in consultation with an expert librarian, we searched MEDLINE (via PubMed), The Cochrane Library, Embase, and PsycINFO from January 1, 2000 to August 18, 2014, for peer-reviewed publications of trials that compared e-interventions with a waitlist control, usual care, or face-to-face therapy in adults who misuse alcohol or who have a diagnosis of AUD.

We used Medical Subject Heading (MeSH) terms and selected free-text terms for the conditions of interest; cognitive-behavioral therapy and closely related therapies, including brief counseling and health education; and the electronic delivery mode, including computer-assisted, Internet, and terms for mobile devices. We added validated search terms for RCTs.<sup>29</sup> We limited the search to RCTs published in English. The exact search strategies used are provided in Appendix A. We further searched the bibliographies of exemplar trials and systematic reviews for missed publications.<sup>23-28,30,31</sup>

To assess for possible publication bias, we searched ClinicalTrials.gov ([www.clinicaltrials.gov](http://www.clinicaltrials.gov)) to identify completed but unpublished studies meeting our eligibility criteria.

All citations were imported into 2 electronic databases (for referencing, EndNote® Version X5, Thomson Reuters, Philadelphia, PA; for data abstraction, DistillerSR; Evidence Partners Inc., Manotick, ON, Canada).

## STUDY SELECTION

Using prespecified inclusion and exclusion criteria, 2 investigators assessed titles and abstracts for relevance to the KQs. Full-text articles identified by either investigator as potentially relevant were retrieved for further review and examined by 2 investigators against the eligibility criteria. Disagreements on inclusion, exclusion, or the major reason for exclusion were resolved by discussion or by a third investigator. The criteria to screen articles for inclusion or exclusion at both the title-and-abstract and full-text screening stages are detailed in Table 2. In addition, trials with 3 or more arms were examined for appropriateness of all arms for inclusion. For example, any active arm that did not include an e-intervention or evidence-based face-to-face treatment was not abstracted for inclusion in the analysis.

**Table 2. Inclusion and Exclusion Criteria**

Study Characteristic	Inclusion Criteria	Exclusion Criteria
Population	Adults 18 years or older with alcohol misuse (eg, positive alcohol screen; KQ 2), at high risk of AUD, or with diagnosis of AUD (KQ 3)	Pregnant women
Intervention	Intervention must be a computer-based therapy adhering to evidence-based treatment principles and providing individually delivered treatment for alcohol misuse delivered by CD-ROM, web-based, IVR, mobile phones, or in-home electronic devices (eg, Health Buddy™) and may be combined with various levels of supplementary human support.	Interventions targeted at dyads (eg, couple) or primary prevention  Computerized screening only
Comparator	The comparator for KQ 2 and KQ 3 was usual care not involving psychotherapy; waitlist; or information or attention control. For KQ 4, the comparator was face-to-face treatment.	Any comparator where the effect of the electronic aspect of the intervention could not be isolated
Outcome	Studies must report effects on at least one of the following relevant outcomes: alcohol consumption, alcohol-related health problems, alcohol-related legal or social problems, HRQOL, functional status measures, medical utilization, or adverse effects from treatment.	–
Timing	Outcomes reported at ≥6 months from randomization and initiation of intervention	Outcomes reported at <6 months
Setting	Outpatients in any setting (general medical, emergency room, and community) or participants not engaged in clinical care who are enrolled through self-assessments. We included studies where enrollment was inpatient but the majority of the intervention was delivered outpatient.	Inpatient settings for intervention delivery
Study design	RCTs with n ≥50. The sample size requirement is designed to exclude small pilot studies that typically are underpowered and have more methodological problems than larger trials. Studies with small samples and no treatment effect are also less likely to be published, increasing the risk of publication bias.	RCTs with n <50

Study Characteristic	Inclusion Criteria	Exclusion Criteria
Publications	English-language publication Published from 2000 to present* Peer-reviewed, full publication Study conducted in North America, the European Union, or Australia/New Zealand†	Non-English language Published before 2000 Abstract only

\*Rationale is that the Internet was developed in the 1990s and not used routinely for interventions until 2000. Based on our assessment of studies included in existing systematic reviews, the earliest relevant publication was in 2004.

†Rationale is to include economically developed countries with sufficient similarities in healthcare system and culture to be applicable to U.S. medical care.

Abbreviations: AUD=alcohol use disorder; CD-ROM=compact disc read-only memory; HRQOL=health-related quality of life; IVR=interactive voice response; KQ=key question; n=number of participants; RCTs=randomized controlled trials

## DATA ABSTRACTION

Before general use, the abstraction form templates, designed specifically for this report, were piloted on a sample of included articles and revised to ensure that all relevant data elements were captured and that there was consistency and reproducibility between abstractors. Data elements included descriptors to assess applicability, quality elements, intervention/exposure details, and outcomes. Key data elements abstracted included patient descriptors (including age, education, baseline alcohol use); setting, features and dose of the e-intervention, characteristics of the comparator, and outcomes as described in Table 2. Key features relevant to applicability included the match between the sample and target populations (*eg*, age, education level) and the training and experience of the clinician. Data from published reports were then abstracted into the final abstraction form by a trained investigator. All data abstractions were confirmed by a second investigator. Disagreements were resolved by consensus or by obtaining a third investigator's opinion. When data from published reports were missing or incomplete, we contacted study authors to request the data.

We abstracted the following key information for each included study:

- Study characteristics and design
  - Study identifiers—Author last name, year published, ID number
  - Study design—RCT, patient level or group level
  - Location (country) and recruitment setting (clinic, *etc*) of study
  - Number of study arms; types of comparison groups
  - Inclusion and exclusion criteria (identified by screen only or diagnostic criteria, *etc*)
  - Recruitment (*eg*, internet, advertisement); source (*eg*, campus, clinic)
  - Number of participants eligible for, randomized, enrolled in, and completing study
  - Analysis method for alcohol consumption: analysis of covariance (ANCOVA), mixed model, other
  - Did analysis account for missing data? (*ie*, mixed models or multiple imputation)
- Population characteristics
  - Sex, race, and age of sample

- Inclusion of active duty or Veteran participants
- Alcohol consumption at baseline (*eg*, drinks/week)
- Are there important differences in alcohol diagnosis or severity (amount of consumption)?
- Description of the intervention and comparator
  - Comparator used (*eg*, waitlist, treatment as usual, informational or attention control, face-to-face treatment)
  - Comparator delivery, if applicable
  - Overall e-intervention classification (degree of support provided)
  - Type, delivery mode, and location of computer platform
  - “Brand” name of intervention
  - Theoretical basis of intervention (cognitive-behavioral therapy, motivational interviewing, *etc*)
  - Components of intervention
    - Group or individual
    - Number of sessions and minutes per session
    - Types of interactivity (*eg*, email, text messaging, telephone, peer component)
    - Techniques used (*eg*, personalized normative feedback [PNF], psychoeducation, negative consequences, tailored materials, *etc*)
  - Therapist credentials, level of therapist support (feedback, email, phone, *etc*)
  - Technical support, if offered
- Outcomes
  - Time points measured (number of follow-up assessments)
  - Study duration (longest follow-up, in weeks)
  - Treatment adherence: mean sessions completed or proportion completing all sessions
  - Patient satisfaction
  - Alcohol consumption (measured at time points closest to 6 and 12 months)
    - Standard drinks per week (and number of grams of alcohol per drink, if given)
    - Heavy drinking episodes (and definition of qualifying episode)
    - Achieved recommended drinking limits
  - Health outcomes
    - HRQOL
    - Alcohol-related accidents
    - Alcohol-related medical problems
    - Mortality
  - Utilization outcomes
    - Hospitalizations
    - Emergency department visits
  - Social or legal problems

## QUALITY ASSESSMENT

We abstracted data necessary to assess the quality (risk of bias) of included trials. Across all included trials, quality criteria were applied for each RCT by 2 independent investigators. Disagreements were resolved between the 2 investigators or, when needed, by arbitration from a third investigator. We used the key risk of bias criteria described in the Agency for Healthcare Research and Quality's (AHRQ's) *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*,<sup>32</sup> adapted to this specific topic and customized to RCTs (Appendix B). These criteria are: adequacy of randomization and allocation concealment; comparability of groups at baseline; blinding; completeness of follow-up and differential loss to follow-up; whether incomplete data were addressed appropriately; validity of outcome measures; and conflict of interest. We assigned a summary risk of bias score (low, moderate, or high) to individual studies. Detailed quality ratings for each included study are provided in Appendix C.

## DATA SYNTHESIS

While synthesizing relevant abstracted data, we developed a summary table describing the key outcomes used to test e-interventions in included RCTs. As part of this process, we classified the e-interventions according to the level of supplementary human support provided as follows:

- Level 1 (minimal): No support; e-intervention only
- Level 2 (low): Some support; e-intervention supplemented by non-counseling meetings with study staff
- Level 3 (moderate to high): Therapeutic support; e-intervention supplemented by counseling with trained staff

We grouped studies into those with eligibility criteria designed to enroll participants with alcohol misuse and those designed to enroll participants at high risk of or with AUD (*ie*, DSM-5 criteria or high threshold on screening test). Controls were grouped into inactive (waitlist, attention/information, and treatment as usual) and face-to-face active controls.

We then determined the feasibility of completing a quantitative synthesis (meta-analysis) to estimate summary effects. Feasibility depends on the volume of relevant literature, conceptual homogeneity of the trials, and completeness of results reporting. Because of important differences in the study samples and intervention designs, we planned a priori to analyze studies conducted in college samples separately from studies conducted in other adult samples.

When meta-analysis was feasible, we computed summary estimates of effect, stratified by condition (alcohol misuse versus at risk of AUD or with AUD), for both end-of-treatment and longest follow-up point  $\geq 6$  months. When trials evaluated more than one e-intervention, we averaged the effect across e-interventions.<sup>33-36</sup> Meta-analyses were feasible only for the e-interventions compared with inactive controls. Because the primary outcome—alcohol consumption—was measured across the trials using different units, the measurements were converted to a common unit (grams/week) and were combined using mean differences (MDs).<sup>37,38</sup> If means and standard deviations (SDs) were not reported, we used other statistics (*eg*, median, F values) to calculate effect sizes. Because of the relatively small number of studies, we used

a random-effects model with the Knapp and Hartung method to adjust the standard errors of the estimated coefficients.<sup>39,40</sup> The dichotomous outcome “met drinking limits” was analyzed using summary risk ratios (RRs). For outcomes that used different measures to evaluate the construct (eg, social or legal problems from alcohol misuse), we computed summary estimates using standardized mean differences (SMDs). At each time point, the SMD was calculated by subtracting the average intake of the intervention group from the average intake of the control group and dividing the result by the pooled standard deviations of the 2 groups. SMDs of 0.2 can be considered small treatment effects; 0.5, moderate effects; and  $\geq 0.8$ , large effects.<sup>41</sup> We evaluated for statistical heterogeneity in treatment effects using Cochran’s Q and I<sup>2</sup> statistics. An I<sup>2</sup> of 0% indicates no observed heterogeneity, and larger values suggest increasing heterogeneity: 25% is interpreted as low, 50% as moderate, and  $\geq 75\%$  as high heterogeneity.<sup>42</sup>

We planned subgroup analyses to explore potential sources of heterogeneity, specifying a priori: follow-up rates, treatment dose, and the level of support given with the intervention. Consistent with AHRQ’s *Methods Guide*, we specified a minimum of 4 studies per subgroup to conduct the planned analyses.<sup>43</sup> However, because of the relatively small number of studies and lack of variability in moderator variables, subgroup analyses were not performed. When there were at least 3 studies at low or moderate risk of bias, we performed sensitivity analyses to compute summary estimates after excluding studies at high risk of bias.

We used R (R Foundation for Statistical Computing, Vienna, Austria) with the metafor package<sup>44</sup> to calculate the summary estimates of treatment effect. Publication bias was assessed using findings from a search of ClinicalTrials.gov. Funnel plots were not used because analyses did not meet the minimum threshold of at least 10 studies for meaningful analysis.<sup>32</sup>

Where quantitative synthesis was not feasible, we analyzed the data qualitatively. We gave more weight to the evidence from higher quality studies with more precise estimates of effect. A precise estimate of effect was defined as one that excluded a difference of  $\leq 3$  standard drinks per week or a SMD of  $\leq 0.4$ . The qualitative syntheses focused on documenting and identifying patterns in efficacy and safety of the interventions across conditions and outcome categories. We also analyzed potential reasons for inconsistency in treatment effects across studies by evaluating differences in the study population, intervention, comparator, and outcome definitions.

## RATING THE BODY OF EVIDENCE

In addition to rating the quality of individual studies, we evaluated the overall strength of evidence (SOE) for each KQ as described in AHRQ’s *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.<sup>32</sup> In brief, this approach requires assessment of 4 domains: risk of bias, consistency, directness, and precision. These domains, along with evidence for publication bias, were considered qualitatively, and a summary rating of high, moderate, low, or insufficient SOE was assigned after discussion by 2 investigators. The 4-level rating scale consists of the following definitions:

- **High**—We are very confident that the true effect lies close to the estimate of the effect.
- **Moderate**—We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

- **Low**—Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect.
- **Insufficient**—Evidence on an outcome is absent or too weak, sparse, or inconsistent to estimate an effect.

When a rating of high, moderate, or low was not possible or was imprudent to make, a rating of insufficient was assigned.

## PEER REVIEW

A draft of this report was reviewed by technical experts and clinical leadership. A transcript of their comments and our responses is provided in Appendix D.

## RESULTS


### LITERATURE FLOW

The flow of articles through the literature search and screening process is illustrated in Figure 1. We identified 704 unique citations from a combined search of PubMed (n=319), Embase (n=154), PsycINFO (n=14), and The Cochrane Library (n=19) and a separate search for IVR studies (n=198) conducted from January 1, 2000, through August 18, 2014. We also searched the bibliographies of seminal, exemplar studies for articles not retrieved by our search (n=18), for a total of 722 citations. After applying inclusion and exclusion criteria at the title-and-abstract level, 84 full-text articles were retrieved and screened. Of these, 53 were excluded at the full-text screening stage, leaving 31 articles (representing 26 unique trials reported in 25 articles and 6 companion papers) for data abstraction. Monahan et al<sup>45</sup> described 2 different trials in the same publication. It was necessary to contact 12 authors for clarification of abstracted elements during the course of the data abstraction process. Eleven authors responded with the requested information.

We searched ClinicalTrials.gov ([www.clinicaltrials.gov](http://www.clinicaltrials.gov)) to identify completed but unpublished studies that might meet our eligibility criteria. We found little evidence of publication bias: there were only 2 completed studies for which we could not identify a publication (NCT000658398<sup>46</sup> and NCT01923246<sup>47</sup>).



Figure 1. Literature Flow Chart



\*Search results from PubMed (319), Embase (154), PsycINFO (14), and The Cochrane Library (19), IVR search (198) and manually identified studies (18)

†Manuscript reference list includes additional references cited for background and methods.

‡One article reported data from 2 trials that we analyzed separately.

## DESCRIPTION OF INCLUDED STUDIES

We identified 26 trials that met eligibility criteria. Eleven trials were conducted in the United States (one in a VA sample); 15 were conducted in the European Union, Australia, or Canada. The populations were divided between college students (n=12) and other groups of adults (n=14). Adult studies enrolled midlife adults, the majority of whom had some college education (median 52%, range 8% to 83%). Across all studies, men and women were well-represented. Twenty-three trials recruited subjects who misused alcohol but did not meet criteria for AUD. Only 3 trials specifically recruited subjects who were at high risk of or had been diagnosed with AUD, but in many of the adult studies the mean weekly alcohol consumption exceeded 14 U.S. standard drinks per week. Study exclusion criteria were limited: 4 trials specifically excluded those in alcohol treatment, 3 excluded those with psychosis, and 2 excluded those with severe or terminal medical illness. Most trials were judged to be at moderate risk of bias (n=14); 5 were at low risk, and 7 were assessed as being at high risk of bias. Twenty-three trials compared an e-intervention with an inactive control, while 3 trials compared IVR (e-intervention via telephone lines) with inactive or active controls. Six studies compared e-interventions or IVR to a face-to-face control condition. A single trial used a mobile device as the delivery platform; none used the Health Buddy™ appliance. The most commonly reported outcomes were the characteristics of the interventions and their effects on alcohol consumption, reductions in consumption to meet drinking limits, binge drinking, and the social and legal consequences of drinking.

**KEY QUESTION 1: For e-interventions targeting adults who misuse alcohol or who have a diagnosis of AUD, what level, type, and modality of user support is provided (eg, daily telephone calls, weekly email correspondence), by whom (eg, professional counselor, technical support staff), and in what clinical context (adjunct to therapy or primary intervention)?**

### Key Points

- The majority of interventions (n=14) did not utilize supplementary human support and were delivered online or at a desktop computer. A single study utilized a mobile device.
- Most interventions were a single session, designed to moderate alcohol consumption in individuals who screened positive on an alcohol questionnaire (eg, Alcohol Use Disorders Identification Test [AUDIT] or AUDIT for Clinicians [AUDIT-C]).
- The main components of the e-interventions were personalized normative feedback (PNF) and psychoeducation or alcohol-specific education including the negative consequences of drinking.
- When supplementary human support was utilized (n=12), it was limited, consisting only of technical support from a research assistant in half the cases. In other cases, it was often given in combination with IVR or combined with telephone or face-to-face treatment in subjects at high risk of or with AUD.

## General Description

We included 26 RCTs, described in 25 publications,<sup>33-36,45,48-67</sup> the paper by Monahan and colleagues<sup>45</sup> reported results from 2 separate trials. All 26 trials were conducted in adults 18 years of age or older, but 46% were conducted specifically in college students. The included RCTs evaluated 21 unique e-interventions—IVR (3 trials) or computer-based programs (23 trials). A single trial used a mobile device as the delivery platform;<sup>52</sup> none used the Health Buddy™ appliance. Seventeen of the interventions used commercially available programs; however, 11 of these were each evaluated in a single study. Three programs were evaluated in 2 studies each—Electronic Check-Up to Go (e-CHUG), Alcohol 101, and Brief Alcohol Screening and Intervention for College Students (BASICS)—and these were all used in student studies. Study characteristics are described in greater details in Appendix E. Detailed descriptions of the e-interventions evaluated are provided in Appendix F. A summary of the adult versus student studies is given in Table 3.

In terms of the level of supplementary human support provided, we classified the e-interventions evaluated in the included studies as follows:

- Level 1 (minimal, n=14): No support; e-intervention only
- Level 2 (low, n=8): Some support; e-intervention supplemented by non-counseling meetings with study staff
- Level 3 (moderate or high, n=4): Therapeutic support; e-intervention supplemented by counseling from trained staff

**Table 3. Characteristics of E-Interventions (KQ 1)**

Characteristic	Adult Studies (n=14)	Student Studies (n=12)
Level of support:		
1 (minimal)	6	8
2 (low)	4	4
3 (moderate or high)	4	0
Number of sessions:		
1	6	8
>1	5 (3=daily IVR)	4 (2–5 sessions)
NR	3	0
Session duration:		
Median (range)	10 (10-90) minutes	30 (5-50) minutes
NR	7	0
NA (IVR)	3	0
Intervention name:		
IVR	3	0
Not named	3	3
Alcohol 101	0	2
e-CHUG	0	2
BASICS	0	2
Program used only once*	8	3

Characteristic	Adult Studies (n=14)	Student Studies (n=12)
Delivery mode:		
Accessed via the Web	8	8
Software on laptop or desktop	1	4
Mobile device	1	0
IVR	3	0
NR	1	0
Delivery location:		
Off-site (home, IVR, etc)	5	2
On-site (clinic, classroom, etc)	3	5
NR	6	5
Content of e-intervention:		
Brief intervention	10	8
PNF <sup>†</sup>	GS: 6; non-GS: 1; NR: 1	GS: 7; non-GS: 2
Psychoeducation	8	8
Alcohol-specific	3 of 8	4 of 8
Goal-setting	7	1
Negative consequences	5	4
Skills training	2	1
Self-monitoring	4	0
Tailored feedback	2	1
Relapse prevention	2	0
Other techniques (used once) <sup>‡</sup>	9	3

\*Programs used only once: in adult trials: FRAMES, eScreen.se, [www.drinktest.nl](http://www.drinktest.nl), Down Your Drink, Check your Drinking, minderdrinken.nl; in student trials: THRIVE, What do You Drink, College Drinkers Check-up

<sup>†</sup>For adults, the comparison group was usually a national population (n=3) or age-matched adults (n=3); for students, the comparison group was usually student peers (n=7).

<sup>‡</sup>Other treatment techniques used: for adults: CBT, computer monitoring, email, GPS, taking responsibility, text messaging, values clarification; for students: homework, decisional balance exercise,

Abbreviations: BASICS=Brief Alcohol Screening and Intervention for College Students; CBT=cognitive-behavioral therapy; e-CHUG=Electronic Check-Up to Go; E-Interventions=electronic interventions; FRAMES=feedback, responsibility, advice, menu of options, empathy, self-efficacy; GPS=global positioning system; GS=gender-specific; IVR=interactive voice response; KQ=key question; NA=not applicable; NR=not reported; PNF=personalized normative feedback; THRIVE=Tertiary Health Research Intervention Via Email

## Level of Support

Over half of the e-interventions (n=14) were classified as level 1 support, which did not provide any supplementary human support, technical or therapeutic.<sup>35,36,45,50,51,53,57,59,61,64-66,68</sup> Eight of these were collegiate studies, and the remainder (n=6) were conducted in other adult groups. All of these studies used computerized PNF as the major component of the intervention.

Four adult<sup>49,52,62,63</sup> and 4 student<sup>34,48,56,58</sup> studies were classified as level 2 support. This usually consisted of technical support from a research assistant, so that human contact was made, but no therapy was given. Among the adult studies, Riper et al<sup>63</sup> used a weekly, moderated peer-support group, and Gustafson et al<sup>52</sup> used both a peer “bulletin board” and email contact with a therapist over the smartphone application. In the other 2 trials,<sup>49,62</sup> interaction with study staff was limited to screening questions and technical aid with the computer program. Among the student studies, one<sup>48</sup> had an in-person assessment in both the active and control conditions in addition to technical help from the staff in the e-intervention arm. In the other 3 level 2 student studies,

the only supplementary human support was technical aid from the research assistant about navigating the program.

All level 3 support studies (n=4) were conducted in non-collegiate adults.<sup>33,54,55,60</sup> Three of these 4 studies<sup>54,55,60</sup> were IVR trials. IVR is a technology that allows a computer to interact with humans through the use of voice and signaling over analog telephone lines. All of the IVR studies were 3-arm studies with one control arm and 2 active arms. In the first,<sup>55</sup> the intervention started with a brief intervention with the participant's primary care provider. Afterwards, participants were randomized to control, IVR, or IVR plus PNF. Participants in IVR groups were instructed to phone in to an interactive program on a daily basis and report their alcohol consumption over the previous 24 hours. Those in the PNF group received monthly PNF from their primary care provider about their consumption in the mail. The second IVR study<sup>54</sup> started with a motivational interview with a counselor who had either a BA or MA in psychology, supervised by a PhD in psychology, which was followed by IVR in one of the active arms. Both active arms received 2 brief (15-minute) follow-up sessions. The third IVR study<sup>60</sup> randomized subjects to IVR only or IVR plus follow-up phone calls with the study coordinator, whose credentials were not specified, for help or guidance in relation to the intervention. The one non-IVR level 3 study<sup>33</sup> also included 3 arms and provided a computerized "expert system intervention" (about which very little information was given) to both active arms, then randomized subjects either to counseling sessions via phone with "trained psychologists" at all follow-up assessment points (full care intervention) or only if needed (stepped care intervention). Whether the extra sessions were needed was determined by the answers to the follow-up questionnaires.

### **Intervention Intensity: Number of Planned Sessions and Mean Duration of Planned Sessions**

Most e-interventions (n=14) consisted of one planned session with the computer program<sup>33,36,45,49-51,53,56,58,59,61,62,66</sup> and offered no further support beyond technical aid. In 5 of these 14 studies, the session lasted 10-20 minutes.<sup>49-51,58,66</sup> Four studies<sup>36,45,56</sup> had slightly more intensive e-interventions that required 30-40 minutes to complete. One study's initial session took 90 minutes to complete.<sup>62</sup> Four studies<sup>33,53,59,61</sup> did not report the length of the e-intervention, but one of these<sup>33</sup> was a level 3 support study in which, although there was an initial computer session, the bulk of the treatment focused on tailored, therapeutic feedback over the phone in up to 4 scheduled phone calls that lasted 30-40 minutes each.

Five studies offered 2 to 5 sessions with the e-intervention;<sup>34,35,48,57,64</sup> 3 studies<sup>52,65,67</sup> offered subjects unlimited access to the program; one was delivered via a mobile phone application.<sup>52</sup> Sinadinovic et al<sup>65</sup> reported in response to a query that the average number of sessions was 2.66 (4.31) in the 69.3% of intervention participants who accessed the site at all. Of studies with multiple, but limited numbers of sessions, Kypri et al offered 2 short, 5- to 10- minute sessions—the initial motivational interview and a short "booster session" 1 month later—in one study<sup>57</sup> and repeated the 10 minute e-intervention at the 1- and 6-month follow-up in another study.<sup>34</sup> Schulz et al<sup>64</sup> provided 3 sessions (length not reported) with the computer program at 3-month intervals. Neighbors et al<sup>35</sup> supplied 2 to 5 computer sessions of 50 minutes each, the number and timing of which was dependent on previous adherence. Barnett et al<sup>48</sup> required a 45-minute session with a 25-minute "booster" session 1 month later to mimic the in-person therapy control.

The other 4 eligible studies included more supplementary human support. Riper and colleagues<sup>63</sup> used a weekly moderated peer support group accessed via the web for 6 weeks after the initial computerized assessment. The length of the group sessions was not reported. The remaining 3 studies were the IVR studies<sup>54,55,60</sup> that included a daily brief check-in call to the program for 2, 3, or 6 months. These check-in calls, averaging 3 to 5 minutes each over 60 to 180 days, totaled 3 to 9 hours of support time. In addition to the IVR time, the IVR studies had either brief initial sessions (<15 minutes) and/or follow-up sessions (40 to 120 minutes).

One study<sup>63</sup> did not report the number or duration of sessions, and the author did not respond to a query for this information.


### **Delivery Mode and Delivery Location: Online, Desktop, IVR, and On- or Off-Site**

Most programs were commercially available (n=17) and accessed online (n=16). The other delivery modes were desktop- or CD-ROM-based programs (n=5), mobile phone application (n=1), or IVR (n=3). Web-delivered interventions were sometimes accessed from dedicated computers in healthcare<sup>49</sup> or research settings, but more often the location was not specified. Desktop-delivered interventions were typically located in a healthcare setting<sup>50,56,62</sup> or research lab.<sup>45</sup> Security features for the programs were not often reported (n=18). In the 3 IVR studies,<sup>54,55,60</sup> the device was linked to the participant's home phone.

### **E-Intervention Content: Theoretical Basis and Most Common Features**

A conceptual framework for the program was reported for 16 of the e-interventions. In most cases (n=13), the e-intervention drew upon principles from motivational interviewing. Cognitive-behavioral therapy (n=4) or the Transtheoretical Model of Behavior Change (n=2) were occasionally used alone or with motivational interviewing.

The most commonly employed strategies used in the e-interventions are summarized by adult and college samples in Figure 2. Overall, PNF was the most common intervention strategy (n=16). PNF consists of a summary of the individual's alcohol intake compared to intake among the subject's peer group. The comparison group was usually gender-specific. In adult studies, they were often age-matched (n=4) or described as local or national population samples (n=3). In student studies, they were described as "student peers." After PNF, the treatment techniques used most often were goal-setting, psychoeducation or alcohol-specific education, and information about the negative consequences of alcohol misuse. Other strategies that were used infrequently included: 1) assessment of attitudes, self-efficacy, and motivation to change; 2) potential strategies to implement change such as decisional balance exercises; 3) relapse prevention strategies; and 4) links to other sources of information. The one mobile application study<sup>52</sup> used techniques specific to this platform— global positioning system (GPS) and texting.

**Figure 2. Strategies Used in E-Interventions**

Abbreviations: A-S=alcohol specific; E-Interventions=electronic interventions; PNF=personalized normative feedback

As described previously, level 1 e-interventions typically consisted of a single session that delivered PNF via computerized feedback or printout. To further illustrate level 2 and 3 interventions, we describe studies in general terms in Table 4.

**Table 4. Examples of Level 2 and Level 3 Interventions (KQ 1)**

**Level 2:** These 8 studies utilized multiple strategies beyond PNF such as the negative consequences and risks associated with alcohol intake, goal-setting, decision-making (including a decisional balance exercise in one case), measurement of motivation to change, correcting misperceptions, and skills training. In 3 cases, subjects completed an assessment that took 30 to 90 minutes.<sup>48,56,62</sup> Five utilized an in-person appointment with a research assistant to conduct parts of the baseline assessment, which is a larger investment on both the part of the subject and the research team.<sup>34,49,56,58,62</sup> Another example of a level 2 intervention used a web-moderated peer group that promoted self-monitoring, skills training, and goal-setting.<sup>63</sup> Finally, a level 2 intervention delivered via mobile phone offered email correspondence with a therapist, an e-bulletin board for posting messages with peers, text-messaging, and a GPS.<sup>52</sup>

**Level 3:** One study<sup>33</sup> used a computer psychoeducation program coupled with a brief intervention delivered by a trained psychologist, followed by 30- to 40-minute phone sessions with the psychologist at 1-, 3-, and 6-month follow-ups. Other level 3 studies<sup>54,55,60</sup> consisted of 1 to 4 face-to-face (n=2) or telephone-delivered (n=1) therapy sessions lasting <30 minutes, combined with daily IVR for 2, 3, or 6 months. In the IVR interventions, therapy was delivered by personnel from various disciplines (study coordinator; MD; BA- or MA-level trainee supervised by a PhD-level psychologist).<sup>54,55,60</sup>

Abbreviations: GPS=global positioning system; KQ=key question; IVR=interactive voice response; PNF=personalized normative feedback

## Summary of Findings

Of 26 studies relevant to KQ 1, only 12 relied on any type of supplementary human support, and only 4 of these included support of a therapeutic nature. This could reflect the fact much of the research to date seems to examine whether a one-time intervention providing data generated from a computer program that compares an individual's alcohol consumption to the peer group norm (PNF) could moderate alcohol consumption in individuals with alcohol misuse. In addition to PNF, goal-setting and psychoeducation/alcohol-specific education including the negative effects and consequences of drinking were the most commonly reported strategies. Although many e-interventions for alcohol misuse are available, few have been evaluated in more than a single study. Most are accessible on the web, but whether or not a user name and password is required is usually not reported. The content of these computer programs varies widely, from one 10-minute session to an unlimited number of sessions that may take up to 45 minutes each to complete. Generally, interventions designed for college students were much less complex, having fewer, shorter sessions and using a more limited number of strategies. Studies in other groups of adults were more intense, containing the studies that used therapeutic support ranging from 1.5 to 5 hours and the studies that targeted subjects with more severe drinking problems.

## KEY QUESTION 2: For adults who misuse alcohol but do not meet diagnostic criteria for AUD, what are the effects of e-interventions compared with inactive controls?

### Key Points

- E-interventions identified were not intensive, typically providing one brief intervention. Existing e-interventions might not have been designed to be robust enough to produce long-term effects ( $\geq 6$  months).
- Included studies generally provided data on alcohol consumption outcomes, but there were limited data on functional outcomes and quality of life.
- The available data suggest that long-term effects of e-interventions on alcohol outcomes are modest or absent.

We identified 22 studies involving 13,929 participants that met eligibility criteria for KQ 2.<sup>33-36,45,49-51,53-59,61-67</sup> Study characteristics are summarized in Table 5; detailed study characteristics are presented in Appendix E. Because one of the studies had 2 e-interventions and one inactive control, allowing 2 comparisons in the same study,<sup>54</sup> there were a total of 23 comparisons between e-interventions and inactive controls. Twelve studies were conducted in adult samples.<sup>33,49-51,53-55,62-65,67</sup> Of these, 4 were judged to be at low risk of bias, 6 moderate risk, and 2 high risk. Another 10 studies were conducted in student samples.<sup>34-36,45,56-59,61,66</sup> Of these, 3 were assessed as being at low risk of bias, 6 moderate risk, and one high risk. Across all studies, there were no differences in baseline alcohol consumption between intervention and control arms (MD 0.32 g/week; 95% confidence interval [CI], -2.87 to 3.50). In this literature, the most common study limitations increasing risk of bias were lack of participant blinding to study condition (demand characteristics) and incomplete or perceived potential for selective reporting of outcome data.



**Table 5. Characteristics of E-Interventions (KQ 2)**

<b>Characteristic</b>	<b>Adult Studies (n=12; 7141 Participants)</b>	<b>Student Studies (n=10; 6788 Participants)</b>
Delivery location/setting:		
Clinic/emergency room:	2	0
College health services	0	2
College classroom	0	1
Research lab	0	1
Home	0	2
NR	10	4
Intervention name:		
escreen	1	0
Drinktest	1	0
Down Your Drink	1	0
Check Your Drinking	1	0
Minderdrinken	1	0
Alcohol-Everything within limits	1	1
Alcohol 101	0	1
e-CHUG	0	2
BASICS	0	2
THRIVE	0	1
What Do You Drink	0	1
College Drinker's	0	1
Not named	6	3
Delivery mode:		
IVR	2	0
Accessed online	8	8
Accessed on desktop computer	1	2
NR	1	0
Number of sessions:		
1	6	8
>1	1	2
NR	2	0
IVR	2	0
Unlimited/unstructured	1	0
Comparator:		
Waitlist	4	5
Attention/information control	6	3
Treatment as usual	2	1
Face-to-face	0	1
Outcomes:		
Alcohol consumption	8	8
HRQOL	2	0
Social/legal effects	1	8
Risk of bias:		
Low	4	3
Moderate	6	6
High	2	1

Abbreviations: BASICS=Brief Alcohol Screening and Intervention for College Students; e-CHUG=Electronic Check-Up to Go; HRQOL=health-related quality of life; IVR=interactive voice response; KQ=key question; NR=not reported; THRIVE=Tertiary Health Research Intervention Via Email

## Results from Studies of Adults


Of the 12 studies conducted in adult samples, 10 provided mean age (range 32.5 to 59.3 years, median 41.1 years). The proportion of females ranged from 0 to 57%, with a median of 41%. Race or ethnicity was reported in only 2 studies. Few data were available on education, though the 7 studies reporting enough data to determine whether participants attended college had a range of 34% to 82% of participants who had attended at least some college. Data on the location of the e-interventions were lacking. Similarly, many studies did not report the name of the e-intervention programs. Of the programs reported in studies reviewed, no specific program was used by more than one study. Most e-interventions were one-session treatments that were accessed online. Half of the studies (n=6) utilized comparison conditions that provided some information or attention as a control, while the other half used waitlist or treatment as usual as comparisons. While 8 studies reported some type of alcohol consumption outcome (grams/week, binge drinking episodes, or meeting alcohol limit guidelines), only 2 studies reported HRQOL, and only one study reported on the social/legal consequences of drinking. We categorized the level of supplementary human support provided as follows: 6 studies had level 1 (minimal) support, 3 studies had level 2 (low) support, and 3 studies had level 3 (moderate or high) support.

To aggregate alcohol consumption across studies, we converted all outcomes to grams (g) of alcohol consumed per week. This generally involved converting standard drinks (U.S., European Union, or Australia) to g/drink, multiplying times the standard drinks/unit time, and then adjusting for unit of time if not given as 1 week. The National Institute on Alcohol Abuse and Alcoholism (NIAAA) defines a U.S. standard drink as a drink that contains 14 g of alcohol, which is often the amount found in a 12-ounce beer, 8 to 9 ounces of malt liquor, 5 ounces of table wine, or 1.5 ounces of hard liquor. The mean baseline alcohol consumption in studies enrolling adult samples ranged from 129 to 436 g/week (median=292 g/wk).

### *Alcohol Consumption*

Figure 3 shows a forest plot of mean differences (MDs) for the 6 studies conducted with adults that had sufficient data at 6-month follow-up assessment to analyze alcohol consumption.<sup>50,51,53,62-64</sup> In these studies, e-interventions were associated with a small, statistically insignificant reduction in alcohol consumption (MD -25.0 g/week; 95% CI, -59.3 to 9.3). Treatment effects varied moderately (Q=11.0; p=0.051; I<sup>2</sup>=55%). Qualitative analyses suggested that studies using more intensive treatment were associated with greater reductions in alcohol consumption. For example, the trial by Riper and colleagues<sup>63</sup> offered access to a moderated peer-to-peer forum, and participants received the recommendation to remain engaged with the e-intervention for 6 weeks. Reductions in alcohol consumption were large. Similarly, Bischof and colleagues<sup>33</sup> observed decreased alcohol consumption in adults at 12-month follow-up in a relatively intensive e-intervention that was combined with 4 phone counseling sessions. A sensitivity analysis that removed the 2 studies at high risk of bias found a small, statistically significant reduction in alcohol consumption (MD -14.7 g/week; 95% CI, -26.4 to -3.0), with little variability across studies (I<sup>2</sup>=0.0%).


Figure 3. Alcohol Consumption at 6 Months in Studies of Adults\*



\*Hansen 2012 and Schulz 2013: Means and SDs were not available, as only mean difference and CI were given. Abbreviations: CI=confidence interval; N=number of participants; NR=not reported; SD=standard deviation

Five studies reported sufficient data to analyze 12-month follow-up assessments of alcohol consumption;<sup>33,51,53,62,67</sup> 3 of the studies were also analyzed at the 6-month time point.<sup>51,53,62</sup> Figure 4 shows a forest plot for these studies. E-interventions were associated with a very small, statistically insignificant reduction in alcohol consumption at 12 months (MD -8.6 g/week; 95% CI, -53.7 to 36.5), with high heterogeneity in treatment effects ( $Q=14.8$ ;  $p=0.005$ ;  $I^2=73\%$ ). A sensitivity analysis that removed the one study at high risk of bias was consistent with the primary analysis (MD -5.5 g/week; 95% CI, -79.0 to 68.1;  $I^2=79.1\%$ ).

Figure 4. Alcohol Consumption at 12 Months in Studies of Adults



Abbreviations: CI=confidence interval; N=number of participants; SD=standard deviation


An additional study with relevance to this question was not included in meta-analyses due to the unique characteristics of its intervention and study sample, namely, face-to-face motivational interviewing plus IVR in individuals with HIV.<sup>54</sup> In this study, a brief motivational interviewing (BMI) intervention was paired with a daily IVR system for tracking drinking behavior. Another arm of the study employed BMI only and is described under KQ 4. Relative to inactive control, the BMI plus IVR arm resulted in similar amounts of alcohol consumption at 6- and 12-month follow-up assessments.

### Alcohol Reduction to Meet Drinking Limit Guidelines

The operational definition of drinking limit guidelines varied across studies. Studies typically defined limits by a weekly total of standard drinks (eg, <20).<sup>49,63</sup> However, one study<sup>62</sup> combined the weekly limit with a daily limit ( $\leq 5$  drinks for men,  $\leq 3$  for women), whereby exceeding either the weekly or the daily limit would count as not meeting the guideline. The studies varied considerably in the proportion of participants meeting drinking limit guidelines at the 6-month follow-up. The 4 studies with data on this outcome reported 17%, 23%, 46%, and 78% meeting drinking limit guidelines in the e-treatment groups at 6 months.

Figure 5 shows a forest plot of risk ratios (RRs) for the 4 studies conducted in adults that had sufficient data at 6-month follow-up assessment to analyze the proportion of participants meeting drinking limit guidelines.<sup>49,62-64</sup> In these studies, e-interventions were associated with a small, statistically insignificant increase in the risk of meeting guidelines (RR 1.22; 95% CI, 0.79 to 1.89). There was low to moderate heterogeneity in effect sizes ( $Q=6.5$ ;  $p=0.088$ ;  $I^2=54\%$ ).

**Figure 5. Alcohol Reduction to Meet Drinking Limit Guidelines at 6 Months in Studies of Adults\***



\*Schulz 2013 did not report event rates. Intervention effects are based on adjusted estimates reported from a logistic regression model.

Abbreviations: CI=confidence interval; N=number of participants; NR=not reported; SD=standard deviation

There were no studies with sufficient data to analyze 12-month follow-up assessments of meeting drinking limit guidelines.

An additional study targeted alcohol misuse with comorbid illicit drug use.<sup>65</sup> This study enrolled participants online and provided a brief online intervention where participants could monitor drinking and get PNF with motivational interviewing-based recommendations. This was compared to an assessment-only condition. Drinking limits in this study were defined as AUDIT <8. At 6-month follow-up, the 2 groups had similar proportions of participants with AUDIT <8.

### Binge Drinking

Two studies in adults reported the effects of e-treatment on binge drinking.<sup>50,53</sup> Both trials reported 6-month data, but no 12-month data. Due to the limited number of trials, we did not conduct meta-analysis. One study enrolled military Veterans with alcohol misuse and compared treatment as usual in a primary care setting to treatment as usual plus a brief online alcohol intervention.<sup>50</sup> Binge drinking was defined as  $\geq 5$  drinks on one occasion for men, and  $\geq 4$  drinks on one occasion for women in the past 30 days. At the 6-month follow-up, participants in the

brief intervention condition had rates of binge drinking (23%) similar to those in the treatment-as-usual condition (25%). The other study reporting binge drinking outcomes in adults compared an online PNF intervention to a no-intervention control group.<sup>53</sup> Binge drinking was defined as drinking  $\geq 5$  drinks on one occasion. At the 6-month follow-up, the binge drinking rate in the online PNF condition (47%) was similar to that of the non-intervention control condition (45%).

### *Consequences of Drinking—Social Problems*

One trial, conducted in Veterans, reported intervention effects on the negative social consequences of drinking at 6-month follow-up.<sup>50</sup> This study is also described immediately above, under “Binge Drinking.” This study measured negative social consequences of drinking with the Short Inventory of Problems, a 15-item measure on which higher scores indicate more social problems. In this study, the brief online alcohol intervention had a mean score on the self-report measure of social problems (mean 5.9, SD 10.2) that was similar to that observed in the treatment-as-usual-only group (mean 6.5, SD 9.3). No 12-month data were available.

### *Other Outcomes*

In addition to the outcomes described above, we examined trials for analyses of HRQOL, alcohol-related health problems, medical utilization, and adverse effects from treatment. No trials with adults reported these outcomes with sufficient data to analyze.

## **Results from Studies of College Students**


Ten trials were conducted in college student samples.<sup>34-36,45,56-59,61,66</sup> Due to theorized potential differences in sample characteristics and responsiveness to treatments, we analyzed these samples separately from the adult samples. Of the 10 trials conducted in college student samples, 8 provided mean age (median of means 19.9 years, range 18.2 to 20.9 years). The proportion of females ranged from 38% to 64%, with a median of 55%. In the 4 trials reporting race/ethnicity, the proportion of participants of non-Caucasian race/ethnicity ranged from 15% to 43%, with a median of 28%. The location of the e-interventions varied considerably, as the 6 studies reporting location included college health services, college classrooms, research labs, and home-based e-interventions. The programs used also varied, and no one program was used in more than 2 studies. E-interventions were primarily delivered online (n=8 studies). Treatment was not intensive, as the modal number of e-intervention sessions was 1 (n=8 studies). Half of the studies used waitlist comparison groups, but more active comparison groups were included, as 3 studies used attention/information comparisons, and one used face-to-face therapy as a comparison. A total of 9 studies provided data on alcohol consumption (either g/week, binge drinking episodes, or meeting alcohol limit guidelines). Eight studies provided data on negative social and/or legal consequences of drinking. No data on HRQOL were provided. Regarding the level of supplementary human support provided, 7 trials had minimal support, and 3 trials had low support. Of the 8 trials that provided baseline alcohol consumption data, baseline means were lower than in adult studies, ranging from 85 g/week to 291 g/week (median 183 g/week).

### *Alcohol Consumption*

Figure 6 shows a forest plot of MDs for the 8 studies conducted in college students that had sufficient data at 6-month follow-up assessment to analyze alcohol consumption.<sup>34-36,45,57,58,61,66</sup> In

these studies, e-interventions were associated with a small, statistically insignificant reduction in alcohol consumption (MD -12.4 g/week; 95% CI, -26.6 to 1.9). There was moderate heterogeneity in treatment effects ( $Q=11.8$ ;  $p=0.11$ ;  $I^2=41\%$ ).


**Figure 6. Alcohol Consumption at 6 Months in Studies of College Students**



Abbreviations: CI=confidence interval; N=number of participants; SD=standard deviation

There were also 3 studies with sufficient data to analyze 12-month follow-up assessments of alcohol consumption.<sup>34,35,56</sup> A forest plot for these studies is shown in Figure 7. In these studies, e-interventions were associated with a small, statistically insignificant reduction in alcohol consumption. (MD -22.2 g/week; 95% CI, -96.6 to 52.3). There was substantial heterogeneity in treatment effects ( $Q=7.5$ ;  $p=0.024$ ;  $I^2=73\%$ ).

**Figure 7. Alcohol Consumption at 12 Months in Studies of College Students**



Abbreviations: CI=confidence interval; N=number of participants; SD=standard deviation

**Alcohol Reduction to Meet Drinking Limits**


One study with college students had data on 6-month outcomes for meeting drinking limit guidelines.<sup>57</sup> In this study, drinking guidelines were defined as a limit of 14 drinks per week for women and 28 drinks per week for men. The e-intervention consisted of psychoeducation about

negative consequences of current alcohol consumption level, including health, financial, and motor vehicle accident risks. In addition, the e-intervention provided normative comparisons and information on treatment resources. Relative to an assessment-only comparison group, the e-intervention resulted in an elevated probability of meeting drinking limits at 6 months (OR 1.53; 95% CI, 1.09 to 2.17).

### Binge Drinking

Figure 8 shows a forest plot of MDs for the 4 trials conducted with college students that had sufficient data at 6-month follow-up assessment to analyze effects of e-interventions on binge drinking episodes.<sup>34,35,45,58</sup> In these studies, e-interventions were associated with a very small, statistically insignificant reduction in binge drinking (MD -0.1 episodes; 95% CI, -1.0 to 0.9), with low to moderate evidence of heterogeneity in effect sizes ( $Q=5.5$ ;  $p=0.14$ ;  $I^2=46\%$ ).

**Figure 8. Episodes of Binge Drinking at 6 Months in Studies of College Students**



Abbreviations: CI=confidence interval; N=number of participants; SD=standard deviation

There were no trials with sufficient data to analyze 12-month follow-up assessments of binge drinking episodes.


Two additional trials reported the proportion of subjects with binge drinking, but these outcomes could not be combined with the mean episodes of binge drinking and thus these studies were excluded from the meta-analysis.<sup>57,66</sup> One study compared a brief online e-intervention to a no-intervention comparison group.<sup>66</sup> At the 6-month follow-up, the proportion of participants reporting binge drinking in the e-intervention group (68%) was similar to that in the control group (66%). Another trial compared a brief online e-intervention to an informational leaflet intervention.<sup>57</sup> Binge drinking was defined as >120 g for males and >80 g for females during one episode in the last 2 weeks. The median number of binge drinking episodes was the same for both groups (median=1). These findings of no intervention effect on episodes of binge drinking are consistent with the finding of no effect on mean number of binge drinking episodes for the studies included in the meta-analysis.

### Consequences of Drinking—Social Problems

Figure 9 shows a forest plot of SMDs for the 7 trials conducted in college students that had sufficient data at 6-month follow-up assessment to analyze the effects of e-interventions on

negative social consequences of drinking.<sup>34-36,57-59,61</sup> In these trials, e-interventions were associated with a very small, statistically insignificant reduction in the negative social consequences of drinking (SMD -0.04; 95% CI, -0.22 to 0.13), with moderate heterogeneity in treatment effects (Q=14.1; p=0.029; I<sup>2</sup>=57%).


**Figure 9. Negative Social Consequences of Drinking at 6 Months in Studies of College Students**



Abbreviations: CI=confidence interval; N=number of participants; SD=standard deviation; SMD=standardized mean difference

Figure 10 shows a forest plot of SMDs for the 4 trials with sufficient data at 12-month follow-up for assessment of negative social consequences of drinking.<sup>34,35,56,59</sup> Similar to results observed in the 6-month data, e-interventions were associated with a very small, statistically insignificant reduction in negative social consequences of drinking at 12-month follow-up (SMD -0.03; 95% CI, -0.59 to 0.54); there was substantial heterogeneity in treatment effects (Q=26.4; p<0.001; I<sup>2</sup>=89%). A sensitivity analysis that removed the one study at high risk of bias was consistent with the primary analysis (MD -0.12 g/week; 95% CI, -0.98 to 0.74; I<sup>2</sup>=90.9%).

**Figure 10. Negative Social Consequences of Drinking at 12 Months in Studies of College Students\***



\*Hester 2012 did not report mean and SD. The estimate of treatment effect was derived from other statistics. Abbreviations: CI=confidence interval; N=number of participants; SD=standard deviation; SMD=standardized mean difference



### *Other Outcomes*

No trials in college students reported HRQOL, alcohol-related health problems, medical utilization, or adverse effects from treatment with sufficient data to analyze.

### **Summary of Findings**

We reviewed a total of 22 studies that compared e-interventions with inactive controls in participants with varying degrees of alcohol misuse. Most interventions were accessed online, used a single session, and had minimal to low levels of supplementary human support. While most studies provided data on alcohol consumption outcomes, other important outcomes were reported infrequently. Treatment effects on weekly alcohol consumption were relatively small and statistically insignificant, and varied importantly across trials. Fewer studies provided data on meeting drinking limit guidelines at the end of the study, reducing binge-drinking episodes, and decreasing social problems related to alcohol. Moderately strong evidence showed no effects on binge drinking or alcohol-related social problems, but these outcomes were reported primarily in student samples. The strength of the current evidence for the longer-term ( $\geq 6$  months) benefits of e-interventions on alcohol consumption and its associated effects on health and well-being is low.

### **Quality of Evidence for KQ 2**

Of the 22 trials relevant to this question, 7 were judged to be at low risk of bias, 12 at moderate risk of bias, and 3 at high risk of bias (Appendix C). We used the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach to rate the overall strength of evidence (SOE) for outcomes that addressed alcohol consumption and social problems related to alcohol use. For 2 of the alcohol consumption outcomes (weekly consumption and meeting alcohol limits), we judged the SOE as low. The SOE was downgraded for inconsistent and imprecise treatment effects and, for the weekly consumption outcome, moderate risk of bias. Moderately strong evidence showed no effects on binge drinking or alcohol-related social problems, but these outcomes were reported primarily in student samples. These outcomes were downgraded for some important inconsistency in treatment effects and indirectness, since most of the studies involved student samples with lower baseline alcohol consumption than the adult samples.

### **KEY QUESTION 3: For adults at high risk of AUD (eg, audit-C $\geq 8$ ), or who have a diagnosis of AUD, what are the effects of e-interventions compared with inactive controls?**

#### **Key Points**

- Only 3 studies compared e-interventions with no-intervention controls in participants with AUD (n=533).
- In a single trial, a multi-component smartphone program used to support recovery following residential treatment increased abstinence (OR 1.94; 95% CI, 1.14 to 3.31) and decreased risky drinking days at 12-month follow-up (SOE=low).
- A computerized feedback system coupled with telephonic counseling in participants recruited from primary care did not reduce alcohol consumption. An IVR system used

for participants discharged from residential treatment did not decrease the risk of relapse (SOE=insufficient).

### **Study Characteristics and Treatment Effects**

We identified 3 trials (Table 6) involving 533 participants that met inclusion criteria for KQ 3.<sup>33,52,60</sup> All studies were conducted in adult, non-student samples and were judged to be at moderate<sup>33,52</sup> or high<sup>60</sup> risk of bias (Appendix C). None of the e-interventions was identified as a commercial product. Because these studies varied considerably in participants enrolled, interventions, and outcomes reported, we describe each separately below.

Table 6. Study Characteristics (KQ 3)

Study	Setting	Mean Age (SD)	N	Inclusion Criteria	E-Intervention	Control	Follow-up Interval	Outcomes	Overall Risk of Bias
Bischof, 2008 <sup>33</sup>	Primary care	36.5 (13.5)	408 (124 AUD subgroup)	AUDIT $\geq 5$ and $\geq 2$ binge episodes in past 4 weeks or $>20$ g (F) or 30 g (M) daily	Computerized feedback on alcohol use and 1-4 telephonic MI sessions	Waitlist	12 months	Alcohol consumption	Moderate
Gustafson, 2014 <sup>52</sup>	Residential care facilities	38.4 (10.4)	349	Completing residential treatment program for DSM-IV alcohol dependence	Multicomponent smartphone application offered for 8 months	Treatment as usual	12 months	Alcohol consumption Social effects	Moderate
Mundt, 2006 <sup>60</sup>	Residential facility	41.9 (9.2)	60	Completing residential treatment program for AUD	IVR prompted or ad lib	No IVR	6 months	Alcohol consumption HRQOL Social effects	High

Abbreviations: AUD=alcohol use disorder; AUDIT=Alcohol Use Disorders Identification Test; DSM-IV=*Diagnostic and Statistical Manual of Mental Disorders*, 4<sup>th</sup> edition; E-Intervention=electronic intervention; F=female; HRQOL=health-related quality of life; IVR=interactive voice response; KQ=key question; M=male; MI=motivational interviewing; N=number of participants; SD=standard deviation

Gustafson and colleagues<sup>52</sup> recruited 349 participants from 5 U.S. residential treatment facilities who were completing treatment for alcohol dependence. The average age was 38.4 (SD 10.4) years old, 39% were women, 80% were white, and mean years of education were not reported. At discharge from residential treatment, participants were randomized to: 1) the multicomponent Addiction-Comprehensive Health Enhancement Support System (A-CHESS) for 8 months; or 2) treatment as usual. Intervention participants received a smartphone with the A-CHESS application, which included both static and interactive features, mobile phone service, and a data plan. Two hundred and eighty-six (286) smartphones were given to the 170 intervention participants due to malfunctioning, lost, stolen, and damaged phones. Participants could elect to share data from the A-CHESS application with their counselors: 93.5% shared data from the weekly Brief Addiction Monitor, and 41.9% shared data about alcohol relapse. At 12-month follow-up, the intervention group had higher rates of abstinence (OR 1.94; 95% CI, 1.14 to 3.31) and fewer risky drinking days per 30-day period (MD -1.47; 95% CI, -0.13 to -2.81). There were no significant effects on any of the 8 items measuring negative consequences of drinking.

Bischof and colleagues<sup>33</sup> recruited 408 participants from German primary care practices with alcohol problems ranging from a positive alcohol screen to alcohol dependence established by a criterion-based structured interview. Average age was 36.5 (SD 13.5) years old, 32% were women, race was not reported, and mean years of education were 10.5 (2.5). We report on the subgroup (n=124) with alcohol dependence. Participants were randomized to: 1) a no-intervention control; 2) computerized feedback plus 4 brief counseling sessions of 30 minutes each delivered by telephone; or 3) computerized feedback and 1 to 4 brief counseling sessions of 30 minutes each delivered by telephone. For the latter group, counseling sessions were discontinued when the patient met criteria for a reduction in alcohol consumption and indicated high self-efficacy to maintain the reduction in alcohol consumption. The computerized feedback was based on the transtheoretical model of behavior change, and telephonic counseling was based on motivational interviewing and delivered by a trained psychologist. At 12-month follow-up, the intervention groups did not differ from the control group in alcohol consumption (p=0.62) or in the proportion meeting criteria for binge drinking (54.5% intervention vs 50% control; p=0.69). Other outcomes prioritized for this review were not reported.

Mundt et al<sup>60</sup> evaluated the effect of an IVR system in 60 participants treated for alcohol dependence at a Wisconsin residential treatment facility. Average age was 41.9 (SD 9.1) years, 45% were women, 95% were white race, and mean years of education were not reported. At discharge from residential treatment, participants were randomized to: 1) a no-intervention control; 2) prompted use of the IVR system; or 3) ad libitum use of the IVR system. The prompted group was called and encouraged to use the system if they failed to make a daily call to the IVR system for 2 consecutive days. The IVR system included assessments of drinking status, alcohol cravings, self-efficacy, difficulty coping, and risk of relapse. Participants were directed to modules that incorporated tailored feedback based on patterns of alcohol consumption: resolve not to drink module, drinking reasons module, and cognitive behavioral advice module. The system included voice mail capability and allowed participants to request a personal phone call from the coordinator. At 6-month follow-up, intervention subjects had placed a mean of 51.4 (SD 22.6) IVR calls; mean calls were higher (p=not statistically significant [NS]) in the ad

libitum group. Self-reported abstinence was reported in 20 of 30 intervention subjects (67%) and 13 of 18 control subjects (72%;  $p$ =not reported [NR]). There were no statistically significant intervention effects on functional status, or work and social adjustment.

### Summary of Findings

Three trials conducted in participants with alcohol use disorders found positive effects from a multicomponent smartphone application, but no intervention effects from a computerized feedback system coupled with telephonic counseling or an IVR system.

### Quality of Evidence for KQ 3

The 3 trials were judged to be at moderate<sup>33,52</sup> or high<sup>60</sup> risk of bias (Appendix C). Using the GRADE framework,<sup>69</sup> the SOE is low (moderate risk of bias, imprecise effects) that a multicomponent smartphone intervention decreases alcohol use in adults with alcohol use disorders. The SOE is insufficient for the effects of IVR and computerized feedback couple with brief telephone counseling.

## **KEY QUESTION 4: For adults who misuse alcohol, are at high risk of AUD, or have a diagnosis of AUD, what are the effects of e-interventions alone or used in combination with face-to-face therapy compared with face-to-face therapy alone?**

### Key Points

- Six trials (1090 participants) compared e-interventions alone or in combination with face-to-face brief motivational interviewing (BMI) to BMI alone. All studies enrolled individuals with alcohol misuse, but varied markedly with regard to setting and subject and intervention characteristics.
- Combination of e-interventions plus BMI versus BMI alone: Two studies in adults compared BMI plus IVR to BMI alone and found no improvement in primary drinking outcomes with the addition of IVR. A subgroup analysis found benefit with combined IVR and BMI for HIV-positive adults who met DSM-IV criteria for alcohol dependence. In a study of college students, the addition of an e-intervention utilizing 1 to 2 sessions of personalized normative feedback (PNF) plus psychoeducation to BMI versus BMI alone showed no improvement in 12-month drinking outcomes.
- E-interventions alone versus BMI alone: In 3 head-to-head comparisons, all involving college students, e-interventions were associated with significantly heavier alcohol intake at follow-up (ranging from 50 to 81 g greater weekly consumption) compared with BMI. Binge drinking frequency was higher in e-intervention participants than in BMI participants (MD 2 to 2.5 more episodes per month).
- This diverse group of studies did not find a benefit of e-interventions alone or as an adjunct to face-to-face BMI, compared with face-to-face BMI alone for college students or midlife primary care patients who misuse alcohol.

## Study Characteristics

Six trials (Table 7), described in 5 papers and involving 1090 participants, compared e-interventions alone or used in combination with face-to-face BMI to face-to-face BMI alone.<sup>36,45,48,54,55</sup> The paper by Monahan and colleagues<sup>45</sup> reported results from 2 separate trials. Two trials (508 participants), conducted with midlife patients in primary care settings, compared the combination of IVR plus face-to-face BMI to face-to-face BMI alone.<sup>54,55</sup> Four trials (582 participants) focused on college students in their late teens.<sup>36,45,48</sup> Three of these compared e-interventions to BMI.<sup>36,45</sup> The fourth compared the combination of an e-intervention plus BMI to BMI alone.<sup>48</sup> All studies enrolled subjects based on recent alcohol misuse (*eg*,  $\geq 1$  episode of heavy drinking in the past 14 or 30 days, acute intoxication, or alcohol policy violation) rather than specific evidence of a diagnosable AUD. Thus, subject inclusion criteria in these studies are similar to those used in the studies described in KQ 2. One study used DSM-IV criteria to assess enrolled subjects, allowing subgroup analysis for those with a diagnosis of alcohol dependence.<sup>54</sup> Two studies<sup>48,54</sup> were judged to be at low risk of bias, and 4 studies were assessed as being at moderate risk of bias.<sup>36,45,55</sup>

Table 7. Study Characteristics (KQ 4)

Author	Setting	Mean Age	N	Inclusion Criteria (M/F)	e-Intervention	Control	Follow-up Interval	Outcomes	Overall Risk of Bias
Barnett 2007 <sup>48</sup>	College health service	18.8	225	Referred due to intoxication or violation	Alcohol 101 + BMI	BMI ± 25-min booster	12 months	Alcohol consumption Social effects	Low
Hasin, 2013 <sup>54</sup>	HIV primary care	46	170	>3 drinks once in 30 days	IVR daily x 60 days + BMI	BMI, 20-25 min	12 months	Alcohol consumption	Low
Helzer, 2008 <sup>55</sup>	Primary care	45	338	>NIAAA drinking limits or CAGE 1+	IVR daily x 6 months + BMI ± PNF	BMI ± mailed PNF	6 months	Alcohol consumption	Moderate
Monahan, 2013a=Study 1 <sup>45</sup>	College health service	21.2	73	>4/3 drinks once in past 30 days	Alcohol 101	BMI 1 x 50-min session	6 months	Alcohol consumption	Moderate
Monahan, 2013b=Study 2 <sup>45</sup>	College course	18.6	91	>4/3 drinks once in past 30 days	e-CHUG	BMI	6 months	Alcohol consumption	Moderate
Walters, 2009 <sup>36</sup>	College campus	19.8	193	>4/3 drinks once in past 14 days	e-CHUG	BMI ± PNF	6 months	Alcohol consumption Social effects	Moderate

Abbreviations: BMI=brief motivational interviewing; CAGE=Cut down/Annoyed/Guilty/Eye-opener; e-CHUG=Electronic Check-Up to Go; F=female; IVR=interactive voice response; KQ=key question; M=male; min=minute(s); NIAA=National Institutes of Alcoholism and Alcohol Abuse; PNF=personalized normative feedback

## E-Intervention Characteristics

E-interventions studied were daily IVR calls for periods of 60 days<sup>54</sup> or 6 months,<sup>55</sup> Alcohol 101 interactive CD-ROM;<sup>45,48</sup> and e-CHUG, a web-based interactive program.<sup>36,45</sup> Comparator face-to-face counseling consisted of single-session of BMI; 2 studies included a brief booster or follow-up counseling sessions. The extent of counselor experience, training, supervision, and fidelity monitoring varied across the 6 studies. Characteristics of the e-interventions evaluated are summarized in Table 8.

**Table 8. Characteristics of E-Interventions (KQ 4)**

Characteristic	Adult studies (n=2; 508 participants)	Student studies (n=4; 582 participants)
Delivery location:		
Primary care	2	0
College health services	0	2
College campus	0	2
Intervention design:		
BMI plus e-intervention	2	1
e-intervention alone	0	3
Intervention name:		
Alcohol 101	0	2
e-CHUG	0	2
Not named	2	0
Delivery Mode:		
IVR	2	0
Accessed online	0	2
Accessed on desktop computer	0	2
Number of e-intervention sessions:		
1	0	3
>1	2 (IVR daily)	1
In-person BMI sessions:		
1	1	3
>1	1	1
Outcomes:		
Alcohol consumption	2	4
HRQOL	0	0
Social/legal effects	0	2
Risk of bias:		
Low	1	1
Moderate	1	3

Abbreviations: BMI=brief motivational interviewing; e-CHUG=Electronic Check-Up to Go; HRQOL=health-related quality of life; IVR=interactive voice response; KQ=key question

## Alcohol Consumption

### *E-intervention Alone versus Face-to-Face Counseling (3 Trials)*

E-interventions were compared with face-to-face interventions only in college student populations. In all 3 head-to-head comparisons of e-interventions versus face-to-face



BMI, e-interventions were associated with poorer alcohol outcomes than was face-to-face counseling.<sup>36,45</sup>

Monahan et al<sup>45</sup> reported 2 trials comparing e-interventions to BMI in college students recruited in a campus health service. The entry criterion for both studies was at least one episode of consuming >5 drinks (males) or >4 drinks (females) in the past month. In Study 1, 60 subjects were randomized to the Alcohol 101 CD-ROM or face-to-face BMI. In Study 2, 68 subjects were randomized to an interactive, web-based program that included personalized feedback (e-CHUG) or face-to-face BMI. At 6-month follow-up in Study 1, participants receiving the e-intervention were consuming more alcohol per week than those assigned to face-to-face BMI, but results were statistically insignificant (MD 52.2 g/week; 95% CI, -35.7 to 140.1). A similar difference was found in Study 2 (MD 50.1 g/week; 95% CI, -52.6 to 152.8). In both studies, the frequency of binge drinking at 6 months was higher in those receiving the e-intervention, but results were statistically significant only in Study 2. Study 1 e-intervention participants reported 5.5 (SD 6.7) past-month binge episodes versus 3.5 (SD 3.3) for BMI participants (MD 2.0; 95% CI, -0.57 to 4.7). In Study 2, e-intervention participants reported 5.8 (SD 5.5) past-month binge episodes versus 3.3 (SD 4.9) with BMI (MD 2.5; 95% CI, 0.14 to 5.01).

Walters et al<sup>36</sup> studied college students (average age 19.8 years) reporting at least one episode of consuming  $\geq 5$  drinks (males) or  $\geq 4$  drinks (females) in the past 2 weeks. Two hundred and seventy-nine subjects were randomized to one of 4 conditions: 1) assessment only; 2) e-CHUG; 3) BMI without structured assessment feedback; 4) BMI that incorporated PNF derived from a structured assessment. The assessment-only group is not relevant to this KQ. At 6-month follow-up, e-intervention participants were consuming 168.9 g of alcohol per week (SD 172.3) versus 162.3 (SD 295.9) for BMI only and 142.7 (SD 133.7) for BMI with feedback. In an analysis that adjusted for differences in baseline values, participants assigned to the e-intervention reported significantly higher alcohol consumption compared with those receiving BMI with feedback (5.79 drinks per week, standard error of the mean [SEM] 2.05,  $p=0.005$ ).

### *E-intervention Plus Face-to-Face Counseling versus Face-to-Face Counseling Alone (3 Trials)*

In the 3 trials comparing e-interventions plus face-to-face counseling versus face-to-face counseling alone, the addition of an e-intervention was not associated with decreased alcohol consumption.<sup>48,54,55</sup> Two studies enrolled midlife adult patients recruited from primary care settings,<sup>54,55</sup> and one recruited participants from a college health service.<sup>48</sup>

Helzer et al<sup>55</sup> studied adult primary care patients (mean age 45) whose average alcohol consumption exceeded NIAAA recommended drinking limits or who scored >1 positive on the Cut down/Annoyed/Guilty/Eye-opener (CAGE) questionnaire. The study was conducted across 15 primary care clinics, involving 112 providers who were trained in conducting brief intervention for excessive alcohol use. All potential subjects received a face-to-face brief intervention delivered by their primary care provider and were then invited to participate in the study, in which 338 subjects were randomized to: 1) face-to-face brief intervention only; 2) face-to-face brief intervention plus 6 months of daily IVR follow-up; or 3) face-to-face brief intervention plus 6 months of daily IVR follow-up plus feedback. Feedback consisted of a mailed, printed graph showing daily consumption as reported via IVR in comparison with

the participant's stated drinking goal, accompanied by a personalized note from Dr. Helzer highlighting the importance of the information. Two hundred and seventy-three subjects (81%) completed the 6-month follow-up interview, with no significant differences in follow-up rate between treatment groups. At 6-month follow-up, mean weekly drinking levels were 25.0 standard drinks (SEM 1.3) in subjects randomized to IVR; 22.4 (SEM 1.4) standard drinks in subjects assigned to IVR plus feedback; and 18.3 (SEM 1.3) for those assigned to brief intervention alone. In a preplanned comparison of all IVR participants (combining those with and without feedback) versus brief intervention-only participants, IVR was associated with significantly heavier weekly drinking (22.6 drinks per week, SEM 0.8) compared with brief intervention only (18.3 drinks, SEM 1.3;  $p=0.01$ ).

Hasin et al<sup>54</sup> compared BMI plus IVR to BMI alone in a primary care clinic serving HIV-positive individuals (mean age 46). The inclusion criterion was consumption of >4 standard drinks on one occasion within the past 30 days. One-hundred and seventy subjects were randomized to a 20- to 25-minute BMI session only versus BMI plus daily IVR for 60 days (BMI+IVR). At 12-month follow-up, there was no difference between groups for the primary outcome measure, number of drinks per drinking day. In subgroup analysis, subjects who met criteria for a DSM-IV diagnosis of alcohol dependence, drinks per drinking day were lower in the BMI+IVR than the BMI-only group ( $p<0.03$ ).

Barnett et al<sup>48</sup> compared the combination of BMI plus the Alcohol 101 CD-ROM program with BMI alone in college students ( $n=225$ ) who were referred to campus health service due to an episode of acute intoxication or violation of campus alcohol policy. At 12-month follow-up, there were no between-group differences in alcohol consumption or alcohol related social or legal problems.

### Other Outcomes

Other outcomes were reported infrequently. Two studies conducted in college students reported the effects of e-interventions on social outcomes at 6<sup>36</sup> or 12 months.<sup>48</sup> Neither study found a differential effect between BMI and the e-intervention for this outcome. No study reported effects on HRQOL, alcohol-related health problems, medical utilization, or adverse effects of treatment.

### Summary of Findings

Six trials (1090 participants) compared e-interventions alone or in combination with face-to-face BMI to BMI alone. Studies were too diverse to compute summary estimates of effect. IVR in combination with BMI was the only e-intervention compared to face-to-face treatment in non-collegiate populations. The addition of IVR did not decrease alcohol consumption in 2 trials that enrolled midlife primary care patients with alcohol misuse. In collegiate populations, BMI was generally more effective than e-interventions alone in decreasing alcohol consumption.

### Quality of Evidence for KQ 4

Two trials were judged to be at low risk of bias and 4 at moderate risk (Appendix C). For e-interventions compared with face-to-face treatment, we judged the SOE as low. These studies had moderate risk of bias, were conducted in collegiate populations, and effect estimates did

not exclude a difference of 3 standard drinks per week or a SMD of  $\leq 0.4$ , the thresholds we set for a precise estimate of effect. When evaluating the overall SOE, we considered a difference of 3 standard U.S. drinks/week or an SMD  $\geq 0.4$  as clinically significant and defined precise effects as those with 95% CIs that excluded smaller effects. For combined e-interventions plus face-to-face treatment versus face-to-face alone, we judged the SOE to be low for effects on alcohol consumption, and insufficient for effects on social and legal outcomes. These studies had moderate risk of bias and primarily used IVR interventions, which have important differences from other e-interventions because the interface is through a telephone or analog line.

## SUMMARY AND DISCUSSION

Substance use disorders, including alcohol use disorder (AUD), are among the most common and costly conditions among Veterans presenting for treatment in the Veterans Health Administration (VHA) system.<sup>12,13</sup> A recent systematic review<sup>8</sup> found moderately strong evidence that in-person, multi-contact, brief behavioral interventions decreased alcohol consumption in primary care patients who screened positive for alcohol misuse. Electronic interventions (e-interventions) for alcohol misuse have been proposed as a useful way to extend the reach of initial intervention and relapse prevention to maintain treatment gains for alcohol misuse.

We identified 26 randomized controlled trials (RCTs) involving 14,497 total participants that were relevant to our key questions (KQs). Participants were selected for these trials based on one or more alcohol consumption criteria, but only 3 studies based inclusion on an assessment of AUD. Most trials compared e-interventions with inactive controls. E-interventions were typically accessed online and consisted of one session completed without supplementary human support; personalized normative feedback (PNF) was the predominant strategy. Studies and participants were roughly equally divided between college students and other groups of adults. Adult participants were typically midlife, the majority of whom had at least some college education, with baseline alcohol consumption in excess of 14 drinks per week.

We assessed the strength of evidence (SOE) for the highest priority outcomes—alcohol consumption and alcohol-related social problems. Overall, there was low SOE that e-interventions compared with inactive controls did not decrease alcohol consumption outcomes in participants with alcohol misuse. In patients with AUD, a multicomponent smartphone application decreased the risk of relapse after residential treatment (SOE=low). In the few studies using face-to-face treatment as the comparator, e-interventions alone or in combination with face-to-face treatment were not associated with decreased alcohol use, but these studies were conducted exclusively in student populations. Interactive voice response (IVR) e-interventions may be less effective than face-to-face treatment. Table 9 details SOE ratings for the highest priority outcomes by KQ.

Table 9. Detailed SOE Ratings

Outcome	SOE Domains				Effect Estimate (95% CI)	SOE
	Number of Studies (Participants)*	Study Design/ Risk of Bias	Consistency Directness	Precision Publication Bias		
<b>KQ 2: E-intervention vs control in individuals who screen positive for alcohol misuse</b>						
Alcohol consumption (weekly)	17 (10,122)	RCT/Moderate	Inconsistent Direct	Imprecise None detected	MD -25.0 g/week (-59.3 to 9.3) Adults, n=6 studies  MD -12.4 g/week (-26.6 to 1.9) Students, n=8 studies	Low Low
Met alcohol consumption limits	6 (4932)	RCT/Low	Some inconsistency Direct	Imprecise None detected	RR 1.22 (0.79 to 1.89) Adults, n=4 studies  OR 1.53 (1.09 to 2.17) Students, n=1 study	Low Low
Alcohol consumption (binge drinking)	8 (5043)	RCT/Low	Some inconsistency Some indirectness	Precise None detected	MD 2% with binge Adults, n=2 studies  MD -0.01 episodes (-1.0 to 0.9) Students, n=4 studies	Moderate Moderate
Alcohol-related social problems	8 (5765)	RCT/Low	Some inconsistency Some indirectness	Precise None detected	No difference Adults, n=1 study  SMD -0.04 (-0.22 to 0.13) Students, n=7 studies	Low Moderate

Outcome	SOE Domains				Effect Estimate (95% CI)	SOE
	Number of Studies (Participants)*	Study Design/ Risk of Bias	Consistency Directness	Precision Publication Bias		
<b>KQ 3: E-intervention vs control in individuals with AUD</b>						
Alcohol consumption (maintain abstinence)	3 (533)	RCT/Moderate	Consistent Direct	Imprecise None detected	Increase in abstinence for adults with smartphone e-intervention: OR 1.94 (1.14 to 3.31)  No difference with IVR or computerized feedback	Low Insufficient
Alcohol-related social problems	2 (409)	RCT/Moderate	NA Direct	Imprecise None detected	No difference in adults	Low
<b>KQ 4: E-intervention vs face-to-face counseling</b>						
Alcohol consumption†	3 (438)	RCT/Moderate	Consistent Some indirectness	Imprecise None detected	Ranged from 50 to 80 g/week higher with e-intervention Students, n=3 studies	Low
Alcohol-related social problems	1 (210)	RCT/Moderate	NA Some indirectness	Imprecise None detected	SMD -0.16 (-0.76 to 0.43) Students, n=1 study	Insufficient
<b>KQ 4: Combined e-intervention + face-to-face counseling vs face-to-face counseling alone</b>						
Alcohol consumption	3 (668)	RCT/ Moderate	Consistent Some indirectness	Imprecise None detected	Adults: No difference in 1 study; higher in 1 study  Students: No difference in 1 study	Low Low
Alcohol-related social problems	0	NA	NA NA	NA NA	No studies	Insufficient

\*Numbers given here are for the total number studies/participants considered in the SOE ratings. Numbers included in the summary effect estimates may vary.

†One standard drink in the U.S contains 14 grams of alcohol (eg, one 12-ounce beer that is 5% alcohol by volume).

Abbreviations: AUD=alcohol use disorder; CI=confidence interval; e-intervention=electronic intervention; g=grams; IVR=interactive voice response; KQ=key question; MD=mean difference; n=number; NA=not applicable; OR=odds ratio; RCT=randomized controlled trial; RR=risk ratio; SMD=standardized mean difference; SOE=strength of evidence

## SUMMARY OF EVIDENCE BY KEY QUESTION

### KQ 1. Characteristics of and User Support for E-Interventions

The e-interventions for alcohol misuse were typically brief, consisting of one session lasting 30 minutes or less. Intervention content was often focused on psychoeducation, PNF, and negative consequences expected from continuation of the participant's current level of alcohol consumption. Interventions including more complex psychotherapy content typically utilized a motivational interviewing approach, with relatively little cognitive-behavioral intervention beyond psychoeducation. Overall, the majority of e-interventions did not utilize supplementary human support. Even when such support was provided, it rarely included counseling beyond brief advice to reduce alcohol consumption levels. Only a single study meeting inclusion criteria used a smartphone as the delivery platform.

### KQ 2. Effects of E-Interventions Compared with Inactive Controls in Adults who Misuse Alcohol

We found low SOE that e-interventions relative to inactive controls (usual care, waitlist, or attention controls) decreased weekly alcohol consumption (n=17 studies) or increased the proportion meeting drinking limits (n=6 studies) in participants with alcohol misuse. Treatment effects varied importantly across studies, and we were unable to definitively explain this variability. Qualitative analyses suggested that studies using more intensive treatment were associated with greater reductions in alcohol consumption. This observation is consistent with a previous systematic review that found short-term effects of single-session PNF interventions were inferior to e-interventions with longer duration.<sup>27</sup> Relative to the variability in the observed effects, the summary effects were modest, ranging from 1 to 2 fewer drinks per week, suggesting a modest long-term ( $\geq 6$  months) effect of e-interventions on alcohol misuse outcomes. These estimates of treatment effect could be biased toward the null because control participants' behaviors may have been affected by assessments of alcohol misuse—similar to a placebo effect in drug studies.

While most studies provided data on the weekly volume of alcohol consumption, other important outcomes targeted in this review, such as effects on alcohol-related health and social or legal consequences, were reported infrequently. Moderately strong evidence showed no effects on binge drinking or alcohol-related social problems, but these outcomes were reported primarily in student samples. The SOE for the longer term ( $\geq 6$  months) benefits of e-interventions on alcohol consumption and its associated effects on health and well-being is low. However, due to the small number of studies evaluating several of the relevant outcomes, the brevity and limited scope of the e-interventions, and the lack of supplementary human support for these interventions, further research is warranted.

### KQ 3. Effects of E-Interventions Compared with Inactive Controls in Adults at High Risk of AUD (eg, AUDIT-C $\geq 8$ ) or with a Diagnosis of AUD

Three studies compared e-interventions with inactive controls or treatment as usual in participants with AUD. One study<sup>33</sup> randomized participants to one of 2 groups utilizing computerized feedback with up to 4 phone counseling sessions lasting 30 minutes each or

a control group receiving no intervention. This study is relatively novel in providing such extensive counseling, but data from the study did not provide evidence of long-term effects on alcohol outcomes. A second study evaluated a multicomponent smartphone application in patients completing residential treatment and found higher rates of abstinence and fewer risky drinking days.<sup>52</sup> The final study<sup>60</sup> recruited inpatients who were randomized at discharge to one of 2 groups using IVR assessments of drinking-related variables and tailored treatment modules with an option to request a counseling phone contact or a control group receiving no intervention. There was no significant effect of the e-intervention on reported alcohol outcomes. Thus, with widely varying interventions, there are conflicting findings on the efficacy of e-interventions for long-term alcohol consumption in participants with AUD. We found low SOE that a smartphone intervention decreased risk of relapse, but the evidence was insufficient for other e-interventions. There was insufficient evidence for evaluating other alcohol-related outcomes.

#### **KQ 4. Effects of E-Interventions Alone or Used in Combination with Face-to-Face Therapy Compared with Face-to-Face Therapy Alone in Adults who Misuse Alcohol**

Six studies compared e-interventions, used either independently or in conjunction with face-to-face counseling, to treatment consisting of face-to-face counseling alone. Three of the 6 studies combined e-interventions with face-to-face treatment. Those studies utilized IVR as the e-intervention and brief motivational interviewing (BMI) as the counseling component.

In collegiate populations, BMI delivered face-to-face was generally more effective than e-interventions alone in decreasing alcohol consumption. No studies compared these interventions in non-college samples. Although the available evidence suggests that e-interventions alone are inferior to face-to-face BMI, the small number of trials and the use of college student samples result in a low SOE. In studies using combined e-intervention and face-to-face counseling, 2 studies found no difference and one study found higher reported alcohol consumption with the combined intervention relative to face-to-face counseling alone. We judged the SOE for combined IVR and BMI interviewing as low. Other types of combined e-interventions have not been researched.

## **CLINICAL AND POLICY IMPLICATIONS**

Though prior systematic reviews have found positive short-term effects on alcohol consumption in students, those effects have generally not been maintained at longer term follow-up.<sup>24</sup> This is consistent with the results of our review, which examined trials reporting alcohol-related outcomes at  $\geq 6$  months and found absent to modest effects. In addition, the few available trials suggested that e-interventions might be less effective than face-to-face counseling, a finding also supported by prior reviews in student samples.<sup>24</sup> Prior syntheses that evaluated studies in adults have found few studies. Khadjesari and colleagues<sup>26</sup> concluded that “computer-based interventions may reduce alcohol consumption compared with assessment-only,” but only 6 of the 24 studies included in this review were conducted in non-student samples, and only 2 of those reported outcomes at  $\geq 6$  months. Riper and colleagues concluded that “E-self-help interventions without professional contact are effective in curbing adult problem drinking in high-income countries.”<sup>27</sup> However, 4 of the 9 studies assessed outcomes at  $\leq 3$  months, and 8



studies included in our review were not evaluated in the Riper review, primarily because they were published after the search date.

The presence of systematic annual screening for alcohol misuse in primary care using the Alcohol Use Disorders Identification Test for Clinicians (AUDIT-C) provides a good means of identifying patients who would be candidates for e-interventions if they decline specialty clinic treatment. However, the evidence in favor of existing e-interventions is low, suggesting that further work is needed to develop effective e-interventions for alcohol misuse. Further, it is not clear whether brief intervention for alcohol is effective for AUD specifically, as the majority of studies target alcohol misuse. This is problematic, as previous reviews of brief alcohol interventions have commented that they do not seem to be as effective in people with AUD.<sup>8,70</sup> Limited data from adult studies suggests a possible benefit of increasing the intensity of e-interventions by extending treatment duration, adding phone counseling, or utilizing mobile applications. More intensive supplementary human support might also improve participant engagement and the effectiveness of e-interventions. More research is needed to determine whether this is effective, as there were not enough e-interventions incorporating a human support component for us to carry out quantitative analyses that resulted in reliable estimates of effect modifiers.

If effective e-interventions can be developed and tested, they have the potential to address many of the barriers to face-to-face treatment of alcohol misuse. For example, some data suggest that Veterans would prefer the convenience of e-interventions<sup>71</sup> as opposed to negotiating the latency to treatment initiation, time investment, and travel barriers to attending clinic visits. In addition, health care providers cite barriers to addressing alcohol misuse at clinic visits that could be circumvented by utilizing e-interventions. The barriers for health care professionals include perceived inadequate training and treatment resources,<sup>72</sup> time constraints,<sup>73</sup> and concerns about negative effects on the doctor-patient relationship.<sup>74</sup>

There are also methods by which e-interventions could be designed to prevent relapse after the termination of intensive face-to-face counseling to achieve initial abstinence from alcohol. The continuing utilization of e-interventions could provide an accessible way for individuals to utilize treatment components originally introduced during face-to-face counseling. New programs could be tailored to a Veteran sample and could incorporate recent developments in treatment as well as be adapted for increasingly prevalent technologies such as smartphones. VHA has introduced some smartphone applications that offer assessment, basic coping tools, and referral to treatment resources. Designing e-intervention materials specifically for mobile devices could be accomplished by translating the content generated in counseling sessions for the patient to carry with them throughout the day, including high-risk situations. Clinicians and patients could also tailor e-intervention content to incorporate personalized motivations to stop drinking, avoid situations that increase the risk of relapse, and use coping strategies. The trial by Gustafson and colleagues<sup>52</sup> incorporated these principles and demonstrated benefit in maintaining abstinence in patients completing treatment for AUD.

Many of the e-interventions we reviewed were focused on psychoeducation and PNF, with relatively few data available on e-interventions incorporating treatment components that are a standard part of psychotherapy for alcohol misuse. Studies including a broader range of treatment

techniques typically reported application of motivational interviewing components. Given the central importance of relational components such as partnership and empathy in conventional face-to-face motivational interviewing, it is unclear the degree to which e-interventions actually replicate the core processes of motivational interviewing. Interventions may be enhanced by faithfully replicating core components of motivational interviewing, incorporating other proven therapies such as cognitive-behavioral techniques or pairing e-interventions with pharmacotherapy for alcohol misuse.

One recent example is a trial by Brief and colleagues<sup>75</sup> that was ineligible for inclusion in our quantitative analyses because the final follow-up assessment was at 3 months post-randomization. However, it is relevant to this review because it was conducted with Veterans and included coping skills for the common comorbidity of alcohol misuse. The e-intervention used 8 online modules lasting approximately 20 minutes each to provide personalized feedback, facilitate decisional balance, set drinking goals, and receive feedback on progress toward goals. Veterans also received information on skills for coping with trigger situations including PTSD-related symptoms (stress, mood, and anger management) and practices to promote normal nighttime sleep. Relative to waitlist, participants receiving the e-intervention significantly reduced alcohol consumption, binge drinking episodes, and alcohol-related problems. The median drinks per day decreased from 24 to 6 at 3-month follow-up. Results suggest the potential utility of e-interventions that are more intensive and broaden their strategies to address related problems that could be contributing to drinking behavior.

One expected contribution of e-interventions is the provision of low-cost clinical services to treat chronic problems such as alcohol misuse over a long period of time without utilizing a great deal of clinical resources. This efficient use of Veterans Affairs (VA) resources to improve the reach and long-term efficacy of alcohol treatment could provide valuable information to determine the value of e-interventions to the VA system. Consequently, clinical researchers could provide important contributions by including measures of cost-effectiveness in trials, especially when comparing e-interventions to traditional, clinic-based, face-to-face treatment.

The use of e-interventions for alcohol misuse will ultimately require attention to privacy and information security risks. Researchers must address these risks, eliminating them when possible and, if not possible, attending to the need to effectively communicate risks to Veterans using the technology. This includes the risk of information being intercepted during transmission to the study site, as well as breaches in the information stored at the study site and on computers and phones accessing the e-intervention.

## LIMITATIONS

Our review builds upon and extends prior reviews by following a protocol-based, transparent process that engaged relevant VA policy makers; including recently published studies; and taking an inclusive approach to electronic interventions. Another strength is the focus on sustained effects of e-interventions, prioritized as clinically important by our research team and technical expert panel. Our review has limitations as well. These include statistical assumptions in transforming data reported in many different formats into a common metric (eg, grams of alcohol per week), and judgments about when to combine multiple e-interventions tested in a single

trial for meta-analyses. We also made the judgment not to conduct subgroup analyses, having specified in advance that these analyses would be conducted only if subgroups contained at least 4 studies per group. Other research teams may have made different decisions.

The literature also had important limitations. Because outcomes were self-reported, daily IVR interventions may have “trained” participants to report more alcohol consumption more accurately, biasing these studies against an intervention effect. Others have found that assessment itself has been associated with reduced alcohol consumption,<sup>8</sup> so alcohol consumption reductions in the e-intervention groups could have been concealed by concurrent reductions in inactive control groups. The relatively low intensity of the interventions in the available literature on e-intervention for alcohol misuse limits the reliability of conclusions drawn about e-interventions as a general approach. In particular, the lack of data on the incremental efficacy of including various levels of supplementary human support prevents evaluation of potential applications of e-interventions for alcohol misuse. The small number of studies or subgroup analyses reporting on effects in participants with AUD limits interpretation for specialty clinic-based clinicians who typically treat patients with AUD specifically, as opposed to alcohol misuse generally. Other study limitations are described below.

### **Publication Bias**

To assess for publication bias, we searched ClinicalTrials.gov to identify completed but unpublished studies that would likely meet our eligibility criteria. Two such trials were identified from this search. The small number of trials reporting given outcomes prevented reliable quantitative evaluation of publication bias. Funnel plots were not used because analyses did not meet the minimum threshold of at least 10 studies for meaningful analysis.

### **Study Quality**

We assessed study quality by having 2 independent investigators provide a rating of risk of bias. Most trials were judged to be at moderate risk of bias (n=14); 5 were at low risk, and 7 were at high risk of bias. In this literature, the most common study limitations increasing risk of bias were lack of participant blinding to study condition and incomplete or perceived potential for selective reporting of outcome data.

### **Heterogeneity**

For effects on alcohol consumption, treatment effects varied from moderate to substantial. Variability in treatment effects were observed despite separate analyses for college samples and other adults. The variability in treatment effects on alcohol consumption limits the conclusions that can be drawn.

### **Applicability of Findings to the VA Population**

In VHA, Veterans are screened annually for alcohol misuse using the AUDIT-C; approximately 15% screen positive. Among those who screen positive, >80% screen positive for alcohol misuse, with <20% screening positive at a threshold (eg, AUDIT-C  $\geq$ 8) suggesting probable AUD. The majority of trials in this review used similar methods to enroll participants, and exclusion criteria were relatively few. In other aspects, these trials may have limited applicability

to VA. Only one trial<sup>50</sup> was conducted in a VA sample, half the trials were conducted in college students, and over half were conducted outside the United States. In the non-college samples, midlife adults, the majority of whom were college educated, participated; this is in contrast to the predominantly older population currently served by VA. Co-existing mental illness is common with alcohol misuse, but other mental illness diagnoses were not reported in these studies, introducing uncertainty about the applicability of these findings to Veterans with alcohol misuse and coexisting depression or anxiety disorders.

## RESEARCH GAPS/FUTURE RESEARCH

Although the current state of the evidence does not strongly support long-term benefit from e-interventions for alcohol misuse, the evidence for short-term effects from low-intensity interventions and longer-term effects from higher intensity interventions suggests the potential for benefit. Table 10 identifies gaps in evidence and study designs that could address these gaps. Although it would be possible to generate an extensive list of gaps in evidence, we restricted this list to the areas judged to be highest priority, given the current state of evidence. To facilitate future literature syntheses, we encourage investigators conducting clinical trials to include these studies in trial registries.

**Table 10. Highest Priority Evidence Gaps**

Evidence Gap	Study Designs
Uncertainty about effects in adults with varying educational levels and co-occurring mental and physical illness	RCTs in well-described samples with analyses for interaction effects by education and other mental or physical illness
Uncertain effects in older adults	Observational studies to evaluate interest, user interface issues, and barriers to e-interventions in older adults
Uncertainty about the effects of multi-session, multi-component interventions	RCTs
Uncertainty about whether effects found in alcohol misuse generally are applicable to individuals meeting criteria for an AUD	RCTs requiring AUD for inclusion or reporting AUD as a subgroup analysis
Uncertainty about effects of robust e-interventions used in combination with face-to-face therapy.	RCTs of combined treatments that include economic outcomes
Uncertainty about the validity of self-reported outcomes for alcohol consumption	Bioverification measures, the use of collateral informants to provide independent reports of drinking behavior, or mobile monitoring that would allow frequent breath alcohol assessments in participants' naturalistic environments

Abbreviations: AUD=alcohol use disorder; RCTs=randomized controlled trials

## CONCLUSIONS

The limited evidence from trials of e-interventions compared with controls on long-term ( $\geq 6$  months) alcohol outcomes in participants who screened positive for alcohol misuse suggests that effects are small or absent. The available data were even more limited for participants

with AUD or comparisons of e-interventions with face-to-face treatment. Further research is needed to determine with higher confidence whether e-interventions can produce long-term benefits for alcohol-related outcomes. In particular, given the limited number and duration of intervention episodes in the studies reviewed, it is possible that these e-interventions were simply not designed to produce enduring effects on alcohol misuse. As reported in previous reviews, brief, in-person interventions can produce sustained reductions in alcohol consumption in those with hazardous drinking who do not meet criteria for AUD, which likely provides benefits at the public health level, and that could be the primary value of these low-intensity interventions. At this time, the evidence does not support substitution of e-interventions for brief, in-person treatment. Future research on e-interventions should include evaluations of more intensive or longer duration alcohol e-interventions.

## REFERENCES

1. U.S. Burden of Disease Collaborators. The state of US health, 1990-2010: burden of diseases, injuries, and risk factors. *JAMA*. 2013;310(6):591-608.
2. Institute of Medicine. *Broadening the Base of Treatment of Alcohol Problems: Report of a Study by a Committee of the Institute of Medicine, Division of Mental Health and Behavioral Medicine*. Washington, DC: National Academies Press; 1990.
3. Cuijpers P, Riper H, Lemmers L. The effects on mortality of brief interventions for problem drinking: a meta-analysis. *Addiction*. 2004;99(7):839-845.
4. Grant BF, Stinson FS, Dawson DA, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2004;61(8):807-816.
5. World Health Organization. Global status report on alcohol and health. Geneva: WHO Press; 2011. [http://www.drugsandalcohol.ie/14675/1/Global\\_status\\_report\\_on\\_alcohol\\_and\\_health.pdf](http://www.drugsandalcohol.ie/14675/1/Global_status_report_on_alcohol_and_health.pdf). Accessed July 19, 2014.
6. Franck J, Jayaram-Lindstrom N. Pharmacotherapy for alcohol dependence: status of current treatments. *Curr Opin Neurobiol*. 2013;23(4):692-699.
7. Centers for Disease Control and Prevention. FastStats: alcohol use. <http://www.cdc.gov/nchs/fastats/alcohol.htm>. Last updated: February 28, 2014. Accessed July 18, 2014.
8. Jonas DE, Garbutt JC, Amick HR, et al. Behavioral counseling after screening for alcohol misuse in primary care: a systematic review and meta-analysis for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2012;157(9):645-654. <http://annals.org/article.aspx?articleid=1361859&resultClick=3>. Accessed September 25, 2014.
9. Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet*. 2009;373(9682):2223-2233.
10. U.S. Census Bureau. S2101. Veteran status. 2010 American Community Survey 1-year estimates. Retrieved July 22, 2014, from [http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_10\\_1YR\\_S2101&prodType=table](http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_10_1YR_S2101&prodType=table).
11. Wagner TH, Harris KM, Federman B, Dai L, Luna Y, Humphreys K. Prevalence of substance use disorders among veterans and comparable nonveterans from the National Survey on Drug Use and Health. *Psychol Serv*. 2007;4(3):149-157.
12. Hawkins EJ, Malte CA, Baer JS, Kivlahan DR. Prevalence, predictors, and service utilization of patients with recurrent use of Veterans Affairs substance use disorder specialty care. *J Subst Abuse Treat*. 2012;43(2):221-230.
13. Yu W, Ravelo A, Wagner TH, et al. Prevalence and costs of chronic conditions in the VA health care system. *Med Care Res Rev*. 2003;60(3 Suppl):146S-167S.

14. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Arlington, VA: American Psychiatric Publishing; 2013.
15. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. Washington, DC: American Psychiatric Association; 1994.
16. Barry KL, Blow FC, Willenbring ML, McCormick R, Brockmann LM, Visnic S. Use of alcohol screening and brief interventions in primary care settings: implementation and barriers. *Subst Abuse*. 2004;25(1):27-36.
17. Nilsen P. Brief alcohol intervention—where to from here? Challenges remain for research and practice. *Addiction*. 2010;105(6):954-959.
18. Williams EC, Johnson ML, Lapham GT, et al. Strategies to implement alcohol screening and brief intervention in primary care settings: a structured literature review. *Psychol Addict Behav*. 2011;25(2):206-214.
19. Pew Research Internet Project. Health fact sheet. <http://www.pewinternet.org/fact-sheets/health-fact-sheet/>. Accessed September 5, 2014.
20. Cunningham JA, Selby PL, Kypri K, Humphreys KN. Access to the Internet among drinkers, smokers and illicit drug users: is it a barrier to the provision of interventions on the World Wide Web? *Med Inform Internet Med*. 2006;31(1):53-58.
21. Vernon ML. A review of computer-based alcohol problem services designed for the general public. *J Subst Abuse Treat*. 2010;38(3):203-211.
22. Tait RJ, Christensen H. Internet-based interventions for young people with problematic substance use: a systematic review. *Med J Aust*. 2010;192(11 Suppl):S15-21.
23. Carey KB, Scott-Sheldon LA, Elliott JC, Bolles JR, Carey MP. Computer-delivered interventions to reduce college student drinking: a meta-analysis. *Addiction*. 2009;104(11):1807-1819.
24. Carey KB, Scott-Sheldon LA, Elliott JC, Garey L, Carey MP. Face-to-face versus computer-delivered alcohol interventions for college drinkers: a meta-analytic review, 1998 to 2010. *Clin Psychol Rev*. 2012;32(8):690-703.
25. Bewick BM, Trusler K, Barkham M, Hill AJ, Cahill J, Mulhern B. The effectiveness of web-based interventions designed to decrease alcohol consumption—a systematic review. *Prev Med*. 2008;47(1):17-26.
26. Khadjesari Z, Murray E, Hewitt C, Hartley S, Godfrey C. Can stand-alone computer-based interventions reduce alcohol consumption? A systematic review. *Addiction*. 2011;106(2):267-282.
27. Riper H, Spek V, Boon B, et al. Effectiveness of E-self-help interventions for curbing adult problem drinking: a meta-analysis. *J Med Internet Res*. 2011;13(2):e42.

28. Rooke S, Thorsteinsson E, Karpin A, Copeland J, Allsop D. Computer-delivered interventions for alcohol and tobacco use: a meta-analysis. *Addiction*. 2010;105(8):1381-1390.
29. Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. [www.cochrane-handbook.org](http://www.cochrane-handbook.org). Accessed July 22, 2014.
30. Elliott JC, Carey KB, Bolles JR. Computer-based interventions for college drinking: a qualitative review. *Addict Behav*. 2008;33(8):994-1005.
31. White A, Kavanagh D, Stallman H, et al. Online alcohol interventions: a systematic review. *J Med Internet Res*. 2010;12(5):e62.
32. Agency for Healthcare Research and Quality. *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. Rockville, MD: Agency for Healthcare Research and Quality. <http://www.effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productid=318>. Accessed July 22, 2014.
33. Bischof G, Grothues JM, Reinhardt S, Meyer C, John U, Rumpf HJ. Evaluation of a telephone-based stepped care intervention for alcohol-related disorders: a randomized controlled trial. *Drug Alcohol Depend*. 2008;93(3):244-251.
34. Kypri K, Langley JD, Saunders JB, Cashell-Smith ML, Herbison P. Randomized controlled trial of web-based alcohol screening and brief intervention in primary care. *Arch Intern Med*. 2008;168(5):530-536.
35. Neighbors C, Lewis MA, Atkins DC, et al. Efficacy of web-based personalized normative feedback: a two-year randomized controlled trial. *J Consult Clin Psychol*. 2010;78(6):898-911.
36. Walters ST, Vader AM, Harris TR, Field CA, Jouriles EN. Dismantling motivational interviewing and feedback for college drinkers: a randomized clinical trial. *J Consult Clin Psychol*. 2009;77(1):64-73.
37. Gual A, Martos AR, Lligona A, Llopis JJ. Does the concept of a standard drink apply to viticultural societies? *Alcohol Alcohol*. 1999;34(2):153-160.
38. Miller WR, Heather N, Hall W. Calculating standard drink units: international comparisons. *Br J Addict*. 1991;86(1):43-47.
39. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177-188.
40. Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Stat Med*. 2003;22(17):2693-2710.
41. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd edition. Hillside, NJ: Erlbaum Associates; 1988.



42. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539-1558.
43. Fu R, Gartlehner G, Grant M, et al. Conducting Quantitative Synthesis When Comparing Medical Interventions: AHRQ and the Effective Health Care Program. In: Agency for Healthcare Research and Quality. *Methods Guide for Comparative Effectiveness Reviews* [posted October 2010]. Rockville, MD. Available at: <http://effectivehealthcare.ahrq.gov/>. Accessed September 5, 2014.
44. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*. 2010;36(3):1-48.
45. Monahan CJ, McDevitt-Murphy ME, Dennhardt AA, Skidmore JR, Martens MP, Murphy JG. The impact of elevated posttraumatic stress on the efficacy of brief alcohol interventions for heavy drinking college students. *Addict Behav*. 2013;38(3):1719-1725.
46. Assistance Publique - Hôpitaux de Paris. Effect of an Interactive Computer Program to Prevent Alcohol Misuse. A Multicentre Randomized Controlled Trial. The BREVALCO Study. In: ClinicalTrials.gov [Internet]. Bethesda, MD: National Library of Medicine (US). <http://clinicaltrials.gov/show/NCT00658398>. Accessed September 11, 2014. NLM Identifier: NCT00658398.
47. Region Skane. Development of IVR and WEB Alcohol Interventions. In: ClinicalTrials.gov [Internet]. Bethesda, MD: National Library of Medicine (US). <http://clinicaltrials.gov/show/NCT01923246>. Accessed September 11, 2014. NLM Identifier: NCT01923246.
48. Barnett NP, Murphy JG, Colby SM, Monti PM. Efficacy of counselor vs. computer-delivered intervention with mandated college students. *Addict Behav*. 2007;32(11):2529-2548.
49. Boon B, Risselada A, Huiberts A, Riper H, Smit F. Curbing alcohol use in male adults through computer generated personalized advice: randomized controlled trial. *J Med Internet Res*. 2011;13(2):e43.
50. Cucciare MA, Weingardt KR, Ghaus S, Boden MT, Frayne SM. A randomized controlled trial of a web-delivered brief alcohol intervention in Veterans Affairs primary care. *J Stud Alcohol Drugs*. 2013;74(3):428-436.
51. Cunningham JA, Wild TC, Cordingley J, van Mierlo T, Humphreys K. A randomized controlled trial of an internet-based intervention for alcohol abusers. *Addiction*. 2009;104(12):2023-2032.
52. Gustafson DH, McTavish FM, Chih MY, et al. A smartphone application to support recovery from alcoholism: a randomized clinical trial. *JAMA Psychiatry*. 2014;71(5):566-572.
53. Hansen AB, Becker U, Nielsen AS, Gronbaek M, Tolstrup JS, Thygesen LC. Internet-based brief personalized feedback intervention in a non-treatment-seeking population of adult heavy drinkers: a randomized controlled trial. *J Med Internet Res*. 2012;14(4):e98.

54. Hasin DS, Aharonovich E, O’Leary A, et al. Reducing heavy drinking in HIV primary care: a randomized trial of brief intervention, with and without technological enhancement. *Addiction*. 2013;108(7):1230-1240.
55. Helzer JE, Rose GL, Badger GJ, et al. Using interactive voice response to enhance brief alcohol intervention in primary care settings. *J Stud Alcohol Drugs*. 2008;69(2):251-258.
56. Hester RK, Delaney HD, Campbell W. The college drinker’s check-up: outcomes of two randomized clinical trials of a computer-delivered intervention. *Psychol Addict Behav*. 2012;26(1):1-12.
57. Kypri K, Hallett J, Howat P, et al. Randomized controlled trial of proactive web-based alcohol screening and brief intervention for university students. *Arch Intern Med*. 2009;169(16):1508-1514.
58. Kypri K, Saunders JB, Williams SM, et al. Web-based screening and brief intervention for hazardous drinking: a double-blind randomized controlled trial. *Addiction*. 2004;99(11):1410-1417.
59. Moreira MT, Oskrochi R, Foxcroft DR. Personalised normative feedback for preventing alcohol misuse in university students: Solomon three-group randomised controlled trial. *PLoS One*. 2012;7(9):e44120.
60. Mundt JC, Moore HK, Bean P. An interactive voice response program to reduce drinking relapse: a feasibility study. *J Subst Abuse Treat*. 2006;30(1):21-29.
61. Neighbors C, Larimer ME, Lewis MA. Targeting misperceptions of descriptive drinking norms: efficacy of a computer-delivered personalized normative feedback intervention. *J Consult Clin Psychol*. 2004;72(3):434-447.
62. Neumann T, Neuner B, Weiss-Gerlach E, et al. The effect of computerized tailored brief advice on at-risk drinking in subcritically injured trauma patients. *J Trauma*. 2006;61(4):805-814.
63. Riper H, Kramer J, Smit F, Conijn B, Schippers G, Cuijpers P. Web-based self-help for problem drinkers: a pragmatic randomized trial. *Addiction*. 2008;103(2):218-227.
64. Schulz DN, Candel MJ, Kremers SP, Reinwand DA, Jander A, de Vries H. Effects of a Web-based tailored intervention to reduce alcohol consumption in adults: randomized controlled trial. *J Med Internet Res*. 2013;15(9):e206.
65. Sinadinovic K, Wennberg P, Berman AH. Targeting problematic users of illicit drugs with Internet-based screening and brief intervention: a randomized controlled trial. *Drug Alcohol Depend*. 2012;126(1-2):42-50.
66. Voogt CV, Poelen EA, Kleinjan M, Lemmers LA, Engels RC. The effectiveness of the ‘what do you drink’ web-based brief alcohol intervention in reducing heavy drinking among students: a two-arm parallel group randomized controlled trial. *Alcohol Alcohol*. 2013;48(3):312-321.

67. Wallace P, Murray E, McCambridge J, et al. On-line randomized controlled trial of an internet based psychologically enhanced intervention for people with hazardous alcohol consumption. *PLoS One*. 2011;6(3):e14740.
68. Collins SE, Witkiewitz K, Larimer ME. The theory of planned behavior as a predictor of growth in risky college drinking. *J Stud Alcohol Drugs*. 2011;72(2):322-332.
69. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383-394.
70. O'Donnell A, Anderson P, Newbury-Birch D, et al. The impact of brief alcohol interventions in primary healthcare: a systematic review of reviews. *Alcohol Alcohol*. 2014;49(1):66-78.
71. Lapham GT, Hawkins EJ, Chavez LJ, et al. Feedback from recently returned veterans on an anonymous web-based brief alcohol intervention. *Addict Sci Clin Pract*. 2012;7(1):17.
72. Griffiths RD, Stone A, Tran DT, Fernandez RS, Ford K. Drink a little; take a few drugs: do nurses have knowledge to identify and manage in-patients at risk of drugs and alcohol? *Drug Alcohol Rev*. 2007;26(5):545-552.
73. Happell B, Carta B, Pinikahana J. Nurses' knowledge, attitudes and beliefs regarding substance use: a questionnaire survey. *Nurs Health Sci*. 2002;4(4):193-200.
74. Nygaard P, Aasland OG. Barriers to implementing screening and brief interventions in general practice: findings from a qualitative study in Norway. *Alcohol Alcohol*. 2011;46(1):52-60.
75. Brief DJ, Rubin A, Keane TM, et al. Web intervention for OEF/OIF veterans with problem drinking and PTSD symptoms: a randomized clinical trial. *J Consult Clin Psychol*. 2013;81(5):890-900.