

Propensity Scores

Todd Wagner, PhD

February 6, 2019

Learning Objectives

We will discuss:

1. What is a propensity score
2. Ways to implement a propensity score
3. The assumptions needed to make causal claims with observational data

Outline

1. Background on assessing causation
2. Define propensity score (PS)
3. Calculate the PS
4. Use the PS
5. Limitations of the PS


Causality

- Researchers are often interested in understanding causal relationships
 - Does drinking red wine affect health?
 - Does a new treatment reduce symptoms?
 - Does job burnout affect risk of suicidality?
 - Does the Veterans Crisis Line reduce the likelihood of suicide?

Randomized Clinical Trial

- RCT provides a methodological approach for understanding causation
- Understanding propensity score is assisted by understanding randomized trials.

Randomization



Note: random sorting can, by chance, lead to unbalanced groups. Most trials use checks and balances to preserve randomization

Just because a RCT can speak to causality, you must ask the question for whom— generalizability is often very limited

Trial analysis

- The expected effect of treatment is

$$E(Y) = E(Y^A) - E(Y^B)$$

Expected effect on group A minus expected effect on group B (i.e., mean difference).

Trial Analysis (II)

- $E(Y) = E(Y^A) - E(Y^B)$ can be analyzed using the following general model

$$y_i = \alpha + \beta x_i + \varepsilon_i$$

Where

- y is the outcome
- α is the intercept
- x is the mean difference in the outcome between treatment A relative to treatment B
- ε is the error term
- i denotes the unit of analysis (person)

Trial Analysis (III)

- The model can be expanded to control for baseline characteristics

$$y_i = \alpha + \beta x_i + \delta Z_i + \varepsilon_i$$

Where

- y is outcome
- α is the intercept
- x is the added value of the treatment A relative to treatment B
- Z is a vector of baseline characteristics (predetermined prior to randomization)
- ε is the error term
- i denotes the unit of analysis (person)

Assumptions

- Right hand side variables are measured without noise (i.e., considered fixed in repeated samples)
- There is no correlation between the right hand side variables and the error term
 $E(x_i \varepsilon_i) = 0$
- If these conditions hold, β is an unbiased estimate of the **causal** effect of the treatment on the outcome



What if...

- The assumptions don't hold in an RCT. Then what?
- You lose the unbiased estimate of causality.

Observational Studies

- Randomized trials may be
 - Unethical
 - Infeasible
 - Impractical
 - Not scientifically justified

Endogenous

- Poll-
- Has anyone heard of this term?
 - Yes, I use the term frequently at the dinner table
 - Yes, I have heard others use it, but it still confuses me
 - No, and I like being honest


Endogenous

- Not attributable to any external factor.
- Example: Does smoking lead to cancer

$$\text{cancer}_i = \alpha + \beta \text{smoking}_i + \varepsilon_i$$

- Smoking is correlated with income, education, parental exposure, etc.
- We aren't controlling for any of those factors, thus $E(\text{smoking}_i, \varepsilon_i) \neq 0$
- Thus, smoking is endogenous


Sorting without randomization




If everything is fully observed and correctly specified;
results are not biased. **Never happens in reality.**

Based on: Maciejewski and Pizer (2007) Propensity Scores and Selection Bias in Observational Studies. HERC Cyberseminar

Sorting without randomization



Sorting without randomization



Propensity Score Defined

- The PS uses observed information, which is multi-dimensional, to calculate a single variable (the score)
- The score is the predicted propensity to get sorted (usually thought of as propensity to get treatment).

Expected treatment effect: $E(Y) = E(Y^A) - E(Y^B)$

Propensity Score is: $\Pr(Y=A \mid X_i)$

Propensity Scores

- What it is: Another way to correct for observable characteristics
- What it is not: A way to adjust for unobserved characteristics
- The only way to make causal claims is to make **huge** assumptions.

Strong Ignorability

- To make statements about causation, you would need to make an assumption that treatment assignment is strongly ignorable.
 - Similar to assumptions of missing at random
 - Equivalent to stating that all variables of interest are observed


Creating a Propensity Score

Calculating the Propensity Score

- You observe treatment: One group receives it and another group doesn't
- Use multivariate logistic regression to estimate the probability that a person received treatment
- The predicted probability from the logistic model is the propensity score

Variables to Include

- Include variables that are related to the observed outcome
- This will decrease the variance of an estimated exposure effect without increasing bias
- Do not include variables affect only correlated with exposure



Brookhart MA, et al Am J Epidemiol. 2006 Jun 15;163(12):1149-56.

Variables to Exclude

- Exclude variables that are related to the exposure but not to the outcome
- These variables will increase the variance of the estimated exposure effect without decreasing bias
- Variable selection is particularly important in small studies ($n < 500$)

Example: Resident Surgery

- Do cardiac bypass patients have better / worse outcomes when their surgery is conducted by a resident or an attending?
- We had a datasets that tracked the primary surgeon for heart bypass

Uses

- Understanding sorting and balance
 - Sorting is multidimensional
 - The PS provides a simple way of reducing this dimensionality to understand the similarity of the treatment groups
- Adjusting for covariance

Example

- Are surgical outcomes worse when the surgeon is a resident?
- Resident assignment may depend on
 - Patient risk
 - Availability of resident
 - Resident skill
 - Local culture

Resident Assignment

	OR	P value
Age	1.00	0.79
Canadian Functional Class		
Class 2	1.93	0.15
Class 3	2.12	0.09
Class 4	4.25	0.02
Urgent priority	0.93	0.89
Artery condition at site		
Calcified	0.67	0.25
Sclerotic	2.63	0.00
site 2	62.89	0.00
site 3	0.67	0.60
site 5	138.16	0.00
site 7	11.66	0.00
site 8	19.85	0.00
site 9	1.76	0.43
endo vascular harvest	0.20	0.01
On pump surgery	1.20	0.75
1-2 grafts	1.70	0.16
4-5 grafts	0.79	0.46


Assignment not associated with age or number of grafts

Assignment associated with angina symptoms and planned harvesting technique




Shared or Common Support

- Concept that measures overlap of people in both treatments
- Conditional on covariates, there exist people who choose both treatments.
- Poor common support suggests that conditional on observables, we cannot control for sorting.

Propensity Score for Resident vs Attending Surgeon






Compare Three Scores



Poll

- Which graph is the most concerning?
Choose one
- A
- B
- C
- All of them
- None of them



Three Scores



RCTs and Propensity Scores

- What would happen if you used a propensity score with data from a RCT?

Shared Common Support



Common Support

- Some evidence that propensity scores provide some advantages when there is considerable shared support
- Yet, growing evidence that it can create problems.

Using the Propensity Score

Using the Propensity Score

1. Compare individuals based on similar PS scores (a matched analysis)
2. Conduct subgroup analyses on similar groups (stratification)
3. Include it as a covariate (quintiles of the PS) in the regression model
4. Use it to weight the regression (i.e., place more weight on similar cases)
5. Use both 3 and 4 together (doubly robust)

PS as a Covariate

- There seems to be little advantage to using PS over multivariate analyses in most cases.¹
- PS provides flexibility in the functional form
- Propensity scores may be preferable if the sample size is small and the outcome of interest is rare.²

1. Winkelmeier. Nephrol. Dial. Transplant 2004; 19(7): 1671-1673.

2. Cepeda et al. Am J Epidemiol 2003; 158: 280-287

Matched Analyses

- The idea is to select controls that resemble the treatment group in all dimensions, except for treatment
- You can exclude cases and controls that don't match, which can reduce the sample size/power.
- Different matching methods

Matching Methods

- Nearest Neighbor: rank the propensity score and choose control that is closest to case.
- Caliper: choose your common support and from within randomly draw controls
- Choice of matching estimator important

Recent Areas of Research

■ Economics: choice of matching estimators

- Busso M et al. New Evidence on the Finite Sample Properties of Propensity Score Reweighting and Matching Estimators. *Review of Economics and Statistics*, 96.5 (2014): 885-897

■ Political Science

- King G, Nielsen R. Why propensity scores should not be used for matching. Copy at <http://j.mp/1sexgVw>. 2016 Dec 16;378.

■ Biostatistics: high dimensional propensity scores using big data

- Schneeweiss, Sebastian, et al. "High-dimensional propensity score adjustment in studies of treatment effects using health care claims data." *Epidemiology* 20.4 (2009): 512.

Limitations

Do the Unobservables Matter?

- Propensity scores focus only on observed characteristics, not on unobserved.
- Improbable that we fully observe the sorting process
 - Thus $E(x_i \varepsilon_i) \neq 0$
 - Multivariate (including propensity score) is biased and we need another method, such as instrumental variables, fixed effects or RCT

Does Using PS Exacerbate Imbalance of Unobservables

- PS is based on observables.
- Brooks and Ohsfeldt, using simulated data, showed that PS models can create **greater** imbalance among unobserved variables.
- King G, Nielsen R. Why propensity scores should not be used for matching. Copy at [http://j. mp/1sexgVw](http://j.mp/1sexgVw) Download. 2016 Dec 16;378.

Brooks and Ohsfeldt (2013): Squeezing the balloon: propensity scores and unmeasured covariate balance. *Health Services Research*.

Summary

Overview

- Propensity scores offer another way to adjust for confound by observables
- Reducing the multidimensional nature of confounding can be helpful
- There are many ways to implement propensity scores and a growing interest in matching estimators

Strengths

- Allow one to check for balance between control and treatment
- Without balance, average treatment effects can be very sensitive to the choice of the estimators.¹

1. Imbens and Wooldridge 2007 http://www.nber.org/WNE/lect_1_match_fig.pdf

Challenges

- Propensity scores are often misunderstood
- Not enough attention is placed on the PS model, itself
- Not enough attention is placed on robustness checks
- While a PS can help create balance on observables, PS models do not control for unobservables or selection bias

Further Reading

- Rosenbaum, P. R., D. B. Rubin. "The central role of the propensity score in observational studies for causal effects." *Biometrika* 70 (1983): 41–55
- Imbens and Wooldridge (2007) www.nber.org/WNE/lect_1_match_fig.pdf
- Imbens, Guido W. "The role of the propensity score in estimating dose-response functions." *Biometrika* 87.3 (2000): 706-710.
- Imbens, Guido W. "Nonparametric estimation of average treatment effects under exogeneity: A review." *Review of Economics and Statistics* 86.1 (2004): 4-29.
- Guo and Fraser (2010) *Propensity Score Analysis*. Sage.
- King G, Nielsen R. Why propensity scores should not be used for matching. Copy at <http://j.mp/1sexgVw>. 2016 Dec 16;378.
- Brooks, John M., and Robert L. Ohsfeldt. "Squeezing the balloon: propensity scores and unmeasured covariate balance." *Health Services Research* 48.4 (2013): 1487-1507.
- Garrido, Melissa M., et al. "Methods for constructing and assessing propensity scores." *Health Services Research* 49.5 (2014): 1701-1720.
- Busso M et al. "New Evidence on the Finite Sample Properties of Propensity Score Reweighting and Matching Estimators." *Review of Economics and Statistics*, 96.5 (2014): 885-897
- Imai, Kosuke, and Marc Ratkovic. "Covariate balancing propensity score." *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 76.1 (2014): 243-263.

Questions?

- HERC@VA.gov

- [@herc_va](https://twitter.com/herc_va)

- [@toddhwagner](https://twitter.com/toddhwagner)



- **Next class: Natural Experiments and Difference-in-Differences**

Jean Yoon, Ph.D. Feb 13.