

















































































































## Assessment and Management of Pain in Inpatients

81. Uden P, Starck CJ, Berger T. Comparison of indomethacin and dihydromorphinone in acute, biliary stone pain. *Current Ther Res.* 1984;36(6):1228-1234.
82. Halbert J, Crotty M. Evidence for the optimal management of acute and chronic phantom pain: a systematic review. *Clin J Pain* 2002;18(2):84-92.
83. Sherman RA, Sherman CJ, Gall NG. A survey of current phantom limb pain treatment in the United States. *Pain.* 1980;8:85-99.
84. Jaeger H, Maier C. Calcitonin in phantom limb pain: a double-blind study. *Pain.* Jan 1992;48(1):21-27.
85. Nikolajsen L, Hansen CL, Nielsen J, Keller J, Arendt-Nielsen L, Jensen TS. The effect of ketamine on phantom pain: a central neuropathic disorder maintained by peripheral input. *Pain.* Sep 1996;67(1):69-77.
86. Huse E, Larbig W, Flor H, Birbaumer N. The effect of opioids on phantom limb pain and cortical reorganization. *Pain.* Feb 1 2001;90(1-2):47-55.
87. Wu CL, Tella P, Staats PS, et al. Analgesic effects of intravenous lidocaine and morphine on postamputation pain: a randomized double-blind, active placebo-controlled, crossover trial. *Anesthesiology.* Apr 2002;96(4):841-848.
88. Bone M, Critchley P, Buggy DJ. Gabapentin in postamputation phantom limb pain: a randomized, double-blind, placebo-controlled, cross-over study. *Regional Anesthesia & Pain Medicine.* Sep-Oct 2002;27(5):481-486.
89. Dertwinkel R, Henrichs C, Senne I, et al. Prevention of severe phantom limb pain by perioperative administration of ketamine - an observational study. *Acute Pain.* 2002;4:9-13.
90. Hayes C, Armstrong-Brown A, Burstal R. Perioperative intravenous ketamine infusion for the prevention of persistent post-amputation pain: a randomized, controlled trial. *Anaesthesia & Intensive Care.* Jun 2004;32(3):330-338.
91. Gehling M, Tryba M. [Prophylaxis of phantom pain: is regional analgesia ineffective?]. *Schmerz.* Jan 2003;17(1):11-19.
92. Flor H, Denke C, Schaefer M, Grusser S. Effect of sensory discrimination training on cortical reorganisation and phantom limb pain. *Lancet.* Jun 2 2001;357(9270):1763-1764.
93. Cook AK, Niven CA, Downs MG. Assessing the pain of people with cognitive impairment. *International Journal of Geriatric Psychiatry.* Jun 1999;14(6):421-425.
94. Weiner D, Peterson B, Ladd K, McConnell E, Keefe F. Pain in nursing home residents: An exploration of prevalence, staff perspectives, and practical aspects of measurement. *Clin J Pain.* 1999;15(2):92-101.
95. Parmelee P. Pain complaints and cognitive status among elderly institution residents. . *Journal of the American Geriatrics Society.* 1993;41(5):517-522.
96. Pautex S, Michon A, Guedira M, et al. Pain in severe dementia: self-assessment or observational scales?. *Journal of the American Geriatrics Society.* 2006;54(7):1040-1045.
97. Buffum MD, Hutt E, Chang VT, Craine MH, Snow AL. Cognitive impairment and pain management: Review of issues and challenges. *Journal of Rehabilitation Research & Development.* 2007;44(2):315-330.

## Assessment and Management of Pain in Inpatients

98. Assessing Pain in the Patient with Impaired Communication: A Consensus Statement from the VHA National Pain Management Strategy Coordinating Committee. In: Affairs V, ed; 2004.
99. Pautex S, Herrmann F, Le Lous P, Fabjan M, Michel JP, Gold G. Feasibility and reliability of four pain self-assessment scales and correlation with an observational rating scale in hospitalized elderly demented patients. . *Journals of Gerontology Series A-Biological Sciences & Medical Sciences*. Apr 2005;60(4):524-529.
100. Defrin R, Lotan M, Pick CG. The evaluation of acute pain in individuals with cognitive impairment: a differential effect of the level of impairment. *Pain*. Oct 2006;124(3):312-320.
101. Chibnall JT, Tait RC. Pain assessment in cognitively impaired and unimpaired older adults: a comparison of four scales. *Pain*. May 2001;92(1-2):173-186.
102. Herr K, Bjoro K, Decker S. Tools for assessment of pain in nonverbal older adults with dementia: a state-of-the-science review. *Journal of Pain & Symptom Management*. Feb 2006;31(2):170-192.
103. van Herk R, van Dijk M, Baar FP, Tibboel D, de Wit R. Observation scales for pain assessment in older adults with cognitive impairments or communication difficulties. *Nurs Res*. 2007;56(1):34-43.
104. Zwakhalen SMG, Hamers JPH, Abu-Saad HH, Berger MPF. Pain in elderly people with severe dementia: a systematic review of behavioural pain assessment tools. *BMC Geriatrics*. 2006;6:3.
105. Definitions Related to the Use of Opioids for the Treatment of Pain. <http://www.ampainsoc.org/advocacy/opioids2.htm>. Accessed December 10, 2007, 2007.
106. Savage SR, Joranson DE, Covington EC, Schnoll SH, Heit HA, Gilson AM. Definitions related to the medical use of opioids: evolution towards universal agreement. . *J Pain Symptom Manage*. 2003;26:655–667.
107. Hojsted J, Sjogren P. Addiction to opioids in chronic pain patients: A literature review. *European Journal of Pain*. 2007;11(5):490-518.
108. American Medical Directors Association. We care: tool kit for implementation of the clinical practice guideline for pain management. Columbia (MD): American Medical Directors Association (AMDA); 2004.
109. ACOVE quality indicators. Assessing the Care of Vulnerable Elders Project. Rand Health, Santa Monica, California. Chodosh J, Ferrell BA, Shekelle PG, et al. Quality indicators for pain management in vulnerable elders. *Ann Intern Med*. 2001; 135 (8pt2):641-758, 731-738.
110. Agency for Healthcare Research and Quality (AHRQ). CAHPS hospital survey. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2006 Feb 1. 4 p. Centers for Medicare & Medicaid Services (CMS). HCAHPS survey [<http://www.hcahpsonline.org>]. Baltimore (MD): Centers for Medicare & Medicaid Services (CMS)