



The Impact of Wearable Motion Sensing Technologies on Physical Activity: A Systematic Review

September 2015

Prepared for:

Department of Veterans Affairs
Veterans Health Administration
Quality Enhancement Research Initiative
Health Services Research & Development Service
Washington, DC 20420

Prepared by:

Evidence-based Synthesis Program (ESP) Center
Durham VA Healthcare System
Durham, NC
John W. Williams, Jr., MR, MHSc, Director

Principal Investigators:

Jennifer M. Gierisch, PhD, MPH
Adam P. Goode, DPT, PhD

Co-Investigators:

Bryan C. Batch, MD
Kim N. Huffman, MD, PhD
Katherine S. Hall, PhD
S. Nicole Hastings, MD, MHS
Kelli D. Allen, PhD
Ryan J. Shaw, PhD, RN
Frances A. Kanach, PhD
Jennifer R. McDuffie, PhD
Andrzej S. Kosinski, PhD
John W. Williams, Jr., MD, MHSc

Research Associate:

Avishek Nagi, MS



VA
HEALTH CARE
Defining
EXCELLENCE
in the 21st Century

PREFACE

The VA Evidence-based Synthesis Program (ESP) was established in 2007 to provide timely and accurate syntheses of targeted healthcare topics of particular importance to clinicians, managers, and policymakers as they work to improve the health and healthcare of Veterans. QUERI provides funding for four ESP Centers, and each Center has an active University affiliation. Center Directors are recognized leaders in the field of evidence synthesis with close ties to the AHRQ Evidence-based Practice Centers. The ESP is governed by a Steering Committee comprised of participants from VHA Policy, Program, and Operations Offices, VISN leadership, field-based investigators, and others as designated appropriate by QUERI/HSR&D.

The ESP Centers generate evidence syntheses on important clinical practice topics. These reports help:

- Develop clinical policies informed by evidence;
- Implement effective services to improve patient outcomes and to support VA clinical practice guidelines and performance measures; and
- Set the direction for future research to address gaps in clinical knowledge.

The ESP disseminates these reports throughout VA and in the published literature; some evidence syntheses have informed the clinical guidelines of large professional organizations.

The ESP Coordinating Center (ESP CC), located in Portland, Oregon, was created in 2009 to expand the capacity of QUERI/HSR&D and is charged with oversight of national ESP program operations, program development and evaluation, and dissemination efforts. The ESP CC establishes standard operating procedures for the production of evidence synthesis reports; facilitates a national topic nomination, prioritization, and selection process; manages the research portfolio of each Center; facilitates editorial review processes; ensures methodological consistency and quality of products; produces “rapid response evidence briefs” at the request of VHA senior leadership; collaborates with HSR&D Center for Information Dissemination and Education Resources (CIDER) to develop a national dissemination strategy for all ESP products; and interfaces with stakeholders to effectively engage the program.

Comments on this evidence report are welcome and can be sent to Nicole Floyd, ESP CC Program Manager, at Nicole.Floyd@va.gov.

Recommended citation: Gierisch JM, Goode AP, Batch BC, Huffman KM, Hall KS, Hastings SN, Allen KD, Shaw RJ, Kanach FA, McDuffie JR, Kosinski AS, Nagi A, Williams JW Jr. The Impact of Wearable Motion Sensing Technologies on Physical Activity: A Systematic Review. VA ESP Project #09-010; 2015.

This report is based on research conducted by the Evidence-based Synthesis Program (ESP) Center located at the **Durham VA Medical Center, Durham, NC**, funded by the Department of Veterans Affairs, Veterans Health Administration, Office of Research and Development, Quality Enhancement Research Initiative. The findings and conclusions in this document are those of the author(s) who are responsible for its contents; the findings and conclusions do not necessarily represent the views of the Department of Veterans Affairs or the United States government. Therefore, no statement in this article should be construed as an official position of the Department of Veterans Affairs. No investigators have any affiliations or financial involvement (eg, employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties) that conflict with material presented in the report.

STRUCTURED ABSTRACT

Background: Participation in regular physical activity is important for improving health, but sedentary behavior is difficult to change. One option is to provide feedback on physical activity with wearable motion sensing technologies (activity devices). This review sets out to synthesize the literature on the effectiveness of these devices for physical activity, weight, and patient satisfaction outcomes, and to describe moderating factors that may impact effectiveness (*ie*, population characteristics, location where device is worn on body, or device role in overall intervention approach).

Methods: We searched MEDLINE, Embase, CINAHL, SPORTDiscus, and Cochrane CENTRAL from January 1, 2000, to January 6, 2015, for peer-reviewed, English-language randomized controlled trials among adults (≥ 18 years of age). Article inclusion, data abstraction, and quality assessment were conducted through a duplicate process, with discussion to resolve discrepancies. Trial quality was evaluated as low, unclear, or high risk of bias (ROB). Strength of evidence (SOE) was summarized as high, moderate, or low. Random-effects models were used to produce standardized mean differences (SMDs) for physical activity outcomes and mean differences (MDs) for weight outcomes. Heterogeneity was measured with I^2 . Qualitative synthesis was conducted for outcomes with < 3 studies.

Results: We identified 4787 unique citations; 14 trials met eligibility criteria. Women comprised 62.5% of the population. Median age was 49.7 years (range 28.7 to 79.8 years). Study sizes ranged from 20 to 544 participants (median 62), with the majority of studies ($n=8$) randomizing < 70 participants. Although all of the interventions had multiple components, in the majority of studies ($n=8$), the wearable device was used in a major role (*ie*, central motivational enhancement). The device was an accelerometer in all 14 studies.

Twelve trials (2 at low ROB, 2 at unclear ROB, 8 at high ROB) examined accelerometer interventions for increasing physical activity; the majority ($n=9$) used an inactive comparator. Overall, a small significant effect was found for increasing physical activity (SMD 0.26; 95% CI 0.04 to 0.49) with high heterogeneity ($I^2=64.7\%$). Moderate SOE was found for small increases in physical activity when compared with an inactive comparator (SMD 0.29; 95% CI 0.03 to 0.55) with high heterogeneity ($I^2=70.3\%$). Low SOE and no statistically significant effect (SMD 0.17; 95% CI -1.09 to 1.43) were found when compared with an active comparator.

Eleven trials (2 at low ROB, 3 at unclear ROB, 6 at high ROB) examined the effect of accelerometer interventions on weight loss or maintenance. The overall pooled estimate showed a small significant effect for weight loss (MD -1.65 kg; 95% CI -3.03 to -0.28) with high heterogeneity ($I^2=81\%$). Moderate SOE and no significant effect were found for accelerometers versus inactive comparators (MD -1.44 kg; 95% CI -3.08 to 0.19). A positive trend with low SOE for accelerometers was found in 2 trials on weight loss, but only one was statistically significant.

No studies reported the outcome of patient satisfaction with healthcare. Also, no moderating factors were found to significantly impact effectiveness or explain heterogeneity.

Conclusions: The small positive effects produced by interventions that include accelerometers may not result in a clinically significant impact on physical activity or weight loss; however, the small sample sizes with moderate to high heterogeneity in the current studies limit the conclusions that may be drawn. Larger, well-designed randomized controlled trials are needed. Clinicians and policymakers should consider these findings and the existing gaps in the literature before widespread use of these technologies.

ABBREVIATIONS TABLE

AHRQ	Agency for Healthcare Research and Quality
CCRB	Cochrane Collaboration Risk of Bias Tool
CI	Confidence interval
COPD	Chronic obstructive pulmonary disease
ESP	Evidence-based Synthesis Program
GPS	Global positioning system
HSR&D	Health Services Research & Development
KQ	Key question
MD	Mean difference
MeSH	Medical Subject Heading
NCP	National Center for Health Promotion and Disease Prevention
PA	Physical activity
PICOTS	Population, intervention, comparator, outcome, timing, and setting
QUERI	Quality Enhancement Research Initiative
RCT	Randomized controlled trial
ROB	Risk of bias
SEM	Standard error of the mean
SMD	Standardized mean difference
SOE	Strength of evidence
VA	Veterans Affairs
VHA	Veterans Health Administration
VISN	Veterans Integrated Service Networks

TABLE OF CONTENTS

INTRODUCTION	1
METHODS	3
Topic Development.....	3
Search Strategy	3
Study Selection	4
Data Abstraction	5
Quality Assessment.....	5
Data Synthesis.....	6
Rating the Body of Evidence	7
Peer Review	7
RESULTS	8
Literature Flow.....	8
Characteristics of Included Studies.....	9
KQ 1: Among adults, what is the effectiveness of wearable motion sensing technologies on: (a) physical activity levels; (b) weight loss or maintenance; (c) patient satisfaction with healthcare?	12
Key Points.....	12
Physical Activity (KQ 1a).....	12
Weight Loss or Maintenance (KQ 1b).....	14
Patient Satisfaction (KQ 1c)	16
Quality of Evidence for KQ 1	16
KQ 2: Among adults, does the impact of wearable motion sensing technologies vary by: (a) characteristics of the population; (b) type of adjunctive interventions; (c) adherence to use of the devices; or (d) characteristics of the device?	20
Key Points.....	20
Variation by Population Characteristics (KQ 2a)	20
Variation by Activity Device Role (Major vs Minor; KQ 2b).....	25
Variation by Adherence to Use of the Activity Device (KQ 2c).....	28
Variation by Characteristics of the Activity Device (Body Location; KQ 2d).....	31
Quality of Evidence for KQ 2.....	35
SUMMARY AND DISCUSSION	36
Strength of Evidence.....	36
Summary of Evidence by Key Question.....	38
Clinical and Policy Implications.....	39
Limitations	40
Research Gaps/Future Research	41
Conclusions.....	42
REFERENCES	44

TABLES

Table 1. Inclusion and Exclusion Criteria..... 4
 Table 2. Approach to Formulating Summary Risk of Bias for Each Outcome Across Domains... 6
 Table 3. Description of Interventions and Comparators..... 9
 Table 4. Summary of Findings: Variation by Population Subgroups (KQ 2a)..... 25
 Table 5. Summary of Findings: Variation by Device Role (KQ 2b)..... 28
 Table 6. Summary of Findings: Accelerometer Adherence Across Individual Studies of
 Physical Activity and Weight Loss (KQ 2c)..... 30
 Table 7. Summary of Findings: Variation by Characteristics of the Device (KQ 2d)..... 35
 Table 8. Summary of Intervention Effects and Strength of Evidence Ratings..... 37
 Table 9. Evidence Gaps and Future Research 41

FIGURES

Figure 1. Literature Flow Chart 8
 Figure 2. Effect of Accelerometer Interventions on Physical Activity..... 13
 Figure 3. Effect of Accelerometer Interventions on Weight Loss or Maintenance 15
 Figure 4. Risk of Bias Graph^a 16
 Figure 5. Risk of Bias Summary: Review Authors’ Judgments About Risk of Bias Items
 for Each Included Study^a 19
 Figure 6. Effect of Accelerometer Interventions on Physical Activity by Population
 Characteristics 22
 Figure 7. Effect of Accelerometer Interventions on Weight (in Kilograms) by Population
 Characteristics 24
 Figure 8. Effect of Accelerometer Interventions on Physical Activity by Device Role..... 26
 Figure 9. Effect of Accelerometer Interventions on Weight (in Kilograms) by Device Role 27
 Figure 10. Effect of Accelerometer Interventions on Physical Activity by Device Location 32
 Figure 11. Effect of Accelerometer Interventions on Weight (in Kilograms) by
 Device Location 34

APPENDICES

Appendix A. Search Strategies48
 Appendix B. Peer Review Comments Table52
 Appendix C. Characteristics of Included Studies59



THE IMPACT OF WEARABLE MOTION SENSING TECHNOLOGIES ON PHYSICAL ACTIVITY: A SYSTEMATIC REVIEW

INTRODUCTION

Participation in regular physical activity is associated with a wide range of mental and physical health benefits. Patients with diabetes, obesity, and musculoskeletal disease, in particular, derive significant benefits from regular physical activity including favorable effects on blood pressure, lipid profiles, joint swelling and pain, weight control and body fat distribution, and psychological well-being.¹ Despite these known benefits, a large proportion of adults, including military Veterans, are insufficiently active. Less than 50% of Veterans report engaging in physical activity at the level to promote health.² Comparisons also show disparities among VA users and non-users, with higher rates of physical inactivity (20%) observed among Veterans using VA services.² This is a significant concern for VHA clinicians, managers, and policymakers given published reports showing that low aerobic fitness, stemming from inactivity, is associated with increased healthcare costs.^{3,4}

Physical activity has previously been measured using self-reported measures (*eg*, questionnaires, interviews); however there are problems with self-report measures, as these measures do little to motivate or change physical activity behavior and can be subject to reporting bias.⁵⁻⁷ Wearable motion sensing technologies (activity monitors), such as pedometers and accelerometers, have become popular tools to overcome some of these barriers. Epidemiologic and observational studies have used activity monitors to characterize activity intensity and daily activity patterns across diverse samples.⁸⁻¹² Pedometers were the first generation of activity monitors and continue to be widely used to monitor daily ambulation activity, as a tool for prescribing increased mobility (*eg*, daily step targets) and motivating individuals to increase their activity level.^{13,14} However, the reduced measurement properties of pedometers in overweight/obese populations and among individuals with slower ambulation speeds continues to be a limitation.^{15,16}

Although pedometers are a cost-effective tool, they are increasingly being replaced by accelerometers. Accelerometers offer the advantage of assessing all accelerative movements, in all directions, in addition to the ambulation data collected by a pedometer, and can be calibrated to detect differences in intensity.¹³ Data from accelerometers also allow for intervention content to be tailored.⁸ Until recently, research-grade activity monitors have not provided real-time feedback to patients due to a lack of a digital display, or required downloading and interpreting of data using specialized, costly analytic software. However, as the *consumer* market for wearable activity monitors has steadily grown, parallel shifts in the *research* market have also occurred, providing more options for personalized, real-time feedback. Activity monitors used by consumers and researchers now have extensive feedback programs that provide real-time data to the wearer via computer programs and mobile applications.^{17,18} Some of these devices even provide an option to relay this information to a third party such as family, friends, or clinicians. This ability to transmit data to patients' physicians and care teams makes these devices attractive

for clinical applications, although this capability is in the early stages of implementation and evaluation.¹⁸

The Veterans Health Administration (VHA), the largest integrated healthcare system in the United States, recognizes the potential benefit of promoting the use of wearable activity monitors to increase physical activity among Veterans. Before care managers in the VHA embrace this technology and implement these tools in clinical care and wellness programs, the disease states affected, the types of devices studied, and the strategies most likely to affect physical activity outcomes need to be identified. The objective of this review is to synthesize the literature on newer wearable activity devices to determine effectiveness for physical activity outcomes and to describe factors that impact the effectiveness of wearable activity devices.

METHODS

TOPIC DEVELOPMENT

The topic was nominated after a process that included a preliminary review of published peer-reviewed literature and consultation with investigators, Veterans Affairs (VA) and non-VA experts, and key stakeholders. The VA National Center for Health Promotion and Disease Prevention (NCP) nominated this project. The goal of this report, as defined by NCP, is to assess whether the VA should invest in accelerometers and other wearable activity devices as a tool to motivate Veterans to be more physically active.

The final key questions (KQs), developed in consultation with stakeholders, were:

- KQ 1: Among adults, what is the effectiveness of wearable motion sensing technologies (*eg*, activity devices such as accelerometer-based fitness trackers, global positioning systems [GPS]) on:
 - a. Physical activity levels?
 - b. Weight loss or maintenance?
 - c. Patient satisfaction with healthcare?

- KQ 2: Among adults, does the impact of wearable motion sensing technologies (*eg*, activity devices such as accelerometer-based fitness trackers, GPS) vary by:
 - a. Characteristics of the population (overweight/obese/sedentary adults, older adults, healthy volunteers, and individuals with chronic medical illnesses)?
 - b. Type of adjunctive interventions (does the activity device play a major or minor role)?
 - c. Adherence to use of the device?
 - d. Characteristics of the device (body location—waist, arm, wrist, or multisite)

We followed a standard protocol for this review, and each step was pilot tested to train and calibrate study investigators. The PROSPERO registration number is CRD42015017343.

SEARCH STRATEGY

In consultation with an expert librarian, we searched MEDLINE (via PubMed), Embase, CINAHL, SPORTDiscus, and Cochrane CENTRAL from January 1, 2000, to January 6, 2015, for peer-reviewed, English-language randomized controlled trials (RCTs). We used Medical Subject Heading (MeSH) terms and selected free-text terms for wearable motion sensing technologies and for activities/outcomes of interest (*eg*, movement, exercise therapy, physical fitness), along with validated search terms for RCTs.¹⁹ The exact search strategies used are provided in Appendix A. We further reviewed the bibliographies of included trials and systematic reviews²⁰⁻²⁶ for missed publications. To assess for possible publication bias, we searched ClinicalTrials.gov (www.clinicaltrials.gov) to identify completed but unpublished studies meeting our eligibility criteria. All citations were imported into 2 electronic databases (for referencing, EndNote[®] Version X5, Thomson Reuters, Philadelphia, PA; for data abstraction, DistillerSR; Evidence Partners Inc., Manotick, ON, Canada).

STUDY SELECTION

Using prespecified inclusion and exclusion criteria (Table 1), 2 trained investigators assessed titles and abstracts for relevance to the KQs. Full-text articles identified by either investigator as potentially relevant were retrieved for further review and examined by 2 investigators against the eligibility criteria. Disagreements on inclusion/exclusion were resolved by discussion or by a third investigator. In addition, trials with 3 or more arms were examined for appropriateness of all arms for inclusion. For example, data from any active arm that did not include a wearable activity device were not abstracted for inclusion in the analysis.

Table 1. Inclusion and Exclusion Criteria

Study Characteristic	Inclusion Criteria	Exclusion Criteria
Population	<ul style="list-style-type: none"> Adults 18 years of age and older 	<ul style="list-style-type: none"> Elite athletes Children Inpatient populations Pregnant women
Interventions	<ul style="list-style-type: none"> Studies will be included if at least one of the groups used wearable activity devices that provide objective feedback on physical activity to the wearer (eg, non-pedometer-based trackers such as accelerometer-based fitness trackers, smartphone applications, GPS-based trackers), alone or in combination with other interventions to enhance physical activity 	<ul style="list-style-type: none"> Pedometer-based (only) studies Non-wearable systems Systems that do not objectively monitor activity Systems that do not provide feedback to the wearer Interventions that use wearable devices to measure the effects of another intervention (eg, drugs) on ability to perform physical activity
Comparators	<ul style="list-style-type: none"> Usual care/standard of care, waitlist control Pedometer-based interventions Other active comparator focused on enhancing physical activity (eg, educational or behavioral interventions) 	<ul style="list-style-type: none"> Validation studies of head-to-head comparisons of different wearable physical activity devices used to assess validity of devices
Outcomes	<ul style="list-style-type: none"> Change in physical activity behavior (eg, total steps, total activity, proportion of participants at activity goal) Change in weight, body mass index, change or proportion of participants at goal Patient satisfaction with healthcare 	<ul style="list-style-type: none"> Any outcomes not listed
Timing	<ul style="list-style-type: none"> ≥3 months postrandomization 	<ul style="list-style-type: none"> <3 months postrandomization

Study Characteristic	Inclusion Criteria	Exclusion Criteria
Setting	<ul style="list-style-type: none"> · Outpatient general medical settings (geriatrics, family medicine, general internal medicine, integrative medicine) · Specialty medical care clinics (eg, orthopedic, rheumatology, endocrine, cardiology) · Community settings 	<ul style="list-style-type: none"> · Intervention delivered primarily in hospital inpatient setting · Studies where monitoring of physical activity is confined to a supervised setting
Study design	<ul style="list-style-type: none"> · RCTs, n>20 	<ul style="list-style-type: none"> · Not a clinical study (eg, editorial, nonsystematic review, letter to the editor, case series) · Prospective and retrospective observational studies · Measurement or validation studies
Publication type	<ul style="list-style-type: none"> · English-language only · Peer-reviewed articles · Published from 2000 forward 	<ul style="list-style-type: none"> · Non-English articles · Abstracts only

Abbreviations: GPS=global positioning system; RCTs=randomized controlled trials

DATA ABSTRACTION

Data from included articles were abstracted into a customized database by a trained investigator and confirmed by a second investigator. Data elements included date of publication, sample size, location of study, intervention/exposure details, descriptors to assess applicability, quality elements, and outcomes. Key population and activity device characteristics abstracted were age, sex, chronic medical illness status, type of wearable activity monitor (eg, brand, location worn on body), type of adjunctive interventions (eg, behavioral weight management strategies, physical activity education) adherence to use of measurement device, and duration and frequency of intervention. Disagreements were resolved by consensus or by obtaining a third investigator’s opinion when consensus could not be reached.

QUALITY ASSESSMENT

We used the key quality criteria described in the Cochrane Collaboration Risk of Bias Tool (CCRB). The CCRBT was designed to evaluate the risk of bias (ROB) in RCTs.²⁷ It evaluates 6 different domains: (1) adequacy of random sequence generation; (2) allocation concealment; (3) blinding of participants and study personnel; (4) incomplete outcome data; (5) reporting bias due to selective outcome reporting; and (6) other forms of bias such as differences in relation to baseline measures, reliable primary outcomes, or protection against contamination. The Cochrane Collaboration provides guidelines to score each item. Each domain is evaluated as low ROB, unclear ROB, or high ROB. To draw conclusions about the overall ROB within trials we summarized assessments across items in the CCRBT and used the approach outlined in Table 2.




Table 2. Approach to Formulating Summary Risk of Bias for Each Outcome Across Domains

Risk of Bias	Interpretation	Criteria
Low ROB	Bias, if present, is unlikely to alter the results seriously	Adequacy of random sequence generation, allocation concealment, and blinding scored as low ROB, and no important concerns related to the other domains
Unclear ROB	ROB that raises some doubts about the results	One or 2 domains are scored “not clear” or “not done”
High ROB	Bias may alter the results seriously	More than 2 domains are scored as “not clear” or “not done”

Abbreviations: ROB=risk of bias

DATA SYNTHESIS

When meta-analysis was feasible, we computed summary estimates of effect. We aggregated outcomes when there were at least 3 studies with the same outcome. For KQ 1, analyses were conducted separately for wearable activity device interventions versus inactive controls (*eg*, waitlist, usual care) and wearable activity device interventions versus active comparators (*eg*, group weight loss, counseling). Three trials had more than one intervention arm.²⁸⁻³⁰ Two of these trials compared different adjunctive interventions to continuous monitoring via accelerometers.^{28,29} For these 2 studies, we selected the intervention conditions with the less-intensive adjuncts (*eg*, monthly counseling vs weekly counseling). The third trial tested the impact of continuous versus intermittent accelerometer feedback.³⁰ For this study, we selected the comparisons between continuous accelerometer use and control, as this was the type of accelerometer use evaluated in all other studies.

Continuous outcomes (difference in change from baseline to follow-up between intervention and control) were analyzed using standardized mean differences (SMDs) for physical activity outcomes and mean differences (MDs) for weight outcomes in a random-effects model with the Knapp-Hartung correction to confidence intervals. The method we used to interpret the SMD as an effect size was as follows: small effect size, SMD=0.2; medium, SMD=0.5; and large, SMD ≥0.80.³¹ We used R (R Foundation for Statistical Computing, Vienna, Austria) with the metafor package³² to calculate the summary estimates of treatment effect. We evaluated for statistical heterogeneity using visual inspection and the I^2 statistic. Publication bias was assessed using findings from the ClinicalTrials.gov search.

We explored potential sources of heterogeneity including characteristics of the population (*eg*, chronic medical illness status), adherence to the use of the wearable activity device, and characteristics of the device defined as location worn on body. We aimed to assess the differential impact of type of adjunctive interventions (*eg*, behavioral weight management intervention, physical activity education, goal-setting) as a source of potential heterogeneity. Because type and quantity of adjunctive interventions varied greatly from study to study, we operationalized this moderator as the role of the wearable activity device (*ie*, major vs minor component of intervention). To be categorized as being a major component of the intervention, the wearable activity device needed to be the central motivational enhancement intervention intended to improve the primary outcome of the study. Other adjunctive interventions might be included but had to play a minor role in enhancing physical activity. To be categorized as being a

minor component of the intervention, the wearable activity device needed to be an integrated component of a suite of other motivation enhancement interventions, such as a structured exercise program, diet or chronic disease counseling/education/monitoring, self-management techniques, or monetary or nonmonetary incentives. Two independent investigators categorized the role of the device, and any discrepancies were reconciled by the co-principal investigators.

If a quantitative synthesis was not feasible, we analyzed the data qualitatively. We gave more weight to the evidence from higher-quality studies with more precise estimates of effect. We focused on documenting and identifying patterns of the intervention across outcome categories. We analyzed potential reasons for inconsistency in treatment effects across studies by evaluating differences in the study population, intervention, comparator, and outcome definitions.

RATING THE BODY OF EVIDENCE

The strength of evidence (SOE) for each KQ was assessed using the approach described in the Agency for Healthcare Research and Quality (AHRQ)'s *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.³³ We focused on the 2 main outcomes of physical activity and weight. The AHRQ approach requires assessment of 4 domains: risk of bias, consistency, directness, and precision. Additional domains are used when appropriate: coherence, dose-response association, impact of plausible residual confounders, strength of association (magnitude of effect), and publication bias. These domains were considered qualitatively, and a summary rating of high, moderate, or low SOE was assigned after discussion by 2 investigators. In some cases, a rating of high, moderate, or low was impossible or imprudent to make. In these situations, a rating of insufficient was assigned. This 4-level rating scale consists of the following definitions:

- High—High confidence that the evidence reflects the true effect. Further research is very unlikely to change our confidence in the estimate of effect.
- Moderate—Moderate confidence that the evidence reflects the true effect. Further research may change our confidence in the estimate of effect and may change the estimate.
- Low—Low confidence that the evidence reflects the true effect. Further research is likely to change the confidence in the estimate of effect and is likely to change the estimate.
- Insufficient—Evidence either is unavailable or does not permit estimation of an effect.

PEER REVIEW


A draft of this report was reviewed by technical experts and clinical leadership. A transcript of their comments and our responses is provided in Appendix B.

RESULTS

LITERATURE FLOW

The literature search (Figure 1), which was limited to RCTs published between January 1, 2000, and January 6, 2015, identified 4787 unique citations from a combined search of MEDLINE (via PubMed; n=3334), Embase (n=839), CINAHL (n=265), SPORTDiscus (n=245), and Cochrane CENTRAL (n=104). After applying inclusion/exclusion criteria at the title-and-abstract screening level, 176 full texts were retrieved for further review. Of these, papers describing 14 unique RCTs were retained for data abstraction.

Figure 1. Literature Flow Chart



CHARACTERISTICS OF INCLUDED STUDIES

Within the 14 included trials, women comprised 62.5% of the populations, median age was 49.7 years (range 28.7 to 79.8 years), and only 4 trials reported race. The majority of trials were conducted in the United States (n=8) and were sponsored either wholly or partially by government agencies (n=9). Study sizes ranged from 20 to 544 participants (median 62), with the majority of studies (n=8) randomizing fewer than 70 participants. All included studies were published in the last 10 years, indicating the relatively new use of these activity devices in studies aimed at promoting physical activity. While we searched for any wearable nonpedometer devices, all identified trials used some form of accelerometer-based motion sensing technology. Characteristics of the individual studies are described in Appendix C. A search of ClinicalTrials.gov identified one completed but unpublished trial that we believe would meet our inclusion criteria (NCT00544245), revealing a small degree of publication bias.

Twelve studies reported on outcomes related to physical activity,^{30,34-44} and 11 on outcomes related to weight.^{28-30,35-41,43} No studies assessed the outcome of satisfaction with healthcare. Four trials were conducted on older adults, 5 on overweight/obese/sedentary adults, 3 on participants with a chronic medical illness, and 2 on healthy volunteers. We did not identify any studies that were specific to Veterans. Comparators were active (*eg*, behavioral counseling) in 3 trials and inactive (*eg*, usual care, waitlist) in 11 trials. The accelerometer, usually worn on the waist (n=8 trials), was a major component of the intervention in 9 trials and a minor component in 5 trials. Only 4 trials examined adherence. Most studies (n=8) were judged to be at high ROB, with 4 studies at unclear ROB, and only 2 at low ROB.

Interventions varied widely. The duration of interventions ranged from 12 to 52 weeks, and the number of planned interactions with participants in the accelerometer conditions ranged from none to 52 weekly contacts. Trials used a wide variety of adjunctive interventions in conjunction with accelerometers, including intensive diet, weight, and physical activity behavioral counseling; tailored written feedback; and web-based supportive educational modules. Selected details of interventions and comparators for each study are provided in Table 3.

Table 3. Description of Interventions and Comparators

Study	Intervention	Comparator
Greene, 2013 ³⁹	6 months of access to online social network to post about weight and physical activity progress + continuous accelerometer use and feedback	Inactive: Printed lifestyle guidelines on diet and exercise
Koizumi, 2009 ⁴²	12 weeks of accelerometer with feedback + goal-setting	Inactive: 12 weeks of blinded accelerometer use

Study	Intervention	Comparator
Luley, 2014 ²⁸	<p>3-arm study (2 interventional):</p> <p><u>Intervention arm 1:</u> 1-time 2-hour instruction on diet and physical activity + 12 months of accelerometer use + 52 weekly individual letters with feedback on weight, diet, and physical activity</p> <p><u>Intervention arm 2:</u> 1-time 2-hour instruction on diet and physical activity + 12 months of accelerometer use + 12 monthly behavioral counseling calls</p>	Inactive: 1-time 1-hour session, consisting of diet education, diet regimen, and physical activity education
Nicklas, 2014 ³⁵	5-month weight loss intervention that included: hypocaloric diet (2 prepared meals a day) + 4 days/week supervised exercise + self-regulatory intervention that involved wearing an accelerometer, documenting activity, and 6 weekly sessions of behavioral counseling	Active: 5-month weight loss intervention consisting of diet education and regimen (individualized hypocaloric intake), physical activity education, structured exercise (supervised treadmill), and in-person counseling
Paschali, 2005 ⁴⁴	12 weeks of continuous accelerometer use and feedback + 4 monthly in-person exercise behavioral counseling sessions + workbook	Active: 12-week blinded accelerometer use with 4 monthly in-person counseling sessions to review exercise diary, a 24-page information book and home-based walking plan with physical activity education, behavioral self-management, weight goal-setting, and chronic disease monitoring
Polzien, 2007 ³⁰	<p>3-arm study (2 interventional):</p> <p><u>Intervention arm 1:</u> <i>Continuous</i> technology-based behavioral weight control program: 7 in-person individualized counseling sessions over 12 weeks + 12 weeks of continuous accelerometer use and feedback</p> <p><u>Intervention arm 2:</u> <i>Intermittent</i> technology-based behavioral weight control program: 7 in-person individualized counseling sessions over 12 weeks + 3 weeks of accelerometer use and feedback over 12 weeks</p>	Active: 7 in-person individualized counseling sessions consisting of diet education, diet regimen (1200 to 1500 kcal/day; dietary fat <20% of total energy intake), physical activity education, and weight goal-setting
Reijonsaari, 2012 ⁴¹	12 months of continuous accelerometer use and feedback + access to telephone counseling (frequency not defined)	Inactive: Explanation of results of physical exams and information on physical activity and occupational healthcare
Shrestha, 2013 ⁴⁰	1-time 1.5-hour lifestyle instruction + 6 months of continuous accelerometer use and feedback	Inactive: Self-directed exercise and/or U.S. Army-mandated physical training



Study	Intervention	Comparator
Shuger, 2011 ²⁹	<p>4-arm study (3 interventional) :</p> <p><u>Intervention arm 1:</u> Group-based behavioral weight loss: 14 group weight loss sessions + 6 individual phone calls + workbook over 9 months</p> <p><u>Intervention arm 2:</u> Accelerometer alone: 9 months of continuous accelerometer use and feedback</p> <p><u>Intervention arm 3:</u> Group-based behavioral weight loss + accelerometer use: 9 months of continuous accelerometer use and feedback + 14 group weight loss sessions + 6 individual phone calls + workbook</p>	Inactive: Self-directed weight loss manual with diet education, physical activity education, and weight goal-setting
Slootmaker, 2009 ⁴³	3 months of continuous accelerometer use + tailored physical activity feedback and motivational tips via web-based portal	Inactive: A single written brochure with brief physical activity recommendations
Tabak, 2014 ³⁴	Self-directed technology-supported care program that included a tailored web-based exercise program, accelerometer-based activity sensor and motivational messaging, COPD self-management module, and as needed web-portal teleconsultation conducted over 9 months	Inactive: Usual care
Thompson, 2014 ³⁸	12 weeks of continuous accelerometer use and feedback + weekly brief counseling sessions on increasing activity + treadmill desk	Inactive: 12 weeks of blinded accelerometer use
Thompson, 2014 ³⁶	24 weeks of continuous accelerometer use and feedback + weekly brief telephone counseling sessions focused on accelerometer feedback + 6 in-person brief counseling sessions	Inactive: 24 weeks of blinded accelerometer use
Wijsman, 2013 ³⁷	12 weeks of continuous accelerometer use and feedback + personal website + personal e-coach who gives updates on activity status and advice via web portal	Inactive: 3-month waitlist control

Abbreviation: COPD=chronic obstructive pulmonary disease

KQ 1: AMONG ADULTS, WHAT IS THE EFFECTIVENESS OF WEARABLE MOTION SENSING TECHNOLOGIES ON: (A) PHYSICAL ACTIVITY LEVELS; (B) WEIGHT LOSS OR MAINTENANCE; (C) PATIENT SATISFACTION WITH HEALTHCARE?

Key Points

- We identified 12 studies (2 judged to be at low ROB, 2 at unclear ROB, and 8 at high ROB) that met eligibility criteria comparing an accelerometer intervention for increasing physical activity outcomes. The majority of the included studies (n=9) compared the accelerometer device against a weak inactive comparator consisting of usual care, waitlist controls, or one-time educational interventions.
- The overall summary estimate (SMD 0.26; 95% CI 0.04 to 0.49) demonstrated a small positive effect of accelerometer interventions on physical activity levels that was statistically significant. When stratified by comparator, a small statistically significant effect (SMD 0.29; 95% CI 0.03 to 0.55) was found for accelerometer interventions on the outcome of increasing physical activity when compared to an inactive control. An effect of smaller magnitude but same direction was found for the active comparator group, but the pooled estimate was not statistically significant (SMD 0.17; 95% CI -1.09 to 1.43).
- Eleven trials (2 judged to be at low ROB, 3 at unclear ROB, and 6 at high ROB) examined the impact of accelerometer interventions on weight loss or maintenance. Nine of these studies used a weak/inactive comparator. The overall pooled estimate demonstrated a small but statistically significant effect (MD -1.65 kg; 95% CI -3.03 to -0.28) of accelerometers on weight loss. Stratified results by inactive comparator mirrored the overall results but were not statistically significant (MD -1.44 kg; 95% CI -3.08 to 0.19). Only 2 studies used active comparators; both demonstrated a positive trend of weight loss, but only one study was statistically significant.
- The large number of trials at high or unclear ROB, small number of participants in the majority of the trials, and moderate to high heterogeneity across summary estimates limits the strength of evidence on the impact of accelerometers on physical activity levels and weight loss or maintenance.

Physical Activity (KQ 1a)


Synthesis of Findings

Twelve studies, (2 judged to be at low ROB,^{37,41} 2 at unclear ROB,^{34,38} and 8 at high ROB^{30,35,36,39,40,42-44}) examined the impact of accelerometers on the outcome of physical activity. There was substantial variability in the mode and metrics of scales used to measure physical activity. Therefore, all summary estimates were calculated as SMDs. We stratified results by comparator type (inactive vs active) and present stratified and overall pooled estimates. The inactive comparator group contained 9 studies (1279 participants; 2 studies judged to be at low ROB,^{37,41} 5 at high ROB,^{36,39,40,42,43} and 2 at unclear ROB^{34,38}). The active comparator group contained only 3 small studies, all judged to be at high ROB (n=109 participants).^{30,35,44}

Figure 2 shows the forest plot of the meta-analysis examining the effect of accelerometers on physical activity with an overall pooled estimate and stratified pooled estimates by inactive and active comparator subgroups. The overall pooled estimate indicated a small, statistically significant effect for interventions using accelerometers to increase physical activity (SMD 0.26; 95% CI 0.04 to 0.49) with a high amount of heterogeneity ($I^2=64.7%$). A similar small effect was found for interventions using accelerometers to increase physical activity when compared to an inactive comparator (SMD 0.29; 95% CI 0.03 to 0.55). This summary estimate had high heterogeneity ($I^2=70.3%$). The high heterogeneity in these pooled estimates may be driven by several intervention characteristics including the duration of the interventions and the intensity of adjunctive interventions (eg, accelerometer only vs brief device-driven feedback vs intensive device-driven behavioral counseling and feedback), which underscore the overall heterogeneity across these intervention approaches.

There was also a very small positive overall effect (SMD 0.17; 95% CI -1.09 to 1.43) that was not statistically significant for accelerometer devices when compared to an active comparator. This summary estimate had moderate heterogeneity ($I^2=52.3%$). All of the studies included in this analysis were judged to be at high ROB.

Figure 2. Effect of Accelerometer Interventions on Physical Activity



Abbreviations: CI=confidence interval; SMD=standardized mean difference

Summary of Findings

We identified 12 studies that assessed the impact of accelerometer interventions on physical activity. Nine of these used weak controls consisting of usual care, waitlists, or one-time educational instruction and were categorized as using inactive comparators. The other 3 studies used more robust comparators and were classified as having active comparators. Compared to inactive control, the impact of accelerometer interventions on physical activity was similar to that observed in the overall summary estimate, favoring a small effect of accelerometer interventions on physical activity. Both the stratified and overall summary estimates displayed high heterogeneity as assessed by I^2 values >50%.

While the 3 trials with active comparators demonstrated a positive trend of increasing physical activity, the negligible pooled summary estimate (SMD 0.17; 95% CI -1.09 to 1.43) was not statistically significant. When compared to the inactive control summary estimate, these findings suggest that accelerometers may not have as substantial an impact on increasing physical activity when compared to a more robust and active comparator. However, the small number of studies, small sample sizes within these studies, and moderate to high heterogeneity limit conclusions that may be drawn.

Weight Loss or Maintenance (KQ 1b)

Synthesis of Findings


Eleven trials (2 judged to be at low ROB,^{37,41} 3 at unclear ROB,^{28,29,38} and 6 at high ROB^{30,35,36,39,40,43}) examined the impact of accelerometers on weight loss or maintenance. All but one study reported weight changes in kilograms (kg). The other study³⁹ reported weight changes in pounds, which we converted to kilograms. Thus, all summary estimates were calculated as changes in weight expressed in kilograms. We stratified results by comparator type (inactive vs active) and present stratified and overall pooled estimates. The inactive comparator group contained 9 studies (n=1505 participants; 4 studies judged to be at high ROB,^{36,39,40,43} 3 at unclear ROB,^{28,29,38} and 2 at low ROB^{37,41}). The active comparator group contained only 2 small studies judged to be at high ROB (n=79 participants).^{30,35}

Figure 3 shows the forest plot of the meta-analysis examining the effect of the accelerometer interventions on weight loss or maintenance. Only the inactive comparator group had sufficient studies to be pooled. Compared to inactive controls, the impact of accelerometer interventions on weight loss was similar to that observed in the overall summary estimate (MD -1.44 vs -1.65 kg, respectively). Thus, both the overall and stratified estimate favored a small impact of accelerometer interventions on weight loss. However, the inactive pooled estimate was not statistically significant. Both the stratified and overall summary estimates displayed high heterogeneity as assessed by I^2 values >80%. High heterogeneity in these pooled estimates is likely driven by several intervention characteristics including the duration of the interventions and intensity of adjunctive interventions (eg, accelerometer only vs brief device-driven feedback vs intensive device-driven behavioral counseling and feedback) and underscores the overall heterogeneity across these intervention approaches.

Two small trials judged to be at high ROB compared interventions that used accelerometers with active comparators.^{30,35} While both trials demonstrated a positive trend of weight loss (MD range

-3.60 to -2.10), only one study³⁵ was statistically significant. Differences in results are likely attributable to differences in the intensity of other intervention strategies used in conjunction with accelerometer use and in the duration of the interventions. The study with statistically significant findings³⁵ used structured and supervised exercise training, meal preparation twice daily, and behavior counseling delivered over 5 months. The study that did not produce statistically significant results was only 12 weeks long and used behavioral counseling only in conjunction with accelerometer use.³⁰

Figure 3. Effect of Accelerometer Interventions on Weight Loss or Maintenance



Abbreviations: CI=confidence interval; MD=mean difference

Summary of Findings

We identified 11 studies that assessed the impact of accelerometer interventions on weight loss or maintenance. Nine of these used weak control conditions consisting of usual care, waitlists, or one-time educational instruction, and were categorized as using inactive comparators. Two studies used more robust comparators and were classified as having active comparators. While both trials with active comparators demonstrated a positive trend of weight loss (MD range -3.60 to -2.10), only one study was statistically significant. Differences in results are likely attributable to differences in the intensity of other intervention strategies used in conjunction with accelerometer use and in the duration of interventions. Compared to inactive controls, the impact of accelerometer interventions on weight loss was similar to that observed in the overall summary estimate, favoring a small impact of accelerometer interventions on weight loss. However, the inactive pooled estimate was not statistically significant. Both the stratified and overall summary estimates displayed high heterogeneity as assessed by I^2 values >80%. High

heterogeneity in these pooled estimates are likely driven by several intervention and population characteristics that underscore the overall variability across these studies.

In the active comparator group, most studies demonstrated a small weight loss that ranged from -0.36 to -2.65 kg. Yet, one study demonstrated a large weight loss of -8 kg (95% CI -10.41 to -5.59).²⁸ This trial evaluated the effects of telemonitoring for management of obesity over 12 months in patients with established metabolic syndrome. Both the control and intervention groups received dietary instruction at the beginning of the trial. The intervention group also received accelerometers and feedback via monthly behavioral counseling calls. Study authors posited that the magnitude of weight loss seen was secondary to the design of the intervention. The active interventions included key strategies associated with successful weight loss such as self-monitoring, counselor feedback and communication, social support, a structured program, and the use of an individually tailored program. Indeed, other studies included in the pooled analysis that included tailored feedback delivered by a coach or device were more likely to be associated with statistically significant weight loss.^{36,37}

Patient Satisfaction (KQ 1c)


No studies reported on this outcome.

Quality of Evidence for KQ 1

Risk of Bias

Figure 4 presents a summary of the evaluation of the ROB, which shows an ROB graph with review authors’ judgments about each ROB item presented as percentages across all included studies.

Figure 4. Risk of Bias Graph^a



^a For the overall score, low ROB required random sequencing, allocation concealment, and blinding in order to be scored low risk with no other important concerns; unclear ROB was assigned if 1 or 2 domains were scored not clear or not done; high ROB was assigned if >2 domains were scored not clear or not done.

Abbreviation: PA=physical activity

Selection Bias

Random Sequence Generation

Across all included studies, treatment allocation was described as random, and no studies were judged to be at high ROB in this domain. However, 7 of the 14 trials (50.0%) did not give details about the method for generating the random sequence, resulting in a rating of unclear ROB.

Allocation Concealment

In 5 of the 14 trials (35.7%), methods for allocation concealment were described in sufficient detail to determine whether the intervention allocation could have been foreseen in advance of or during enrollment, resulting in a judgment of low ROB. In the majority of trials (9 of 14 [64.3%]), there was an unclear ROB due to inadequate detail about allocation concealment provided by authors.

Performance Bias

For the outcome of physical activity, the blinding of participants and personnel was highly variable, as there are challenges to blinding physical activity interventions. Of the 12 trials measuring physical activity as an outcome, only 2 were judged to be at low ROB as a result of adequate reporting of blinding. In 7 of 12 trials (58.3%), there was an unclear ROB due to inadequate information regarding blinding. One study was judged to be at high ROB due to lack of blinding.

Similar to physical activity outcomes, for the weight outcome, the blinding of participants and personnel was highly variable, as there are challenges to blinding weight change interventions. Of the 11 trials measuring weight change as an outcome, 4 (36.4%) were judged to be at low ROB as a result of adequate reporting of blinding. In 6 of 11 trials (54.5%), there was an unclear ROB due to inadequate information regarding blinding. One study was judged to be at high ROB due to lack of blinding.

Satisfaction with healthcare was not identified as an outcome in any of the included studies. Therefore, no ROB is reported for this outcome.

Detection Bias

For the outcome of physical activity, in 5 of the 12 trials (41.7%) there was sufficient information provided by the authors regarding outcome blinding assessment to judge this domain as low ROB. In the remaining trials (7 of 12 [58.3%]), insufficient information regarding outcome blinding assessment was provided by authors, resulting in a judgment of unclear ROB.

For the outcome of weight, 6 of the 11 trials (41.7%) gave sufficient information provided by the study authors about outcome blinding assessment to judge this domain to be at low ROB. In the remaining trials (4 of 11 [36.4%]), insufficient information regarding outcome blinding assessment was provided by authors, resulting in a score of unclear ROB.

Satisfaction with healthcare was not identified as an outcome in any of the included studies. Therefore, no ROB is reported for this outcome.

Attrition Bias

All trials reported the numbers randomized to each group. The majority of trials (13 of 14 [92.9%]) reported complete outcome data that included information on attrition and exclusions from analysis. One trial did not disclose the reason for attrition/exclusion in sufficient detail, resulting in a judgment of high ROB.

Reporting Bias

The majority of trials (12 of 14 [85.7%]) reported details of the measured outcomes sufficient to be judged to be at low ROB.

Other Bias

The majority of trials (11 of 14 [78.6%]) provided sufficient details not to raise concerns about bias of a nature not covered within the other domains mentioned. Two trials did not provide sufficient methodological detail to judge this domain and were given a judgment of unclear ROB, whereas one trial was judged to be at high ROB stemming from use of a questionable analytical approach during the interim analysis and stopping the trial due to insignificant results.

Overall Risk of Bias

Overall ROB was assessed for each included study (Figure 5). The majority of studies (8 of 14 [57.1%]) were judged to be at high ROB, 4 (28.6%) were at unclear ROB, and only 2 studies (14.3%) were judged to be at low ROB.

Figure 5. Risk of Bias Summary: Review Authors' Judgments About Risk of Bias Items for Each Included Study^a

Study	Was the allocation sequence adequately generated?	Was allocation adequately concealed?	Was knowledge of the allocation intervention adequately prevented during the study?(PA)	Was knowledge of the allocation intervention adequately prevented during the study?(Weight)	Was knowledge of the allocation intervention adequately prevented during the study?(Satisfaction)	Was knowledge of the allocation intervention adequately prevented from outcome assessors?(PA)	Was knowledge of the allocation intervention adequately prevented from outcome assessors?(Weight)	Was knowledge of the allocation intervention adequately prevented from outcome assessors?(Satisfaction)	Were incomplete outcome data adequately addressed?	Are reports of the study free of suggestion of selective outcome reporting?	Are reports of the study free from other bias due to problems not covered above?	Overall
Greene 2013	?	+	?	?		?	?		+	+	+	-
Koizumi 2009	?	?	?	?		?	?		+	+	+	-
Luley 2014	?	?	?	?		?	?		+	+	+	?
Nicklas 2014	?	?	?	?		+	+		+	+	+	-
Paschall 2005	+	?	?	?		?	?		+	+	+	-
Polzien 2007	?	?	?	?		?	?		+	+	+	-
Reijonsaari 2012	+	+	-	-		+	+		+	+	+	+
Shrestha 2013	+	?	?	?		?	?		+	+	+	-
Shurger 2011	+	?	?	+		?	+		+	-	+	?
Slootmaker 2009	?	+	?	?		?	?		+	+	+	-
Tabak 2014	+	?	?	?		?	?		-	?	?	?
Thompson 2014a	?	?	?	?		?	?		+	+	?	-
Thompson 2014b	+	+	+	+		+	+		+	+	-	?
Wijisman 2013	+	+	+	+		+	+		+	+	+	+

^a For the overall score, low ROB required random sequencing, allocation concealment, and blinding in order to be scored low risk with no other important concerns; unclear ROB was assigned if 1 or 2 domains were scored not clear or not done; high ROB was assigned if >2 domains were scored not clear or not done.

Abbreviation: PA=physical activity

KQ 2: AMONG ADULTS, DOES THE IMPACT OF WEARABLE MOTION SENSING TECHNOLOGIES VARY BY: (A) CHARACTERISTICS OF THE POPULATION; (B) TYPE OF ADJUNCTIVE INTERVENTIONS; (C) ADHERENCE TO USE OF THE DEVICES; OR (D) CHARACTERISTICS OF THE DEVICE?

Key Points

We explored the variable impact of wearable motion sensing technologies (activity devices) on physical activity and weight by multiple single factors that may contribute to heterogeneity (*ie*, recruited populations, role of device relative to other motivational enhancement strategies, location of device on body as a proxy for ease of use). None of these individual factors was a robust predictor of heterogeneity.

- Across 4 population subgroups (overweight/obese/sedentary, older adults, healthy volunteers, and chronic medical illnesses), there were no major subpopulation differences in the magnitude and direction of effect estimates on physical activity or weight. Only the older adult subgroup for physical activity and the overweight/obese/sedentary subgroups for weight displayed relatively lower heterogeneity.
- The impact of accelerometer interventions was not well-explained by the role of the device, operationalized as major role versus minor role of the accelerometer in the overall intervention approach. Pooled estimates mirrored the results of the overall summary estimate. The direction and magnitude of the effects were similar for both major and minor role groups; subgroups displayed moderate to high heterogeneity.
- Only 4 of 14 studies reported on adherence to the activity device, and these reports largely focused on the process measures related to accelerometer adherence (hours the device was worn, uploads to websites). The role of accelerometer adherence as a moderating factor of outcome remains to be fully examined in the literature.
- The predominant location of accelerometers was on the waist, followed by the arm. The impact of accelerometer interventions was not well explained by activity device characteristics operationalized as location of the device on the body. Pooled results on physical activity and weight were similar to the overall summary estimate.

Variation by Population Characteristics (KQ 2a)

Synthesis of Findings

We classified studies and organized findings by physical activity and weight. While satisfaction with healthcare was also an outcome of interest, no study reported on this outcome. We planned to perform meta-analytic regression to assess the impact of population characteristics on the outcomes of interest, but we had insufficient studies to perform these analyses. Thus, we categorized studies into population subgroups as determined by study trial recruitment criteria (*ie*, overweight/obese/sedentary, older adults, healthy volunteers, and chronic medical illnesses). For each subgroup with 3 or more studies, we calculated summary estimates. Other results are synthesized qualitatively.

Physical Activity

To assess whether the effects of accelerometer interventions vary by population characteristics, we conducted exploratory subgroup analyses based on study eligibility criteria. For the overweight/obese/sedentary population, there were 158 participants in 4 studies. All 4 studies were judged to be at high ROB.^{30,38,40,43} For older adults, there were 393 participants in 4 studies, including one study judged to be at low ROB,³⁷ one at unclear ROB,³⁶ and 2 at high ROB.^{35,42} For healthy volunteers, there were 783 participants in 2 studies, with one study each at low⁴¹ and high³⁹ ROB. For chronic medical illnesses, there were 54 participants in 2 studies, with one study each at unclear³⁴ and high⁴⁴ ROB.


Figure 6 shows the forest plot of the meta-analysis examining the effect of accelerometer interventions on physical activity by the population subgroups. We had sufficient studies to pool effects for 2 population subgroups—the overweight/obese/sedentary population and older adults. Both subgroups displayed a similar magnitude and direction of effect—namely, a small positive effect of accelerometer interventions on increased physical activity—but neither estimate was statistically significant. The pooled SMD was 0.35 (95% CI -0.52 to 1.22) for the overweight/obese/sedentary group and displayed moderate heterogeneity ($I^2=51.3\%$). For the older adult subgroup, the pooled SMD was 0.34 (95% CI -0.11 to 0.80), with low heterogeneity ($I^2=38.0\%$).

Two additional subgroups—healthy volunteers^{39,41} and those with chronic medical illnesses^{34,44}—were evaluated in only 2 studies each and therefore could not be pooled. Results were mixed in the 2 studies in healthy volunteers. The differences between the 2 studies were likely due to differences in the population baseline activity levels and in participant interactions with adjunctive interventions. In a study at low ROB,⁴¹ healthy insurance company employees (n=521 total) receiving a combined accelerometer and counseling intervention did not increase physical activity when compared to a control group (SMD -0.11; 95% CI -0.28 to 0.06). At baseline, the majority of employees in both arms met physical activity guidelines. In a study judged to be at high ROB,³⁹ there was a small effect indicating increased physical activity (SMD 0.30; 95% CI 0.05 to 0.54) in 262 healthy employees and family members of PeaceHealth Oregon who used an intervention containing an accelerometer plus an online social network versus an education-only control group. The level of social interaction was much greater in the intervention using an online social network compared with the intervention with online or telephone counseling.

For the 2 studies in chronic medical illness populations, accelerometer interventions produced no significant effect on physical activity in one study judged to be at unclear ROB³⁴ (SMD 0.00; 95% CI -0.80 to 0.80), and a moderate statistically significant effect on physical activity in the other study, which was at high ROB⁴⁴ (SMD 0.80; 95% CI 0.06 to 1.55). These differences were likely due to differences in populations (chronic obstructive pulmonary disease [COPD]³⁴ vs type 2 diabetes⁴⁴) and interventions, with one intervention more focused on physical activity.⁴⁴ In the COPD study³⁴ 24 participants received either usual care or a telehealth program where an accelerometer plus a smartphone activity coach was one of 4 components. The intervention arm did not increase physical activity; however, adherence to the physical activity portion of the program was low compared with the portion focused on COPD self-management. In the type 2

diabetes study (n=30 participants),⁴⁴ accelerometers plus counseling increased physical activity as compared to counseling alone.

Figure 6. Effect of Accelerometer Interventions on Physical Activity by Population Characteristics



Abbreviations: CI=confidence interval; SMD=standardized mean difference

Weight

For the outcome of weight, we conducted subgroup analyses by the following populations based on study eligibility criteria: overweight/obese/sedentary populations (1170 participants, 5 studies, with 4 at high ROB^{30,38,40,43} and one at unclear ROB²⁹); older adults (315 participants, 3 studies at low,³⁷ high,³⁵ and unclear³⁶ ROB); healthy volunteers (870 participants, 2 studies at high³⁹ and low⁴¹ ROB); and chronic medical illness populations (120 participants, 1 study at unclear ROB²⁸). There were insufficient studies to pool results for healthy volunteers and chronic medical illness populations.


Figure 7 shows the forest plot displaying the effects of accelerometer interventions on weight by the population subgroups. Among overweight/obese/sedentary persons, 5 studies assessed the impact of accelerometer interventions on change in weight in kilograms. Compared with inactive and active control conditions combined, accelerometer interventions resulted in an MD of -1.22 kg (95% CI -2.46 to 0.02). This estimate displayed relatively low heterogeneity ($I^2=26.8%$). In older adults, accelerometer interventions produced a decrease in weight of a similar magnitude. Again, the pooled effect estimate was not statistically significant (MD -1.08 kg; 95% CI -5.22 to

3.06) and displayed high heterogeneity ($I^2=74.4\%$). This heterogeneity likely reflected populations somewhat different in age and other study eligibility criteria (*eg*, also recruiting for other comorbidities such as overweight).

We identified only 2 studies in healthy volunteers,^{39,41} and both showed a small positive effect of accelerometer interventions on weight. In one study at high ROB,³⁹ 262 healthy employees and family members of PeaceHealth Oregon participated in an intervention with accelerometers where the major focus was on developing an online social network for sharing goal-setting and attainment. When compared to control, this intervention produced an MD of -1.63 kg (95% CI -2.48 to -0.78). In the other study, at low ROB,⁴¹ when compared to a control intervention, a combined accelerometer and counseling intervention produced an MD of -0.50 kg (95% CI -1.00 to 0.00) in health insurance company employees. As above, baseline physical activity was higher in the latter study, and the intensity of the social component of the intervention was greater in the former.

One study judged to be at unclear ROB assessed the effect of accelerometer interventions on weight among study participants with chronic medical illnesses.²⁸ Persons with metabolic syndrome were randomized to a one-time education-only control or 12 months of accelerometer use plus weight, diet, and physical activity feedback via monthly telephone calls or weekly letters. More weight loss was observed in the accelerometer condition than in the control arm (MD -8.00 kg; 95% CI -10.41 to -5.59).

Figure 7. Effect of Accelerometer Interventions on Weight (in Kilograms) by Population Characteristics



Abbreviations: CI=confidence interval; MD=mean difference

Summary of Findings

Physical Activity

For subgroups where quantitative analyses were performed (overweight/obese/sedentary and older adults), pooled effect sizes were not statistically significant (SMD 0.35; 95% CI -0.52 to 1.22 and SMD 0.34; 95% CI -0.11 to 0.80), but were of similar magnitude and direction when compared to one another and to the overall summary estimate. For subgroups where only qualitative analyses could be performed (healthy and chronic medical illness), there were 2 studies each with mixed results. For both subpopulations, one study showed a small, statistically significant increase in physical activity, while the other showed little to no effect that was not statistically significant. Thus, taken together, all subgroups support a small positive effect, with no apparent differential effectiveness on physical activity.

Weight

For the outcome of weight loss, there were no major differences in effect sizes across the 4 population subgroups with the exception of the chronic medical illness subgroup, where a single study at high ROB showed a larger effect size (MD -8.00 kg; 95% CI -10.41 to -5.59). This difference might be attributed to the metabolic syndrome population studied, in which counseling convention strongly promotes lifestyle change above pharmacologic interventions.

Additionally, this intervention was focused on weight loss and included counseling and monitoring of nutrition—as well as physical activity and weight—over a 12-month period compared to a weak control of a one-time educational intervention.

Summary

Table 4 provides a summary of findings for KQ 2a.

Table 4. Summary of Findings: Variation by Population Subgroups (KQ 2a)

Outcome	Population			
	Overweight/Obese/Sedentary	Older Adults	Healthy Volunteers	Chronic Medical Illnesses
Physical activity	4 studies: SMD 0.35 (95% CI -0.52 to 1.22)	4 studies: SMD 0.34 (95% CI -0.11 to 0.80)	2 studies: SMD range: -0.11 to 0.30	2 studies: SMD range: 0.00 to 0.80
Weight	5 studies: MD -1.22 kg (95% CI -2.48 to 0.02)	3 studies: MD -1.08 kg (95% CI -5.22 to 3.06)	2 studies: MD range: -1.63 to -0.50 kg	1 study: MD -8.00 kg (95% CI -10.41 to -5.59)

Abbreviations: CI=confidence interval; MD=mean difference; SMD=standardized mean difference

Variation by Activity Device Role (Major vs Minor; KQ 2b)

Synthesis of Findings

We classified studies and organized findings by physical activity and weight. While satisfaction with healthcare was also an outcome of interest, no study reported on this outcome. We categorized studies into subgroups as determined by activity device role (major or minor). A device was determined to have a major role if the device (accelerometer) was the only or central motivational enhancement intervention intended to improve the primary outcome of the study. In this case, other intervention components were centered on the feedback provided by the accelerometer. Other adjunctive interventions may have been included but played a minor role in enhancing physical activity or could not have been conducted without the feedback of accelerometer data. An activity device was determined to have a minor role if the accelerometer was an integrated component of a suite of other motivation enhancement interventions which may have included, but were not limited to, a structured exercise program, diet counseling/education, chronic disease counseling/monitoring, and/or self-management techniques.

Physical Activity


To assess whether the effects of the accelerometer interventions on physical activity vary by activity device role, we conducted an exploratory subgroup analysis by major versus minor role of the device. The major role subgroup contained 8 studies (n=1041 participants), 5 judged to be at high ROB,^{30,40,42-44} one at unclear ROB,³⁶ and 2 at low ROB.^{37,41} The minor role subgroup contained 4 studies (n=374 participants), 3 at high ROB^{35,38,39} and one at unclear ROB.³⁴

Figure 8 shows the forest plot of the meta-analysis examining the effect of accelerometer interventions on physical activity by activity device role. Both subgroups displayed a similarly



small, positive effect size for accelerometer interventions on increased physical activity, but neither group was statistically significant. The pooled SMD was 0.26 (95% CI -0.02 to 0.54) for the major device role subgroup and displayed high heterogeneity ($I^2=71.6\%$). The subgroup including studies where the activity device played a minor role showed a pooled SMD of 0.28 (95% CI -0.43 to 1.00), with moderate heterogeneity ($I^2=43.2\%$).

Figure 8. Effect of Accelerometer Interventions on Physical Activity by Device Role



Abbreviations: CI=confidence interval; SMD=standardized mean difference


Weight

To assess whether the effects of the accelerometer interventions on weight vary by activity device role, we conducted an exploratory subgroup analysis by major role versus minor role of the device. In the major role subgroup, there were 8 studies (n=1174 participants), 3 judged to be at high ROB,^{30,40,43} 2 at low ROB,^{37,41} and 3 at unclear ROB.^{28,29,36} In the minor role subgroup, there were 3 studies (n=410 participants), all at high ROB.^{35,38,39}

Figure 9 shows the forest plot of the meta-analysis examining the effect of the interventions on weight by activity device role. The direction and magnitude of the effects were similar for both groups, favoring a small impact of less than 2 kg of weight loss, but neither group was statistically significant. Compared with inactive and active control conditions combined, the pooled MD was -1.47 kg (95% CI -3.47 to 0.53) for the major role subgroup and displayed high heterogeneity ($I^2=83.1\%$). The minor role subgroup showed a pooled MD of -1.99 kg (95% CI -4.10 to 0.12) and displayed low heterogeneity ($I^2=24.2\%$).



Figure 9. Effect of Accelerometer Interventions on Weight (in Kilograms) by Device Role



Abbreviations: CI=confidence interval; MD=mean difference

Summary of Findings

The impact of accelerometer interventions on physical activity and weight were not well-explained by the role of the activity device, operationalized as major role versus minor role of accelerometer in the overall intervention approach. These estimates mirrored the results of the overall summary estimate. For physical activity, 8 studies were categorized as using accelerometers as the major intervention strategy,^{30,36,37,40-44} and 4 were categorized as using accelerometers as a minor component in the overall intervention approach.^{34,35,38,39} For weight loss, 8 studies were categorized as major^{28-30,36,37,40,41,43} versus 3 categorized as minor.^{35,38,39} The direction and magnitude of the effects were similar for both subgroups, favoring a small positive impact on increasing physical activity and decreasing weight that mirrored the overall summary estimate. Yet neither group was statistically significant. Across physical activity and weight loss, the major subgroups displayed high heterogeneity. High heterogeneity in the major role of device subgroup is likely driven by several intervention characteristics that underscore the overall variability in the intervention approaches included in this subgroup. For example, studies in this group ranged from accelerometer-alone strategies²⁹ to interventions that included in-person behavioral counseling and accelerometer-driven feedback used to enhance the self-monitoring and motivational aspects of device feedback.³⁰ In general, interventions that included an accelerometer as a major component of the intervention and that capitalized on the self-monitoring and tailored device-driven feedback capabilities of these devices were associated with a greater decrease in weight (see Figure 9).²⁸⁻³⁰ The same trend was not seen for greater

increases in physical activity with better implementation of activity device feedback and self-monitoring functionality.

Table 5 provides a summary of findings for KQ 2b.

Table 5. Summary of Findings: Variation by Device Role (KQ 2b)

Outcome	Role of Accelerometer	
	Major Role	Minor Role
Physical activity	SMD 0.26 (95% CI -0.02 to 0.54)	SMD 0.28 (95% CI -0.43 to 1.00)
Weight	MD -1.47 kg (95% CI -3.47 to 0.53)	MD -1.99 kg (95% CI -4.10 to 0.12)

Abbreviations: CI=confidence interval; MD=mean difference; SMD=standardized mean difference

Variation by Adherence to Use of the Activity Device (KQ 2c)

Synthesis of Findings

The primary aim of KQ 2c was to examine accelerometer adherence as a moderator of the outcomes of physical activity, weight loss, and satisfaction with healthcare. Only 4 of 14 studies reported any measure of accelerometer adherence.^{30,35,41,43} Across these 4 studies, adherence to the activity device was defined variably by total arm-band time use, categorical measure of time the device was worn (eg, often vs regularly), percentage of days device was worn at least 10 hours per day, and change in web-based service use at 2 time points. In light of dropout rates (range of 12% to 31%), adherence data were reported only on study completers. While these 4 studies assessed physical activity levels and body weight as outcome measures, only one of these studies examined accelerometer adherence relationship to these outcomes (weight only). Satisfaction with healthcare was not assessed as an outcome in any of these studies. Individual determinants of accelerometer adherence were not examined in any of these studies. The synthesis of findings presented here combines results for physical activity and weight loss outcomes.

A large worksite physical activity promotion study judged to be at low ROB⁴¹ randomized 544 employees to either 12 months of continuous accelerometer use and distance counseling (by phone or the web) or a control group that received baseline fitness results only. Participants in the intervention arm were instructed to enter their physical activity score from the accelerometer manually to the web-based service. Use of the web-based service was assessed as a proxy of accelerometer adherence in this study. Overall, use of the web-based service was low among participants in the intervention arm, averaging just 15 logins during the last 6 months of the trial (0.6 logins per week). Comparisons of the 0- to 6-month versus 6- to 12-month time periods of the study show that use of the web-based service also decreased over the trial for both participants and coaches. Participants entered fewer days of physical activity scores (14% of days vs 9% of days), coaches sent fewer messages to participants (7.2 vs 6.1 messages), and participants sent fewer messages to coaches (4.3 vs 1.7 messages). This study did not examine the impact of accelerometer adherence on physical activity or weight loss outcomes.

A second, smaller worksite physical activity promotion study judged to be at high ROB⁴³ randomized 102 healthy young employees to either 3 months of continuous accelerometer use and tailored web-based support or a control group that received a single brochure with brief activity recommendations at baseline. Participants in the intervention arm were encouraged to



enter their physical activity score from the accelerometer manually to the web-based service. Uploaded accelerometer scores were paired with automatic, tailored physical activity advice and motivational tips. Seventy-three percent of those in the intervention arm reported wearing the accelerometer “regularly” or “often” and accessed the web-based service on average nearly once per week to upload accelerometer data. Fifty-two percent of intervention arm participants set a personal goal using the web-based service. However, 41% of participants found the tailored advice on the website unappealing. Barriers to compliance included a limited menu of preferred personal activities on the web-based service, and physical activity advice not relevant to the participant’s preferred personal activity. This study did not examine the impact of accelerometer adherence on physical activity or weight loss outcomes.

In a small 12-week behavioral weight-loss study judged to be at high ROB,³⁰ 57 healthy adult participants were randomized to: (1) standard in-person weight control program; (2) standard program + continuous technology-based behavioral program (continuous armband-accelerometer wear and feedback); or (3) standard program + intermittent technology-based behavioral program (3 weeks of armband accelerometer wear and feedback). Average total accelerometer wear time was low in both the intermittent and continuous groups (63.8 minutes/week and 71.0 minutes/week, respectively) and was highly variable (range 51 to 350 minutes/week). Adherence to the accelerometer, defined as total arm band time, declined significantly across all 4-week time periods in the continuous technology group. Self-monitoring of exercise also decreased across the intervention period in the standard and intermittent groups. This was the only study to examine the relationship between accelerometer adherence and weight loss. Significant correlations were reported between change in body weight and total accelerometer wear time in both the intermittent ($r=-0.68$, $p<0.01$) and continuous ($r=-0.71$, $p<0.01$) technology arms.

In a pilot study (high ROB) of spontaneous physical activity and weight loss,³⁵ 48 older adults were randomized to a 5-month weight loss intervention that included a hypocaloric diet and aerobic exercise or the same weight loss intervention plus a self-regulatory intervention (which included an accelerometer) to promote spontaneous physical activity. Following post-weight loss assessments at 5 months, both groups transitioned to a self-directed diet and exercise program during a 5-month follow-up period. Adherence to the accelerometer was high (see Table 6). Eighty-five percent of participants ($n=17$) submitted accelerometer data for the entire 10 months of the study, of which 87% wore the accelerometer for the prescribed ≥ 10 hours per day. Eighty-one percent of participants met the daily spontaneous physical activity goal recorded by the accelerometer. Reported barriers to accelerometer wear compliance included device malfunction/battery change (13%), illness/health (9%), forgot to wear the monitor (7%), and too busy/time conflict (7%). The transition from the 5-month, intensive intervention to the self-monitored program also seemed to pose a barrier for some who stopped wearing the monitor (15%). This study did not examine the impact of accelerometer adherence on physical activity or weight loss outcomes.

Summary of Findings

Only 4 of 14 studies provided any data on adherence to the activity device. Although our primary aim was to examine accelerometer adherence as a moderator of physical activity, weight loss, and satisfaction outcomes, none of these studies reported such analyses. Instead, these studies focused largely on the process measures related to accelerometer adherence. Despite the major

role played by the accelerometer in 3 of these studies, a consistent pattern of decline in participants’ use of the device was evident over study duration, and generally low levels of adherence over time were reported. Two studies conducted in workplace settings showed significant participant disengagement expressed in low levels of self-motivated, self-initiated, stated goals to be achieved during the study time period. In summary, we are unable to determine from this literature whether accelerometer adherence has an effect on the outcomes of interest when accelerometers are used as part of behavioral interventions.

Table 6 provides a summary of findings for KQ 2c.

Table 6. Summary of Findings: Accelerometer Adherence Across Individual Studies of Physical Activity and Weight Loss (KQ 2c)

Study	Role of Accelerometer	Accelerometer Adherence Description
Nicklas, 2014 ³⁵	Minor	<ul style="list-style-type: none"> • 85% of participants submitted accelerometer data for entire 10 months • 87% of days completers wore accelerometer at least 10 hours/day • 81% of days spontaneous physical activity goal met • Barriers to compliance: device malfunction/battery change (13%), illness/health (9%), forgot to wear (7%), too busy/time conflict (7%)
Polzien, 2007 ³⁰	Major	<ul style="list-style-type: none"> • Accelerometer wear time was low and highly variable in both continuous and intermittent technology groups • Significant decline in total accelerometer wear time use by continuous technology group across all 4-week time periods • Standard and intermittent groups decreased self-monitoring of exercise across the 12-week intervention period • Longer total accelerometer wear time was significantly associated with greater weight loss in both the continuous and intermittent technology groups
Reijonsaari, 2012 ⁴¹	Major	<ul style="list-style-type: none"> • Use of web-based service decreased with time (comparing 0 to 6-month vs 6- to 12-month study periods) for both participants and coaches • Declines in web-based service across time included entries of participant physical activity scores (14% of days vs 9% of days); coach communication to participants (7.2 vs 6.1 messages); participant communication to coaches (4.3 vs 1.7 messages)
Slootmaker, 2009 ⁴³	Major	<ul style="list-style-type: none"> • 73% of participants wore accelerometer “often” or “regularly” • 52% of participants set personal goal on web-based service • Barriers to compliance: Limited menu of preferred, personal activities on web-based service; physical activity advice not relevant to participant-preferred, personal activity; 41% found advice on website unappealing



Variation by Characteristics of the Activity Device (Body Location; KQ 2d)

Synthesis of Findings

We classified studies and organized findings by physical activity and weight. While satisfaction with healthcare was also an outcome of interest, no study reported on this outcome. We categorized studies into subgroups as determined by characteristics of the activity device, operationalized as location of the device on the body. Locations of interest were waist, arm, wrist, and multisite. One study that did not specify device location was not included. We had sufficient studies to pool effects only for one subgroup (waist); other results are synthesized qualitatively.

Physical Activity

To assess whether the effects of the accelerometer interventions on physical activity vary by device characteristics, we conducted exploratory subgroup analysis by the following device location sites: waist (n=809 participants; 7 studies, 5 judged to be at high ROB,^{35,38,42-44} one at low ROB,⁴¹ and one at unclear ROB³⁶); arm (n=38 participants; 2 studies, both at high ROB^{30,40}); wrist (n=235 participants; one study at low ROB³⁷); and multisite (n=24 participants; one study at unclear ROB³⁴).

Figure 10 shows the forest plot examining the effect of the interventions on physical activity by activity device location. The pooled estimate displayed a weakly positive association regarding effect of location of accelerometer at the waist on increasing physical activity (SMD 0.24; 95% CI -0.15 to 0.63) and moderate heterogeneity ($I^2=62.3%$). This effect was similar in direction and magnitude to the overall summary estimate.

Three additional subgroups—arm, wrist, and multisite—contained fewer than 3 studies and could not be pooled. The 2 studies where the device was worn on the arm^{30,40} showed mixed results, neither of which was statistically significant. Of the 2 studies, one (at high ROB)³⁰ showed a negative impact on physical activity (SMD -0.15; 95% CI -0.78 to 0.49). The second (also at high ROB)⁴⁰ showed a small positive effect on physical activity (SMD 0.60; 95% CI -0.30 to 1.50). The differences in the 2 studies were likely due to differences in the duration of the intervention (12 vs 24 weeks), comparator arms (active vs weak or active vs active), and overall analytic strategy of each study. Of note, both studies had high dropout rates, but only one study specifically stated that they used an intention-to-treat analysis. In Shrestha et al⁴⁰ an accelerometer paired with an interactive website versus usual care did not show a statistically significant mean change in total physical activity between groups. The lack of difference in effect on physical activity was attributed to low initial recruitment (28 participants total) and a high dropout rate (attrition 64% at 6 months). Polzien et al³⁰ used a technology-based program added to an in-person behavioral weight loss program. Although each group showed a significant increase in leisure time physical activity, there was no statistically significant difference between the active intervention arms.


For one study where the accelerometer was worn on the wrist (low ROB),³⁷ the intervention produced a modestly statistically significant increase in physical activity (SMD 0.56; 95% CI 0.30 to 0.82). In this study, 119 adults were randomized to receive a 12-week web-based physical activity program focused on increasing daily activity. The intervention was composed

of 3 elements: an accelerometer-based activity monitor, a personal website, and a personal e-coach who gave feedback about ways to increase physical activity. After 13 weeks, daily physical activity increased in the intervention versus control group (11% vs 5%, $p < 0.11$).

Finally, a pilot study (at unclear ROB)³⁴ showed no effect on physical activity of an accelerometer that could be worn at multiple sites as compared to a usual care group (SMD 0.00; 95% CI -0.80 to 0.80). The intervention arm included 4 modules: a web-based exercise program for home exercising, an activity coach with real-time feedback, self-management of COPD exacerbations via diary on web portal, and teleconsultation. The activity coach consisted of an accelerometer-based sensor and a smartphone that displayed cumulative activity and sent motivational cues based on activity already performed. Objective activity at baseline and 3 months for the intervention group (n=15) was 536.3 (standard error of the mean [SEM] 42.6) and 511.0 (SEM 44.1), respectively. Objective activity at baseline and 3 months for usual care (n=14) was 360.5 (SEM 44.7) and 335.2 (46.3), respectively. The notable baseline difference may be influenced by the poorer clinical characteristics including a higher prevalence of dyspnea among usual care ($p=0.03$) despite computerized randomization distributed in sealed envelopes.

The lack of difference in physical activity based on multisite location is likely secondary to very low use of the exercise module by physiotherapists and low use and adherence to the exercise module and activity coach by participants. Of note, median adherence to prescribed exercise schemes was 21%.

Figure 10. Effect of Accelerometer Interventions on Physical Activity by Device Location



Abbreviations: CI=confidence interval; SMD=standardized mean difference


Weight

To assess whether the effects of the accelerometer interventions on weight vary by device characteristics, we conducted exploratory subgroup analysis by the following device location sites: waist (n=852 participants; 6 studies, 3 judged to be at high ROB,^{35,38,43} one at low ROB,⁴¹ and 2 at unclear ROB^{28,36}); arm (n=157 participants; 3 studies, 2 at high^{30,40} and one at unclear²⁹ ROB); and wrist (n=226 participants; one study at low ROB³⁷).

Figure 11 shows the forest plot of the meta-analysis examining the effect of the interventions on weight by device location. The overall pooled estimate demonstrated a small statistically significant effect on weight (MD -1.65 kg; 95% CI -3.03 to -0.28), with high heterogeneity ($I^2=81\%$). We had sufficient studies to pool effects for 2 subgroups (waist and arm). Both produced effects of the same direction and magnitude, which mirrored those of the overall summary estimate. Compared with control conditions, the pooled MD was -2.01kg (95% CI -4.99 to 0.97) for the waist subgroup and displayed high heterogeneity ($I^2=89.1\%$). High heterogeneity in the waist subgroup is likely driven by differences in intervention approach (self-monitoring vs self-monitoring plus individualized counseling/feedback), intensity (weekly counseling via phone calls or letters vs self-driven interaction with device/website), and length of follow up (range 3 to 12 months). The arm location subgroup showed a similar pooled MD of -2.08 kg but was statistically significant (95% CI -4.13 to -0.02), with low heterogeneity ($I^2=0.00\%$).

One additional subgroup, wrist, contained fewer than 3 studies and could not be pooled but demonstrated small positive impacts on weight loss. In the one study (low ROB) where the accelerometer was worn on the wrist,³⁷ the intervention produced a small impact on weight (MD -0.67 kg; 95% CI -1.33 to -0.01). In this study, 119 adults were randomized to receive either a 12-week web-based physical activity program focused on increasing daily activity or a 3-month waitlist control. The intervention was composed of 3 elements: an accelerometer-based activity monitor, a personal website, and a personal e-coach who gave feedback about ways to increase physical activity.

Figure 11. Effect of Accelerometer Interventions on Weight (in Kilograms) by Device Location



Abbreviations: CI=confidence interval; MD=mean difference

Summary of Findings

The predominant location of accelerometers was the waist, followed by the arm. The impact of accelerometer interventions was not well explained by where the device was worn on the body; pooled estimates mirrored those of the overall summary estimate. For physical activity, activity devices worn on the waist (n=7 studies) produced effects that ranged from negative and not statically significant to modest positive effects that were statistically significant. Of note, only one small (n=20) study of a waist-worn device, judged to be at high ROB, produced positive and statistically significant effects on physical activity.³⁸ Activity devices worn on the arm (n=2 studies) also displayed mixed results. One study (at low ROB) including a wrist-worn device³⁷ produced a positive and statistically significant impact on physical activity, while a study that allowed multisite use³⁴ showed no impact on physical activity.

For weight loss, activity devices worn on the waist (n=6 studies) produced effects that ranged from no effect to positive and statistically significant effects. High heterogeneity in the waist subgroup is likely driven by differences in intervention approach (self-monitoring vs self-monitoring plus individualized counseling/feedback), intensity (weekly counseling via phone calls or letters vs self-driven interaction with device/website), and differences in length of follow up (range 3 to 12 months). Activity devices worn on the arm (n=3 studies) displayed modestly positive effects on weight that were statistically significant and displayed low heterogeneity. One study including a wrist-worn device produced a small positive and statistically significant effect

on weight. This was the only device location that produced positive and statistically significant effects on both physical activity and weight, but this group included only one study (low ROB).³⁷

Table 7 provides a summary of findings for KQ 2d.

Table 7. Summary of Findings: Variation by Characteristics of the Device (KQ 2d)

Outcome	Location of Accelerometer			
	Waist	Arm	Wrist	Multisite
Physical activity	7 studies: SMD 0.24 (95% CI -0.15 to 0.63)	2 studies: SMD range: -0.15 to 0.60	1 study: SMD 0.56 (95% CI 0.30 to 0.82)	1 study: SMD 0.00 (95% CI -0.80 to 0.80)
Weight	6 studies: MD -2.01 (95% CI -4.99 to 0.97)	3 studies: MD -2.08 (95% CI -4.13 to -0.02)	1 study MD -0.67 (95% CI -1.33 to -0.01)	No studies

Abbreviations: CI=confidence interval; MD=mean difference; SMD=standardized mean difference

Quality of Evidence for KQ 2

The same 14 studies were included for KQs 1 and 2. The quality of evidence is discussed above, under KQ 1.



SUMMARY AND DISCUSSION

Physical activity is universally endorsed as a cornerstone for maintaining optimal physical and cognitive health.^{45,46} Increased physical activity is also considered an important therapeutic modality for patients with many chronic conditions including diabetes,⁴⁷ arthritis,⁴⁸ various cancers,⁴⁹ obesity,⁵⁰ and depression.⁵¹ Despite the proven benefits and widespread public health and clinical calls to increase physical activity, sedentary behavior has proven difficult to change.

Self-monitoring is a key behavioral strategy to increase physical activity, and objective self-monitoring, as opposed to self-report, is considered the gold standard.⁵² Pedometers have been found to be associated with significant increases in physical activity and significant decreases in both body mass index and blood pressure.⁵³ Although pedometers are a cost-effective tool, they are increasingly being replaced by accelerometers. Accelerometers offer the advantage of assessing accelerative movements, in all directions. In addition, accelerometers can detect differences in intensity¹³ and can be tethered with other electronic devices to provide real-time, tailored feedback. Therefore this review sought to identify and summarize the evidence on the use of accelerometers on 2 major outcomes associated with many chronic conditions: physical activity and weight loss or maintenance. In addition, we sought to examine the effects of accelerometer use on satisfaction with healthcare.

We identified 14 unique RCTs that assessed the impact of accelerometers on physical activity and weight loss or maintenance. All included studies were published in the last 10 years, indicating the relatively new use of motion sensing devices in studies aimed at promoting physical activity or weight loss. Although we searched for wearable motion sensing technologies in general, no studies included newer wearable motion sensing technologies (*eg*, GPS, hand gesture, eye gesture, hand swipe) beyond an accelerometer. Twelve studies assessed the impact of accelerometers on physical activity and 11 on weight. No identified studies reported on the impact of accelerometers on satisfaction with healthcare. Four trials explicitly recruited older adults, 5 recruited obese/sedentary adults, 3 recruited participants with a chronic medical illness (diabetes, metabolic syndrome, COPD), and 2 recruited healthy volunteers. No studies specifically recruited Veterans.

STRENGTH OF EVIDENCE

Table 8 presents an overview of findings and strength of evidence (SOE) by major outcomes. We found moderate SOE for small increases in physical activity level (SMD 0.29; 95% CI 0.03 to 0.55) and small decreases in weight (MD -1.44 kg; 95% CI -3.08 to 0.19) over 3- to 12-month periods (median 5.5 months) when interventions that integrated accelerometers were compared with weak or inactive controls. We found low SOE for very small increases in physical activity levels (SMD 0.17; 95% CI -1.09 to 1.43) and small decreases in weight (MD range -3.60 to -2.10 kg) for accelerometer interventions compared with active control conditions.

Table 8. Summary of Intervention Effects and Strength of Evidence Ratings

Outcome and Comparison	Number of Studies (Subjects)	Domains and Ratings Pertaining to SOE	SOE and Summary Effect Estimate (95% CI)
Physical Activity: Inactive Comparator	9 (1279)	<p>Risk of Bias: All RCTs, but some limitations due to poor study quality. Majority of RCTs were judged to be at high (n=5) or unclear (n=2) ROB. This domain was not rated as high ROB because 2 of the larger studies were at low ROB, and a large proportion (59.1%) of the weight for the subgroup summary estimate was generated from these 2 studies.</p>	Moderate SMD 0.29 (0.03 to 0.55)
		<p>Consistency: Some inconsistency. The individual study estimates for the most part had the same direction of effect as the subgroup summary estimate. One study, which was the largest and judged to be at low ROB, had an individual effect size on the opposite side of the null value. The magnitude of effect sizes varied across individual studies (SMD range -0.11 to 1.22).</p>	
		<p>Directness: Direct</p>	
		<p>Precision: Some imprecision. Most of the included studies had small sample sizes. The subgroup summary estimate was precise and similar to the overall summary estimate. The impact of bias on heterogeneity was inconclusive, as there was no consistency in the relationship between studies of variable ROB and direction of effect, and the effect estimate from the largest study, scored at low ROB, was on the opposite side of the null.</p>	
Physical Activity: Active Comparator	3 (109)	<p>Risk of Bias: All RCTs, but all studies judged to be at high ROB.</p>	Low SMD 0.17 (-1.09 to 1.43)
		<p>Consistency: Minor inconsistency. Two of the 3 individual estimates had a similar effect size, but the range of individual effect sizes varied substantially from -0.05 to 0.80.</p>	
		<p>Directness: Direct</p>	
		<p>Precision: Imprecise. All of the included studies had small sample sizes. The subgroup summary estimate and the overall summary estimate had wide confidence intervals. The impact of bias on heterogeneity could not be ruled out, as one of the individual estimates was on the opposite side of the null and was judged to be at high ROB. This may have influenced heterogeneity and subsequently the overall summary estimate confidence interval.</p>	

Outcome and Comparison	Number of Studies (Subjects)	Domains and Ratings Pertaining to SOE	SOE and Summary Effect Estimate (95% CI)
Weight: Inactive Comparator	9 (1505)	Risk of Bias: All RCTs, but some limitations due to poor study quality. Most studies were judged to be at high (n=4) or unclear (n=3) ROB. This domain was not rated as high ROB because 2 of the larger studies were scored at low ROB, and a large proportion (49.6%) of the weight for the subgroup summary estimate was generated from these 2 studies.	Moderate MD -1.44 kg (-3.08 to 0.19)
		Consistency: Consistent in direction and magnitude. The individual estimates for the active comparator group were on the same side of the null value. Aside from one study, the MDs were consistent in magnitude when compared to the subgroup summary estimate.	
		Directness: Direct	
		Precision: Precise. A few of the included studies had small sample sizes, which contributed to imprecision. The subgroup summary estimate was precise and similar to the overall summary estimate. The impact of bias on heterogeneity is possible, as one study judged to be at unclear ROB was outside the MD range of other included studies in this subgroup.	
Weight: Active Comparator	2 (79)	Risk of Bias: Both studies RCTs, but both judged to be at high ROB.	Low MD range: -3.60 kg (-5.82 to -1.38) to -2.10 kg (-4.30 to 0.10)
		Consistency: Minor inconsistency. The individual studies had similar MDs on the same side of the null value.	
		Directness: Direct	
		Precision: Some imprecision. Both included studies had small sample sizes.	

Abbreviations: CI=confidence interval; MD=mean difference; SMD=standardized mean difference; RCTs=randomized controlled trials; ROB=risk of bias; SOE=strength of evidence

SUMMARY OF EVIDENCE BY KEY QUESTION

Overall, we found a small positive effect of accelerometers on increasing physical activity and decreasing weight. Twelve studies provided estimates for the effect of accelerometers on level of physical activity. Pooled estimates demonstrated a small and favorable impact on increasing physical activity levels (SMD 0.26; 95% CI 0.04 to 0.49) with moderate heterogeneity ($I^2=64.7\%$). In stratified analysis by type of comparator, the effects versus the inactive comparator were similar; however, no significant effect was found when accelerometers were compared with an active comparator. A similar pattern was observed for the 11 studies that provided estimates for the effect of accelerometers on weight loss. The overall pooled estimates demonstrated a small and favorable impact on weight (MD -1.65 kg; 95% CI -3.03 to -0.28) but exhibited high heterogeneity ($I^2=81\%$). In stratified analysis by type of comparator, effects were muted when accelerometer interventions were compared with more robust controls than with weaker controls. The moderate to high heterogeneity in these pooled estimates may be driven by

several intervention characteristics, including the duration of the interventions, intensity of adjunctive interventions (*eg*, accelerometer only vs brief device-driven feedback vs intensive device-driven behavioral counseling and feedback) and study design features that resulted in most studies being judged to be at high or unclear ROB. For physical activity outcomes, an additional source of heterogeneity may stem from the substantial variability in the mode (self-report vs objectively measured) and metrics (step counts per day vs minutes per day active) used to measure this outcome.

We also sought to explore the variable impact of wearable activity devices on physical activity and weight by multiple single factors that may contribute to heterogeneity (recruited populations, role of device relative to other motivational enhancement strategies, location of device on body as a proxy for ease of use). None of these individual factors was a robust predictor of heterogeneity. Thus, the variability in pooled estimates is likely due to a combination of factors related to underlying differences in populations, comparators, interventions, and study design and quality issues. However, other quantitative patterns were observed. In general, interventions that capitalized on the self-monitoring and tailored activity device-driven feedback capabilities of these devices were associated with greater decreases in weight loss. Effects were even greater when these strategies were paired with behavioral counseling focused on device feedback. The same trend was not consistently seen for greater increases in physical activity with better implementation of device feedback and self-monitoring functionality. Lastly, while we sought to explore the role of adherence to the activity device as a moderator of effects, only 4 studies provided any data on adherence to the device, and none of these studies reported such analyses. Instead, these studies focused largely on the process measures related to accelerometer adherence. In general, these studies reported a consistent pattern of decline in participants' use of activity devices over time.

CLINICAL AND POLICY IMPLICATIONS

Commercially available wearable activity devices have surged in popularity in recent years, including accelerometers marketed for their ability to assist individuals who want to increase their physical activity. Accelerometers have proven to be a valid and reliable means of tracking step counts and activity levels; however, their effectiveness as an agent of behavior change has been less clear. Also it is unclear whether accelerometers offer additional advantages over pedometers as a means of self-monitoring physical activity.

The results of this review provide important new information by demonstrating a small positive effect of wearable accelerometers on increasing physical activity and decreasing weight. These findings are important for both clinician and policymakers who may be considering programs to encourage more widespread use of these technologies as a public health strategy to improve health and well-being. Although we found small but significant effects related to activity device use, the heterogeneity of the included studies precludes firm conclusions about how they can be deployed most effectively in clinical practice as a tool to facilitate increased physical activity. Furthermore, we did not find any studies that sought to integrate physical activity data from wearable accelerometers into the patient's medical records to facilitate ongoing primary care and chronic disease management. Such research could be of real value to clinicians and policymakers.

No studies actively recruited Veterans and, germane to the VA healthcare system, we found surprisingly few trials (n=3) that focused on chronic disease populations. Use and effectiveness of wearable accelerometers may differ among participants who are motivated to use these devices to achieve different goals; for example, those who are trying to increase physical activity to reduce pain from osteoarthritis compared to participants whose goal is to lose weight. Thus, it is unclear if these devices should be offered to all patients or a subset of patients who may derive the most benefits from increased physical activity.

For accelerometers to function optimally, patients need to wear the devices and understand the information they receive from them. In this review, we found limited data on activity device adherence. Other research suggests that more than half of individuals who purchase a wearable activity device stop using it; of these, one-third stop in the first months.⁵⁴ Future research should (1) measure how often participants wear accelerometers; (2) measure how participants interact with their generated data; and (3) explore facilitators and barriers to adoption of wearable technologies. This would be valuable for clinicians, policymakers, and organizations by helping programs target individuals who are most likely to wear and therefore benefit from these devices. Last, accelerometers are more costly than traditional pedometers, which may impact decisions regarding purchase and usage. In this review, we were unable to assess whether accelerometers increase physical activity beyond pedometers.

LIMITATIONS

Our review has a number of strengths, including a protocol-driven design, a comprehensive search, and careful quality assessment. We conducted both quantitative and qualitative synthesis when possible. Our review—and the literature—have limitations: the number of studies is small; many had design limitations (8 of 14 were judged to be at high ROB); the range of interventions evaluated was diverse; and the number and reporting of studies precluded any analyses of variability in accelerometers by more than one variable at a time. Our review was limited to English-language publications, but the likelihood of identifying relevant data unavailable from English-language sources is low.

The small sample sizes of most included trials and the populations recruited also limited our findings and may have resulted in a type II error as a result. No studies actively recruited Veterans, yet we did identify 3 studies among those with chronic medical illnesses, 5 studies among overweight/obese/sedentary populations, and 4 studies among older adults. It is likely that the results of these studies are highly applicable to the VA, because these conditions and populations are common among the patients seen in the VA healthcare system. However, having so few studies with large sample sizes leaves unanswered questions of feasibility of integrating accelerometers into clinics and healthcare systems with large, heterogeneous patient populations and multiple providers.

Many of our pooled analyses suggested moderate to high heterogeneity. While we conducted subgroup analysis to explore multiple single factors that may contribute to heterogeneity (recruited populations, role of activity devices relative to other motivational enhancement strategies, location of device on body as a proxy for ease of use), none of these individual factors was a robust predictor of heterogeneity. Thus, the observed heterogeneity is likely attributable to a combination of factors that relate to underlying differences in populations, comparators,

interventions, and study design and quality issues. The overall low number and small size of included trials per outcome precluded us from conducting multivariable analyses.

RESEARCH GAPS/FUTURE RESEARCH

This comprehensive review of the literature identified several gaps in the current evidence that warrant future investigation. We used the framework recommended by Robinson et al⁵⁵ to identify gaps in evidence and classify why these gaps exist (Table 9). This approach considers the PICOTS (population, intervention, comparator, outcome, timing, and setting) to identify gaps and classifies them as due to (1) low strength of evidence or imprecise information, (2) biased information, (3) inconsistency or unknown consistency, and (4) not the right information. VA and other healthcare systems should consider their clinical and policy needs when deciding whether to invest in research to address gaps in evidence.

Table 9. Evidence Gaps and Future Research

Evidence Gap	Reason	Type of Studies to Consider
Population		
<ul style="list-style-type: none"> · Absence of trials that actively recruited Veterans · Limited studies (n=3) among those with chronic medical illnesses 	Insufficient information	<ul style="list-style-type: none"> · RCTs · Quasi-experimental studies · Prospective cohort studies
Interventions		
<ul style="list-style-type: none"> · What are the optimal adjunctive interventions that enhance functionality of accelerometers in motivating behavior change? · How does intensity or dose of adjunctive interventions enhance functionality of accelerometers in motivating behavior change? · How is adherence to use of the activity device associated with key outcomes? Are there strategies to enhance adherence? · How do aspects of the activity device (location on body, ease of use, ability to pair with other technologies) impact physical activity and weight? · Is accelerometer-based feedback inferior, equivalent, or superior to pedometer-based feedback for increases in physical activity or decreases in weight? 	Insufficient information	<ul style="list-style-type: none"> · RCTs of head-to-head comparisons of different types of adjunctive intervention packages · Stepped and adaptive trial designs
Comparators		
<ul style="list-style-type: none"> · Relatively few studies that used active or more robust comparators · No studies that assessed accelerometers vs pedometers 	Insufficient information	<ul style="list-style-type: none"> · RCTs · RCTs of head-to-head comparisons of accelerometers and pedometers
Outcomes		
Uncertain effects on: <ul style="list-style-type: none"> · Patient satisfaction with healthcare Limited information on:	Insufficient information	<ul style="list-style-type: none"> · RCTs · Prospective cohort studies · Non-randomized controlled

Evidence Gap	Reason	Type of Studies to Consider
<ul style="list-style-type: none"> · Physical activity · Weight loss or maintenance · Adherence to devices over time 		<ul style="list-style-type: none"> before-and-after studies · Secondary analyses of existing trial data
<ul style="list-style-type: none"> · None of these studies examined factors that impact accelerometer adherence. Future research aimed at identifying individual characteristics associated with accelerometer adherence will inform efforts to create a hierarchy of who is more likely to adhere to these devices. · Determine how participants interact with their generated data as feedback. · Explore facilitators and barriers to adoption of wearable activity devices. 	Insufficient information	<ul style="list-style-type: none"> · RCTs · Prospective or retrospective cohort studies · Non-randomized controlled before-and-after studies · Qualitative studies · Secondary analyses of existing trial data
Setting		
<ul style="list-style-type: none"> · No studies that recruited through clinical settings 	Insufficient information	<ul style="list-style-type: none"> · RCTs · Prospective or retrospective cohort studies · Nonrandomized controlled before-and-after studies

Abbreviation: RCTs=randomized controlled trials

CONCLUSIONS

Overall this review found low to moderate strength of evidence that the use of wearable accelerometers produces small positive effects on physical activity and weight. These changes, however, may not result in clinically significant changes to health-related outcomes. The small number of studies with small sample sizes and the moderate to high heterogeneity limit the conclusions that may be drawn. What appears evident is that accelerometers may be necessary—but not sufficient—to change behavior. Unfortunately, the data available for this review did not allow us to determine the most effective means of using wearable accelerometers to facilitate behavior change. Nevertheless, we observed qualitatively that studies with more robust effects used adjunctive interventions (*eg*, tailored feedback and behavioral counseling) to foster motivation, had longer intervention durations, and compared accelerometers to weaker controls. Focusing on specific goals, using evidence-based strategies, personalizing designs, and integrating social support are additional, critical components for achieving increased physical activity or reduced weight. Stepped and adaptive trial designs, as well as those that incorporate patient preferences, may be particularly well-suited to address questions about the effectiveness of accelerometers within the context of complex behavioral interventions to increase physical activity and other related health outcomes.^{56,57}

The overall paucity of studies that used accelerometers and the generally low quality of the included trials may be due, in part, to the only recent availability of accelerometers with visible displays to allow for self-monitoring and the ability to interface with behavioral programs (*ie*, ability to collect and share data in real time). This review raises a number of important questions and illuminates key gaps in the evidence. This includes a need to understand whether accelerometers outperform traditional pedometers, how activity device features may enhance adherence and outcomes, and the effects on long-term outcomes. The relatively low response rates and attrition trends over time from the included studies in this report underscore the



importance of integrating these devices in tailored, patient-focused behavioral interventions that offer dynamic content, real-time and personalized behavioral counseling, and progressive goal-setting.

These findings are important for both clinicians and policymakers who may be considering programs to encourage more widespread use of these technologies as a public health strategy to improve health and well-being. Although we found small significant effects related to activity device use, the heterogeneity of the included studies precludes firm conclusions about how they can be deployed most effectively in clinical practice as a tool to facilitate increased physical activity. Furthermore, we did not find any studies that sought to integrate physical activity data from wearable accelerometers into the patient's medical records to facilitate ongoing primary care and chronic disease management.

REFERENCES

1. U.S. Department of Health and Human Services. *Physical Activity and Health: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion; 1996. www.cdc.gov/nccdphp/sgr/contents.htm. Accessed June 23, 2015.
2. Littman AJ, Forsberg CW, Koepsell TD. Physical activity in a national sample of veterans. *Med Sci Sports Exerc*. 2009;41(5):1006-1013.
3. Myers J. The health benefits and economics of physical activity. *Curr Sports Med Rep*. 2008;7(6):314-316.
4. Weiss JP, Froelicher VF, Myers JN, Heidenreich PA. Health-care costs and exercise capacity. *Chest*. 2004;126(2):608-613.
5. Troiano RP, Macera CA, Ballard-Barbash R. Be physically active each day. How can we know? *J Nutr*. 2001;131(2S-1):451S-460S.
6. Sallis JF, Saelens BE. Assessment of physical activity by self-report: status, limitations, and future directions. *Res Q Exerc Sport*. 2000;71(2 Suppl):S1-14.
7. Duncan GE, Sydemann SJ, Perri MG, Limacher MC, Martin AD. Can sedentary adults accurately recall the intensity of their physical activity? *Prev Med*. 2001;33(1):18-26.
8. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008;40(1):181-188.
9. Steeves JA, Murphy RA, Crainiceanu CM, Zipunnikov V, Van Domelen DR, Harris TB. Daily patterns of physical activity by type 2 diabetes definition: comparing diabetes, prediabetes, and participants with normal glucose levels in NHANES 2003-2006. *Prev Med Rep*. 2015;2:152-157.
10. Loprinzi PD, Walker JF, Lee H. Association between physical activity and inflammatory markers among U.S. adults with chronic obstructive pulmonary disease. *Am J Health Promot*. 2014;29(2):81-88.
11. Chastin SF, Mandrichenko O, Helbostadt JL, Skelton DA. Associations between objectively-measured sedentary behaviour and physical activity with bone mineral density in adults and older adults, the NHANES study. *Bone*. 2014;64:254-262.
12. Lee J, Song J, Hootman JM, et al. Obesity and other modifiable factors for physical inactivity measured by accelerometer in adults with knee osteoarthritis. *Arthritis Care Res (Hoboken)*. 2013;65(1):53-61.
13. Bassett DR, Jr., Dinesh J. Use of pedometers and accelerometers in clinical populations: validity and reliability issues. *Phys Ther Rev*. 2010;15(3):135-142.
14. Tudor-Locke C. Taking steps toward increasing physical activity: using pedometers to measure and motivate. *President's Council on Physical Fitness and Sports Research Digest*, Series 3, No. 17; June 2002. www.presidentschallenge.org/informed/digest/docs/200206digest.pdf. Accessed June 23, 2015.
15. Le Masurier GC, Tudor-Locke C. Comparison of pedometer and accelerometer accuracy under controlled conditions. *Med Sci Sports Exerc*. 2003;35(5):867-871.



16. Tyo BM, Fitzhugh EC, Bassett DR, Jr., John D, Feito Y, Thompson DL. Effects of body mass index and step rate on pedometer error in a free-living environment. *Med Sci Sports Exerc.* 2011;43(2):350-356.
17. Lyons EJ, Lewis ZH, Mayrsohn BG, Rowland JL. Behavior change techniques implemented in electronic lifestyle activity monitors: a systematic content analysis. *J Med Internet Res.* 2014;16(8):e192.
18. Gonzalez C, Herrero P, Cubero JM, et al. PREDIRCAM eHealth platform for individualized telemedical assistance for lifestyle modification in the treatment of obesity, diabetes, and cardiometabolic risk prevention: a pilot study (PREDIRCAM 1). *J Diabetes Sci Technol.* 2013;7(4):888-897.
19. Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions.* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. www.cochrane-handbook.org. Accessed July 22, 2014.
20. Baker PR, Francis DP, Soares J, Weightman AL, Foster C. Community wide interventions for increasing physical activity. *Cochrane Database Syst Rev.* 2015;1:Cd008366.
21. Mansi S, Milosavljevic S, Baxter GD, Tumilty S, Hendrick P. A systematic review of studies using pedometers as an intervention for musculoskeletal diseases. *BMC Musculoskelet Disord.* 2014;15:231.
22. Mays RJ, Rogers RK, Hiatt WR, Regensteiner JG. Community walking programs for treatment of peripheral artery disease. *J Vasc Surg.* 2013;58(6):1678-1687.
23. Vaes AW, Cheung A, Atakhorrami M, et al. Effect of 'activity monitor-based' counseling on physical activity and health-related outcomes in patients with chronic diseases: A systematic review and meta-analysis. *Ann Med.* 2013;45(5-6):397-412.
24. De Bourdeaudhuij I, Van Dyck D, De Meester F, et al. Built environment, physical activity, and obesity. *Int J Obes.* 2011;35(Suppl 1):S149.
25. Bort-Roig J, Gilson N, Puig-Ribera A, Contreras R, Trost S. Measuring and Influencing Physical Activity with Smartphone Technology: A Systematic Review. *Sports Med.* 2014;44(5):671-686.
26. Bellet RN, Adams L, Morris NR. The 6-minute walk test in outpatient cardiac rehabilitation: validity, reliability and responsiveness—a systematic review. *Physiotherapy.* 2012;98(4):277-287.
27. Higgins JPT, Altman DG, Sterne JAC. Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions.* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. www.cochrane-handbook.org. Accessed June 23, 2015.
28. Luley C, Blaik A, Gotz A, et al. Weight loss by telemonitoring of nutrition and physical activity in patients with metabolic syndrome for 1 year. *J Am Coll Nutr.* 2014;33(5):363-374.
29. Shuger SL, Barry VW, Sui X, et al. Electronic feedback in a diet- and physical activity-based lifestyle intervention for weight loss: a randomized controlled trial. *Int J Behav Nutr Phys Act.* 2011;8:41.
30. Polzien KM, Jakicic JM, Tate DF, Otto AD. The efficacy of a technology-based system in a short-term behavioral weight loss intervention. *Obesity (Silver Spring).* 2007;15(4):825-830.

31. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hillsdale, NJ: Lawrence Earlbaum Associates; 1988.
32. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*. 2010;36(3):1-48.
33. Agency for Healthcare Research and Quality. *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. Rockville, MD: Agency for Healthcare Research and Quality. www.effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productid=318. Accessed November 6, 2014.
34. Tabak M, Brusse-Keizer M, van der Valk P, Hermens H, Vollenbroek-Hutten M. A telehealth program for self-management of COPD exacerbations and promotion of an active lifestyle: a pilot randomized controlled trial. *Int J Chron Obstruct Pulmon Dis*. 2014;9:935-944.
35. Nicklas BJ, Gaukstern JE, Beavers KM, Newman JC, Leng X, Rejeski WJ. Self-monitoring of spontaneous physical activity and sedentary behavior to prevent weight regain in older adults. *Obesity (Silver Spring)*. 2014;22(6):1406-1412.
36. Thompson WG, Kuhle CL, Koeppe GA, McCrady-Spitzer SK, Levine JA. "Go4Life" exercise counseling, accelerometer feedback, and activity levels in older people. *Arch Gerontol Geriatr*. 2014;58(3):314-319.
37. Wijsman CA, Westendorp RG, Verhagen EA, et al. Effects of a web-based intervention on physical activity and metabolism in older adults: randomized controlled trial. *J Med Internet Res*. 2013;15(11):e233.
38. Thompson WG, Koeppe GA, Levine JA. Increasing physician activity with treadmill desks. *Work*. 2014;48(1):47-51.
39. Greene J, Sacks R, Piniewski B, Kil D, Hahn JS. The impact of an online social network with wireless monitoring devices on physical activity and weight loss. *J Prim Care Community Health*. 2013;4(3):189-194.
40. Shrestha M, Combest T, Fonda SJ, Alfonso A, Guerrero A. Effect of an accelerometer on body weight and fitness in overweight and obese active duty soldiers. *Mil Med*. 2013;178(1):82-87.
41. Reijonsaari K, Vehtari A, Kahilakoski OP, van Mechelen W, Aro T, Taimela S. The effectiveness of physical activity monitoring and distance counseling in an occupational setting - results from a randomized controlled trial (CoAct). *BMC Public Health*. 2012;12:344.
42. Koizumi D, Rogers NL, Rogers ME, Islam MM, Kusunoki M, Takeshima N. Efficacy of an accelerometer-guided physical activity intervention in community-dwelling older women. *J Phys Act Health*. 2009;6(4):467-474.
43. Sloomaker SM, Chinapaw MJ, Schuit AJ, Seidell JC, Van Mechelen W. Feasibility and effectiveness of online physical activity advice based on a personal activity monitor: randomized controlled trial. *J Med Internet Res*. 2009;11(3):e27.
44. Paschali AA, Goodrick GK, Kalantzi-Azizi A, Papadatou D, Balasubramanyam A. Accelerometer feedback to promote physical activity in adults with type 2 diabetes: a pilot study. *Percept Mot Skills*. 2005;100(1):61-68.
45. Institute of Medicine. *Cognitive Aging: Progress in Understanding and Opportunities for Action*. Washington, DC: National Academies Press; 2015.

46. Pahor M, Guralnik JM, Ambrosius WT, et al. Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA*. 2014;311(23):2387-2396.
47. Umpierre D, Ribeiro PA, Kramer CK, et al. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA*. 2011;305(17):1790-1799.
48. Dunlop DD, Song J, Semanik PA, et al. Relation of physical activity time to incident disability in community dwelling adults with or at risk of knee arthritis: prospective cohort study. *BMJ*. 2014;348:g2472.
49. Fong DY, Ho JW, Hui BP, et al. Physical activity for cancer survivors: meta-analysis of randomised controlled trials. *BMJ*. 2012;344:e70.
50. Thorogood A, Mottillo S, Shimony A, et al. Isolated aerobic exercise and weight loss: a systematic review and meta-analysis of randomized controlled trials. *Am J Med*. 2011;124(8):747-755.
51. Herring MP, Puetz TW, O'Connor PJ, Dishman RK. Effect of exercise training on depressive symptoms among patients with a chronic illness: a systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med*. 2012;172(2):101-111.
52. Burke LE, Wang J, Sevick MA. Self-monitoring in weight loss: a systematic review of the literature. *J Am Diet Assoc*. 2011;111(1):92-102.
53. Bravata DM, Smith-Spangler C, Sundaram V, et al. Using pedometers to increase physical activity and improve health: a systematic review. *JAMA*. 2007;298(19):2296-2304.
54. Nielsen Newswire. Are consumers really interested in wearing tech on their sleeves? March 20, 2014. www.nielsen.com/us/en/insights/news/2014/tech-styles-are-consumers-really-interested-in-wearing-tech-on-their-sleeves.html. Accessed July 2, 2015.
55. Robinson KA, Saldanha IJ, Mckoy NA. *Frameworks for Determining Research Gaps During Systematic Reviews*. Methods Future Research Needs Report No. 2. (Prepared by the Johns Hopkins University Evidence-based Practice Center under Contract No. HHSA 290-2007-10061-I.) AHRQ Publication No. 11-EHC043-EF. Rockville, MD: Agency for Healthcare Research and Quality. June 2011. www.effectivehealthcare.ahrq.gov/reports/final.cfm. Accessed July 2, 2015.
56. Craig P, Dieppe P, Macintyre S, et al. Developing and evaluating complex interventions: the new Medical Research Council guidance.[Reprint in *Int J Nurs Stud*. 2013 May;50(5):587-92; PMID: 23159157]. *BMJ*. 2008;337:a1655.
57. Moore GF, Audrey S, Barker M, et al. Process evaluation of complex interventions: Medical Research Council guidance. *BMJ*. 2015;350:h1258.

APPENDIX A. SEARCH STRATEGIES

Database: MEDLINE (via PubMed)

Search date: 01/06/15

Set #	Search Terms	Results
#1	"Accelerometry"[Mesh] OR "Magnetometry"[Mesh] OR "Motor Activity/instrumentation"[Mesh] OR fitness track*[tiab] OR activity track*[tiab] OR fitness monitor*[tiab] OR gps[tiab] OR "global positioning"[tiab] OR activity monitor*[tiab] OR motion sens*[tiab] OR accelerometer[tiab] OR accelerometers[tiab] OR accelerometry[tiab] OR gyroscope[tiab] OR gyroscopic[tiab] OR gyroscopes[tiab] OR actograph[tiab] OR actographic[tiab] OR actography[tiab] OR actographs[tiab] OR wearable system[tiab] OR wearable systems[tiab] OR wearable sensor[tiab] OR wearable sensors[tiab] OR ((step[tiab] OR steps[tiab]) AND (counting[tiab] OR counted[tiab] OR counter[tiab] OR counters[tiab] OR count[tiab])) OR actigraph[tiab] OR (basis[tiab] AND peak[tiab]) OR "bowflex boost"[tiab] OR "fit link"[tiab] OR (misfit[tiab] AND shine[tiab]) OR (polar[tiab] AND loop[tiab]) OR bodybugg[tiab] OR bodymedia[tiab] OR fitbit[tiab] OR fitbug[tiab] OR fuelband[tiab] OR garmin[tiab] OR gowear[tiab] OR gruve[tiab] OR ibitz[tiab] OR iqua[tiab] OR lumo[tiab] OR motoactiv[tiab] OR runtastic[tiab] OR scosche[tiab] OR smartband[tiab] OR striiv[tiab] OR tomtom[tiab] OR vivofit[tiab] OR vivosmart[tiab] OR wahoo[tiab] OR wakemate[tiab] OR withings[tiab]	52,751
#2	"Movement"[Mesh] OR "Exercise Movement Techniques"[Mesh] OR "Exercise Therapy"[Mesh] OR "Physical Fitness"[Mesh] OR "Physical Endurance"[Mesh] OR "Physical Exertion"[Mesh] OR fitness[tiab] OR activity[tiab] OR active[tiab] OR walk*[tiab] OR run*[tiab] OR step[tiab] OR steps[tiab] OR exercise[tiab] OR move*[tiab]	3,555,057
#3	(randomized controlled trial[pt] OR controlled clinical trial[pt] OR randomized[tiab] OR randomised[tiab] OR randomization[tiab] OR randomisation[tiab] OR randomly[tiab] OR trial[tiab] OR groups[tiab]) NOT (Editorial[ptyp] OR Letter[ptyp] OR Case Reports[ptyp] OR Comment[ptyp])	2,079,904
#4	#1 AND #2 AND #3	4858
#5	#4 NOT ("Child"[Mesh] NOT "Adult"[Mesh])	4355
#6	#5, English, 2000 - present	3506

Database: Embase

Search date: 01/06/15

Set #	Search Terms	Results
#1	'accelerometry'/exp OR 'magnetometry'/exp OR (fitness NEAR/2 track*):ab,ti OR (activity NEAR/2 track*):ab,ti OR (fitness NEAR/2 monitor*):ab,ti OR gps:ab,ti OR 'global positioning':ab,ti OR (activity NEAR/2 monitor):ab,ti OR (motion NEAR/2 sens*):ab,ti OR accelerometer:ab,ti OR accelerometers:ab,ti OR accelerometry:ab,ti OR gyroscope:ab,ti OR gyroscopic:ab,ti OR gyroscopes:ab,ti OR actograph:ab,ti OR actographic:ab,ti OR actography:ab,ti OR actographs:ab,ti OR 'wearable system':ab,ti OR 'wearable systems':ab,ti OR 'wearable sensor':ab,ti OR 'wearable sensors':ab,ti OR ((step OR steps):ab,ti AND (counting OR counted OR counter OR counters OR count):ab,ti) OR actigraph:ab,ti OR (basis NEAR/3 peak):ab,ti,df OR 'bowflex boost':ab,ti,df OR 'fit link':ab,ti,df OR (misfit NEAR/3 shine):ab,ti,df OR (polar NEAR/3 loop):ab,ti,df OR bodybugg:ab,ti,df OR bodymedia:ab,ti,df OR fitbit:ab,ti,df OR fitbug:ab,ti,df OR fuelband:ab,ti,df OR garmin:ab,ti,df OR gowear:ab,ti,df OR	45,316

Set #	Search Terms	Results
	gruve:ab,ti,df OR ibitz:ab,ti,df OR iqua:ab,ti,df OR lumo:ab,ti,df OR motoactiv:ab,ti,df OR runtastic:ab,ti,df OR scosche:ab,ti,df OR smartband:ab,ti,df OR striiv:ab,ti,df OR tomtom:ab,ti,df OR vivofit:ab,ti,df OR vivosmart:ab,ti,df OR wahoo:ab,ti,df OR wakemate:ab,ti,df OR withings:ab,ti,df	
#2	'movement (physiology)/exp OR 'physical activity, capacity and performance'/exp OR 'kinesiotherapy'/exp OR 'fitness'/exp OR fitness:ab,ti OR activity:ab,ti OR active:ab,ti OR walk*:ab,ti OR run*:ab,ti OR step:ab,ti OR steps:ab,ti OR exercise:ab,ti OR move*:ab,ti	4,564,954
#3	('randomized controlled trial'/exp OR 'crossover procedure'/exp OR 'double blind procedure'/exp OR 'single blind procedure'/exp OR random*:ab,ti OR factorial*:ab,ti OR crossover*:ab,ti OR (cross NEAR/1 over*):ab,ti OR (doubl* NEAR/1 blind*):ab,ti OR (singl* NEAR/1 blind*):ab,ti OR assign*:ab,ti OR allocat*:ab,ti OR volunteer*:ab,ti) NOT ('case report'/exp OR 'case study'/exp OR 'editorial'/exp OR 'letter'/exp OR 'note'/exp)	1,431,100
#4	#1 AND #2 AND #3	3250
#5	#4 NOT ('child'/exp NOT 'adult'/exp)	2888
#6	#5 AND [embase]/lim NOT [medline]/lim	1051
#7	#6, Limits: English, 2000-	988

Database: CINAHL

Search date: 01/06/15

Set #	Search Terms	Results
#1	(MH "Accelerometry") OR (MH "Magnetics+") OR TI ("fitness track*" or "activity track*" or "fitness monitor*" or gps or "global positioning" or "activity monitor*" or "motion sens*" or accelerometer or accelerometers or accelerometry or gyroscope or gyroscopic or gyroscopes or actograph or actographic or actography or actographs or "wearable system" or "wearable systems" or "wearable sensor" or "wearable sensors" or ((step or steps) and (counting or counted or counter or counters or count)) or actigraph or (basis and peak) or "bowflex boost" or "fit link" or (misfit and shine) or (polar and loop) or bodybugg or bodymedia or fitbit or fitbug or fuelband or garmin or gowear or gruve or ibitz or iqua or lumo or motoactiv or runtastic or scosche or smartband or striiv or tomtom or vivofit or vivosmart or wahoo or wakemate or withings) OR AB ("fitness track*" or "activity track*" or "fitness monitor*" or gps or "global positioning" or "activity monitor*" or "motion sens*" or accelerometer or accelerometers or accelerometry or gyroscope or gyroscopic or gyroscopes or actograph or actographic or actography or actographs or "wearable system" or "wearable systems" or "wearable sensor" or "wearable sensors" or ((step or steps) and (counting or counted or counter or counters or count)) or actigraph or (basis and peak) or "bowflex boost" or "fit link" or (misfit and shine) or (polar and loop) or bodybugg or bodymedia or fitbit or fitbug or fuelband or garmin or gowear or gruve or ibitz or iqua or lumo or motoactiv or runtastic or scosche or smartband or striiv or tomtom or vivofit or vivosmart or wahoo or wakemate or withings)	14,089
#2	(MH "Movement+") OR (MH "Exercise+") OR (MH "Therapeutic Exercise+") OR (MH "Physical Activity") OR (MH "Physical Fitness+") OR (MH "Exertion+") OR TI (OR fitness OR activity OR active OR walk* OR run* OR step OR steps OR exercise OR move*) OR AB (OR fitness OR activity OR active OR walk* OR run* OR step OR steps OR exercise OR move*)	361,653
#3	(MH "Treatment Outcomes+") OR randomized OR PT clinical trial	317,587
#4	#1 AND #2 AND #3	636
#5	#4, English, 2000-	602



Database: SPORTDiscus

Search date: 01/06/15

Set #	Search Terms	Results
#1	DE "ACCELEROMETERS" OR TI ("fitness track*" or "activity track*" or "fitness monitor*" or gps or "global positioning" or "activity monitor*" or "motion sens*" or accelerometer or accelerometers or accelerometry or gyroscope or gyroscopic or gyroscopes or actograph or actographic or actography or actographs or "wearable system" or "wearable systems" or "wearable sensor" or "wearable sensors" or ((step or steps) and (counting or counted or counter or counters or count)) or actigraph or (basis and peak) or "bowflex boost" or "fit link" or (misfit and shine) or (polar and loop) or bodybugg or bodymedia or fitbit or fitbug or fuelband or garmin or gowear or gruve or ibitz or iqua or lumo or motoactiv or runtastic or scosche or smartband or striiv or tomtom or vivofit or vivosmart or wahoo or wakemate or withings) OR AB ("fitness track*" or "activity track*" or "fitness monitor*" or gps or "global positioning" or "activity monitor*" or "motion sens*" or accelerometer or accelerometers or accelerometry or gyroscope or gyroscopic or gyroscopes or actograph or actographic or actography or actographs or "wearable system" or "wearable systems" or "wearable sensor" or "wearable sensors" or ((step or steps) and (counting or counted or counter or counters or count)) or actigraph or (basis and peak) or "bowflex boost" or "fit link" or (misfit and shine) or (polar and loop) or bodybugg or bodymedia or fitbit or fitbug or fuelband or garmin or gowear or gruve or ibitz or iqua or lumo or motoactiv or runtastic or scosche or smartband or striiv or tomtom or vivofit or vivosmart or wahoo or wakemate or withings)	6204
#2	(random* OR trial)	56299
#3	#1 AND #2	639
#4	#3, English, 2000-, Academic Journals	543

Database: Cochrane CENTRAL

Search date: 01/06/15

Set #	Search Terms	Results
#1	[mh Accelerometry] OR [mh Magnetometry]	341
#2	"fitness track*":ab,ti or "activity track*":ab,ti or "fitness monitor*":ab,ti or gps:ab,ti or "global positioning":ab,ti or "activity monitor*":ab,ti or "motion sens*":ab,ti or accelerometer:ab,ti or accelerometers:ab,ti or accelerometry:ab,ti or gyroscope:ab,ti or gyroscopic:ab,ti or gyroscopes:ab,ti or actograph:ab,ti or actographic:ab,ti or actography:ab,ti or actographs:ab,ti or "wearable system":ab,ti or "wearable systems":ab,ti or "wearable sensor":ab,ti or "wearable sensors":ab,ti or ((step:ab,ti or steps:ab,ti) and (counting:ab,ti or counted:ab,ti or counter:ab,ti or counters:ab,ti or count:ab,ti)) or actigraph:ab,ti or (basis:ab,ti and peak:ab,ti) or "bowflex boost":ab,ti or "fit link":ab,ti or (misfit:ab,ti and shine:ab,ti) or (polar:ab,ti and loop:ab,ti) or bodybugg:ab,ti or bodymedia:ab,ti or fitbit:ab,ti or fitbug:ab,ti or fuelband:ab,ti or garmin:ab,ti or gowear:ab,ti or gruve:ab,ti or ibitz:ab,ti or iqua:ab,ti or lumo:ab,ti or motoactiv:ab,ti or runtastic:ab,ti or scosche:ab,ti or smartband:ab,ti or striiv:ab,ti or tomtom:ab,ti or vivofit:ab,ti or vivosmart:ab,ti or wahoo:ab,ti or wakemate:ab,ti or withings:ab,ti	2945
#3	#1 OR #2	3204
#4	[mh "Movement"] OR [mh "Exercise Movement Techniques"] OR [mh "Exercise Therapy"] OR [mh "Physical Fitness"] OR [mh "Physical Endurance"] OR [mh "Physical Exertion"]	29,268



Set #	Search Terms	Results
#5	fitness:ab,ti OR activity:ab,ti OR active:ab,ti OR walk*:ab,ti OR run*:ab,ti OR step:ab,ti OR steps:ab,ti OR exercise:ab,ti OR move*:ab,ti	134,034
#6	#4 OR #5	140,071
#7	#3 AND #6	1630
#6	#5, 2000 – present, In Trials	1281

APPENDIX B. PEER REVIEWER COMMENTS TABLE

Question Text	Reviewer Number	Comment	Response
Are the objectives, scope, and methods for this review clearly described?	3	Yes	N/A
	4	Yes	N/A
	5	Yes	N/A
	6	<p>No - My primary concern with this document is the broad conceptualization of the difference between an accelerometer and a pedometer.</p> <p>A pedometer is something that counts steps.</p> <p>An accelerometer is something that counts accelerations possibly with some additional metric of intensity.</p> <p>In the old days, pedometers counted steps using a spring and magnet internal mechanism. Now, most pedometers are actually using internal accelerometers to measure step counts. So, is a pedometer that uses an internal mechanism of an accelerometer a pedometer or an accelerometer. These are not technically mutually exclusive categories. Examples of accelerometer based pedometers include Stepwatch, Omron HJ-720 ITC and Fitbits. Many interventions using classic accelerometers such as actigraphs actually only feed back step count data even if they could feed back the full accelerometer data so would these count in this synthesis?</p> <p>It appears that you chose to exclude studies of interventions that use activity monitors with internal accelerometer mechanisms that report step counts back to the user rather than the full accelerometer data. The decision to exclude interventions that use accelerometer based activity trackers to give step count feedback over the internet seems arbitrary. Accelerometer based pedometers have all of the advantages of accelerometers except that they filter out non-step count accelerations before presenting the data (step count) to the wearer.</p> <p>If you want to do a synthesis of trials that feedback non-step count physical activity metrics obtained from an accelerometer, then the intro and justification needs to be rewritten to reflect this focus. As it is written, it does not</p>	<p>Thank you for these comments. This report and other products have been explicitly developed to meet the needs of our stakeholder partners. There have been several excellent reviews of the effects of pedometers on physical activity and other health outcomes (Bravata, 2007, JAMA; Richardson, 2008, Annals of Family Medicine). As such, we developed a “white paper” that provides a review of recent good- and fair-quality systematic reviews on the effect of pedometer use on weight change, physical activity, and specific conditions of interest (eg, osteoarthritis, COPD, heart disease) to summarize the evidence for our primary stakeholders.</p> <p>Stakeholders were also interested in wearable technologies that measured activity beyond step counts (non-pedometer–based wearable motion sensing technologies), as those technologies tend to be more costly than traditional pedometers that only capture or report step counts. We agree that distinguishing between pedometers and accelerometers can be challenging. Our inclusion criteria consisted of the use of a wearable activity device that provided objective feedback to the wearer and was not defined as a pedometer-only device by study authors. To determine this, we used the author’s definition of the device. If the underlying mechanics of a device were unclear, we searched online resources to determine each device type and manufacturer to ascertain if the predominate technology used something beyond, or in addition to, pedometers for step counts. We have added this information to Appendix C. All of the included wearable devices were considered to be accelerometers by manufacturer’s definition. Our exclusion criteria did seek to exclude pedometer-based studies only if they</p>

Question Text	Reviewer Number	Comment	Response
		<p>currently make sense. It is simply not true that a) pedometers are less accurate than accelerometers (because many pedometers these days actually are accelerometers), b) accelerometers can link to electronic networks and devices but pedometers cannot (since the early days of the sportbrain, even spring and magnet pedometers have been able to upload step count data to networks and devices, now fitbits etc are accelerometer based pedometers that link up) .</p> <p>I think you need to rewrite the criteria and clearly state that your question is - does feedback related to accelerometer count and intensity change physical activity , weight and other health outcomes. It is the data that is fed back to the user that varies - either step counts vs accelerations with intensity ratings.</p>	<p>identified themselves as a pedometer and feedback was not part of the intervention. We did not exclude studies where the intervention gave step count feedback over the internet. We did not restrict the type of feedback that was provided to the participant, and several articles included studies that measured physical activity on different scales and with different measures including step counts.</p>
	7	Yes	N/A
Is there any indication of bias in our synthesis of the evidence?	3	No	N/A
	4	No	N/A
	5	No	N/A
	6	No	N/A
	7	No	N/A
Are there any <u>published</u> or <u>unpublished</u> studies that we may have overlooked?	3	No	N/A
	4	No	N/A
	5	No	N/A
	6	<p>Yes - If you decide to include accelerometer based interventions that limit feedback to step count data then you have missed many studies. If the focus and exclusion criteria are really intended to exclude accelerometer based step count feedback interventions and if these criteria are made more coherent, then I think you may have all of the relevant studies.</p>	<p>Thank you for this comment. Our eligibility criteria did not exclude studies that limit feedback to step count data. We included any objective feedback that was provided to the participant (wearer) and that was not defined as a pedometer-only study by study authors.</p>
	7	No	N/A
Additional suggestions or comments	3	<p>Overall thorough and appears to be well-done. Key questions are addressed as well as can be for the amount of data available.</p>	<p>Thank you.</p>

Question Text	Reviewer Number	Comment	Response
<p>can be provided below. If applicable, please indicate the page and line numbers from the draft report.</p>		<p>1) In the abstract, you report the SMD for the 9 studies using an inactive comparator for physical activity, and in the conclusions, you choose to highlight the SMD for the overall. I recommend being consistent in which estimate you highlight to avoid confusion.</p> <p>2) In the abstract, you conclude that the devices produce small positive effects on physical activity and weight that are not likely to have a clinically significant impact on health outcomes. This statement strikes me as too definite of a conclusion. I think that your support for the small positive effects is well-substantiated in the methods, results, and conclusions. I think your support for the interpretation of the lack of clinically significant impact on health outcomes is lacking. If you wish to keep this statement in the conclusions of the abstract, you need to provide support in the Discussion for the fact that these small differences in physical activity and weight do not translate to differences in clinical outcomes (ok to make this judgment using data from other studies). If you don't provide adequate support for this statement, you may need to leave this more open-ended. Furthermore, the confidence intervals for effects are still relatively wide, and do not exclude a medium-sized effect.</p> <p>3) On Forrest plots, it would be easier/faster to interpret if the subgroup "Summary" line was labeled (e.g. Inactive Comparator Summary, Active Comparator Summary).</p> <p>4) It wasn't a pre-defined KQ2 item, but it is mentioned a couple of times - would it be worth it to do a formal analysis on differences by duration of intervention?</p>	<p>1) We agree about being consistent with the estimates we provide. We now provide inactive, active, and overall summary estimate for both outcomes.</p> <p>2) We agree that our conclusion was originally written too definitively. We have made changes to soften the conclusion and highlight the limitations in this body of literature. Specifically, we have indicated that the small positive effects found for these devices on physical activity and weight may not result in clinically important outcomes. We have also highlighted in the Abstract's conclusion that the small number of studies with small sample sizes and the moderate to high heterogeneity limit the conclusions that may be drawn.</p> <p>3) We have added labels to all the stratified summary estimates.</p> <p>4) We did abstract the information on duration of intervention from individual studies. We did not, however, include intervention duration in the <i>a priori</i> analysis plan developed in collaboration with our primary stakeholders. As such, we report on those analyses that are <i>a priori</i>. We agree that formal analysis on differences by duration of intervention would be interesting, and we will explore including</p>

Question Text	Reviewer Number	Comment	Response
		I think the section on Evidence Gaps and Future Research will end up being very helpful!	these in other analyses beyond this report. Thank you.
	4	<p>Overall, I commend the authors on an extremely well written synthesis of this literature. I was unable to locate any additional RCTs that would meet the search criteria. I believe this report to be unbiased and a fair assessment of or current knowledge of these technologies. I have made some minor comments below for clarification and consideration.</p> <p>(p5, Abstract) – The abstract briefly mentions KQ2 in the background (i.e. factors affecting use), but nowhere else. The authors may consider adding a line about the paucity of literature in this regard should text limit be available.</p> <p>(pg5, Abstract) – In my opinion, the concluding statements could be revised somewhat to acknowledge the limitations of these data and relative immaturity of this field.</p> <p>(pg9, Introduction) – As the targeted audience for this report is VA clinicians, managers and policymakers – the authors may consider adding in some stats that are specific to Veterans and physical activity. For example, I believe there is some evidence that VA users in comparison to non-users are less likely to meet PA recommendations (Littman et al. 2009, MSSE). In addition, Consider including some relevant studies from VHA describing health-care costs and exercise capacity in Veterans (Weiss et al. 2004 Chest; Myers 2008 Curr Sports Med Report, etc.) including data derived from the Veterans Exercise Testing Study from Jon Myers’ group in Palo Alto.</p> <p>(p61, Appendix B) – I’m not sure a column for risk of bias</p>	<p>Thank you.</p> <p>We agree. We have added a sentence to the Results section that we did not identify any moderating factors that significantly impacted the effectiveness of the outcomes. This was our KQ 2 objective.</p> <p>We agree. We have made changes to soften the conclusion and highlight the limitations in this body of literature.</p> <p>Thank you for the suggestion to more clearly highlight the relevance of this inactivity problem for our stakeholders and intended audience. We have expanded the Introduction to include specific statistics on Veterans and physical (in)activity as well as the healthcare cost implications of sedentary behavior/lifestyle, citing the compelling literature you provided.</p> <p>Thank you for this comment. We decided to add the risk of</p>

Question Text	Reviewer Number	Comment	Response
		is necessary here as this was thoroughly covered in a prior section.	bias to the Appendix C table (formerly Appendix B table) since it was a standalone table and we felt that many readers may want to view the risk of bias assessment alongside the individual studies and study characteristics.
	5	<p>This report is a very elegant piece of work that clearly represents a tremendous amount of work undertaken to synthesize all of the available data on this topic. My concern is that the major take-home message of this report, from my perspective, may be a bit misleading which may lead to erroneous assumptions about the outcomes. "We found that use of the wearable devices produces small positive effects that are not likely to have a clinically significant impact on health outcomes. Clinicians and policymakers should consider these findings before widespread use of this technology etc". I believe some of the evidence gaps should be noted in the concluding statement. Policy makers might be inclined to conclude that there is no point in investing in this technology when in fact your data suggest that there is a paucity of well designed studies and that more research is needed that could better inform this decision. For example, on page 14 regarding KQa, the final sentence of the summary states that the "small number of studies, small sample sizes within these studies, and the moderate to high heterogeneity limits conclusions that may be drawn." This statement sounds like a more overall accurate conclusion than what is currently stated on the abstract main conclusion which is what most people will be most likely to read. I also fear that it is a big jump to make a conclusion about clinical impact on health outcomes when the focus of this report is on behaviors (activity and weight management).</p> <p>On page 28 - Summary of findings. The last sentence is not clear. It has a double negative which I am not sure makes sense 'The same trend was not seen for decreases in PA etc.' It seems that it should be 'The same trend was not seen for greater increases in PA' etc.</p>	<p>Thank you for your thorough and thoughtful review. We agree about the conclusions in the previous draft report. We have made changes to soften the conclusions and highlight the limitations in this body of literature. Specifically, we have indicated that the small positive effects found for these devices on physical activity and weight may not result in clinically important outcomes. We have also highlighted in the Abstract's conclusion that the small number of studies with small sample sizes and the moderate to high heterogeneity limit the conclusions that may be drawn.</p> <p>Thank you for identifying this error. We have changed to "greater increases in PA" as suggested.</p>

Question Text	Reviewer Number	Comment	Response
		<p>On page 39 in the bottom paragraph of Clinical and Policy implications. This paragraph seems to be "promoting" use of devices and has some cautionary verbiage about the limitations of the evidence which seems to be contradictory to the conclusion in the structured abstract. It would seem to me that you can't have it both ways. These seem like mixed messages. I believe the data support this second conclusion more than the one in the structured abstract.</p> <p>The gaps and future research table is very useful. Hopefully this will guide future research. It was interesting to note the final row in which it was stated that no studies were recruited through clinical settings which should have a huge bearing on clinical policy makers in that we truly have no evidence in favor or against the use of wearable monitors in clinical settings.</p>	<p>We agree. We have changed the structured Abstract as mentioned above to highlight the limitations in the body of literature.</p> <p>Thank you.</p>
	6	<p>2) There are several intervention characteristics that we know have an impact on behavior change. Goal-setting is one that may be the most critical given the focus on self-monitoring. Others include strategies to overcome barriers, motivational interviewing etc. It would be helpful to understand if / how participants set or were prescribed activity count goals for each of the included studies. Did all the trials include some kind of coaching around goal setting. If not, does explicit attention to goal setting impact outcomes</p> <p>3) The second question related to heterogeneity of treatment effect is so profoundly limited by small sample size that it probably should be reported as unable to judge due to risk of type 2 error rather than as not difference detected.</p> <p>4) Further limitations should be emphasized – there is insufficient evidence to determine if accelerometer based feedback is inferior, equivalent or superior to pedometer</p>	<p>2) Thank you for posing this question. Due to the high heterogeneity in adjunctive interventions delivered alongside the activity devices, we were unable to explore these additions quantitatively. Table 3 outlines the types of adjunctive interventions used for each study. Thus, we chose to categorize the use of the activity device alongside the adjunct intervention as “major vs minor.” It would be interesting for future analyses to examine the impact on outcomes that goal-setting may have with these devices.</p> <p>3) We agree. We have highlighted in the report the small sample sizes found in individual trials. We added to the Limitations section that the small sample sizes found within this literature may have resulted in a type II error.</p> <p>4) Thank you for this suggestion. We have added this information to the Research Gaps/Future Research section of the report.</p>

Question Text	Reviewer Number	Comment	Response
	7	<p>based feedback for any of the outcomes.</p> <p>These comments are submitted on behalf of the National Center for Health Promotion and Disease Prevention (NCP). The report was thorough and organized and the information was presented in a clear manner. There were two points of potential clarification that we wanted to bring to the attention of the authors: 1. Pg. 1 (lines 22-24)- physical activity diaries are included as a self-reported measure that "does little to motivate or change physical activity behavior..." While diaries and other real time records of self-monitored behavior are self-reported, they are not intended to be the same as questionnaires and interviews that gather self-reported data retrospectively. In addition, the act of attending to a target behavior using self-monitoring strategies has been shown to result in behavior change. Unless there is evidence to suggest that self-monitoring of behavior IS NOT associated with behavior change, or self-monitoring of physical activity is NOT associated with changes in physical activity, it may be helpful to reword this sentence.</p> <p>2. Pg. 16 (lines 33-54), Figure 4 - What does the absence of color/white colored bar represent? Also, it appears that the overall percentage of high risk of bias (as indicated by the proportion of the bar colored red in the "Overall" row) is much larger than one would expect given the relatively low risk of bias (as depicted by minimal red coloring in each of the individual risk of bias items). Additional description of the information in Figure 4 may provide clarity on these issues.</p>	<p>Thank you for the positive comments about the thoroughness and organization of the report.</p> <p>1) We agree with the review team that the way the sentence on page1, lines 22-24, is written is misleading. We initially intended to indicate that these self-report measures, such as diaries, may not be an optimal measurement strategy rather than imply them as intervention strategies. We have since revised the report to read, "Physical activity has previously been measured using self-reported measures (eg, questionnaires, interviews.); however, these do little to motivate or change physical activity behavior and can be subject to reporting bias."</p> <p>2) Thank you for noticing this error. We have revised the Risk of Bias figure and removed the "clear" space from the PA and weight questions. This occurred since overall we had 14 studies included; however, some studies did not include both outcomes of interest. There is high risk of bias in many of the studies even without individual questions being rated as high risk because of the way the overall evaluation is calculated. In the Methods section, we describe the scoring process for "high risk" which includes two or more domain questions that are "not clear." As such, many of the studies had "not clear" domain questions, leading to an overall high risk of bias. We have added the scoring as a footnote to Figure 4.</p>

APPENDIX C. CHARACTERISTICS OF INCLUDED STUDIES

Study	Population N Randomized Device Name	Intervention	Comparator	Outcomes	Duration (Weeks)	Funding Source	Risk of Bias
Greene, 2013 ¹	Healthy volunteers 513 NR	6 months access to online social network to post about weight and PA progress + continuous accelerometer use and feedback	Printed lifestyle guidelines on diet (sample daily meal plan with recommended serving sizes), recommendations on daily level of exercise, and articles on benefits of exercise and healthy eating	PA: Yes Weight: Yes Satisfaction: No	12 to 24	Industry	High
Koizumi, 2009 ²	Older adults 68 Kenz Lifecorder accelerometer	12 weeks of accelerometer with feedback + goal-setting	12-week blinded accelerometer with instruction to continue normal daily activity	PA: Yes Weight: No Satisfaction: No	12 to 24	NR	High
Luley, 2014 ³	Chronic medical illness 184 Aipermotion 440 accelerometer	3-arm study (2 interventional): <u>4s intervention:</u> 1-time, 2-hour instruction on diet & PA + 12 months of accelerometer use + 52 weekly individual letters with feedback on weight, diet, and PA <u>ABC intervention:</u> 1-time, 2-hour instruction on diet & PA + 12 months of accelerometer use + 12 monthly behavioral counseling calls	1-time, 2-hour session, consisting of diet education, diet regimen, and PA education	PA: No Weight: Yes Satisfaction: No	25+	Gov	Unclear

Study	Population N Randomized Device Name	Intervention	Comparator	Outcomes	Duration (Weeks)	Funding Source	Risk of Bias
Nicklas, 2014 ⁴	Older adults 48 Lifecorder PlusVR tri-axial accelerometer	5-month weight loss intervention that included hypocaloric diet (2 prepared meals a day) + 4 days/week supervised exercise + self-regulatory intervention that involved wearing an accelerometer, documenting activity, and 6 weekly sessions of behavioral counseling	5-month weight loss intervention consisting of diet education and regimen (individualized hypocaloric intake), PA education, structured exercise (supervised treadmill) and in-person counseling	PA: Yes Weight: Yes Satisfaction: No	25+	Gov	High
Paschali, 2005 ⁵	Chronic medical illness 30 BioTrainer accelerometer	12 weeks of continuous accelerometer use and feedback + 4 monthly in-person exercise behavioral counseling sessions + workbook	12-week blinded accelerometer with 4 monthly in-person counseling sessions to review exercise diary. A 24-page information book and home based walking plan with PA education, behavioral self-management, weight goal-setting, and chronic disease-monitoring.	PA: Yes Weight: No Satisfaction: No	12 to 24	Gov	High

Study	Population N Randomized Device Name	Intervention	Comparator	Outcomes	Duration (Weeks)	Funding Source	Risk of Bias
Polzien, 2007 ⁶	Overweight/ obese and/or sedentary 57 SenseWear Pro Armband	3-arm study: <u>Standard in-person behavioral weight control program:</u> 7 in-person individualized counseling session over 12 weeks <u>Continuous technology-based behavioral weight control program:</u> 7 in-person individualized counseling session over 12 weeks + 12 weeks of continuous accelerometer use and feedback <u>Intermittent technology-based behavioral weight control program:</u> 7 in-person individualized counseling session over 12 weeks + 3 weeks of accelerometer use and feedback over 12 weeks	7 in-person individualized counseling sessions consisting of diet education, diet regimen (1200 to 1500 kcal/day; dietary fat <20% of total energy intake), PA education, and weight goal-setting	PA: Yes Weight: Yes Satisfaction: No	12 to 24	Industry	High
Reijonsaari, 2012 ⁷	Healthy volunteers 544 Uni-axial accelerometer	12 months of continuous accelerometer use and feedback + access to telephone counseling (frequency NR)	Written and verbal explanation of results of physical exams, general PA, and health information and occupational healthcare	PA: Yes Weight: Yes Satisfaction: No	25+	Fdn/Indiv	Low

Study	Population N Randomized Device Name	Intervention	Comparator	Outcomes	Duration (Weeks)	Funding Source	Risk of Bias
Shrestha, 2013 ⁸	Overweight/ obese and/or sedentary 28 Polar FA20 accelerometer	1-time, 1.5 hour lifestyle instruction + 6 months of continuous accelerometer use and feedback	Self-directed exercise and/or US Army mandated physical training	PA: Yes Weight: Yes Satisfaction: No	12 to 24	Gov	High
Shuger, 2011 ⁹	Overweight/ obese and/or sedentary 197 BodyMedia SenseWear Armband	4-arm study (3 interventional arms) : <u>Group-based behavioral weight loss:</u> 14 group weight loss sessions + 6 individual phone calls + workbook over 9 months <u>Accelerometer alone:</u> 9 months of continuous accelerometer use and feedback <u>Group-based behavioral weight loss + accelerometer use:</u> 9 months of continuous accelerometer use and feedback + 14 group weight loss sessions + 6 individual phone calls + workbook	Self-directed weight loss manual with diet education, PA education, and weight goal-setting	PA: No Weight: Yes Satisfaction: No	25+	Industry	Unclear

Study	Population N Randomized Device Name	Intervention	Comparator	Outcomes	Duration (Weeks)	Funding Source	Risk of Bias
Slootmaker, 2009 ¹⁰	Overweight/ obese and/or sedentary 102 PAM (model AM101), uniaxial activity monitor + smartphone	3 months of continuous accelerometer use + tailored PA feedback and motivational tips via web-based portal	A single written brochure with brief PA recommendations	PA: Yes Weight: Yes Satisfaction: No	12 to 24	Gov	High
Tabak, 2014 ¹¹	Chronic medical illness 29 Activity Coach (accelerometer-based activity sensor + smartphone)	Self-directed technology supported care program that included a tailored web-based exercise program, accelerometer-based activity sensor and motivational messaging, COPD self-management module, and as needed web-portal teleconsultation conducted over 9 months	Usual care with regular physiotherapy, if prescribed	PA: Yes Weight: No Satisfaction: No	25+	Gov	Unclear
Thompson, 2014 ¹²	Overweight/ obese and/or sedentary 20 GRUVE triaxial accelerometer	12 weeks of continuous accelerometer use and feedback + weekly brief counseling sessions on increasing activity + treadmill desk	12 weeks of monitoring without feedback, then crossed over to receive monitoring, plus feedback, counting, and treadmill desk for 12 weeks	PA: Yes Weight: Yes Satisfaction: No	12 to 24	Fdn	High

Study	Population N Randomized Device Name	Intervention	Comparator	Outcomes	Duration (Weeks)	Funding Source	Risk of Bias
Thompson, 2014 ¹³	Older adults 49 FitBit	24 weeks of continuous accelerometer use and feedback + weekly brief telephone counseling sessions focused on accelerometer feedback + 6 in-person brief counseling sessions	24 weeks of accelerometer without feedback	PA: Yes Weight: Yes Satisfaction: No	25+	Gov/Fdn	Unclear
Wijsman, 2013 ¹⁴	Older adults 235 DirectLife	12 weeks of continuous accelerometer use and feedback + personal website + personal e-coach who gives updates on activity status and advice via web portal	3-month waitlist control	PA: Yes Weight: Yes Satisfaction: No	12 to 24	Gov/ Industry	Low

Abbreviations: COPD=chronic obstructive pulmonary disease; Fdn=foundation; Gov=government; Indiv=individual; N=number; NR=not reported; PA=physical activity

References in Appendix C

1. Greene J, Sacks R, Piniewski B, Kil D, Hahn JS. The impact of an online social network with wireless monitoring devices on physical activity and weight loss. *J Prim Care Community Health*. 2013;4(3):189-194.
2. Koizumi D, Rogers NL, Rogers ME, Islam MM, Kusunoki M, Takeshima N. Efficacy of an accelerometer-guided physical activity intervention in community-dwelling older women. *J Phys Act Health*. 2009;6(4):467-474.
3. Luley C, Blaik A, Gotz A, et al. Weight loss by telemonitoring of nutrition and physical activity in patients with metabolic syndrome for 1 year. *J Am Coll Nutr*. 2014;33(5):363-374.
4. Nicklas BJ, Gaukstern JE, Beavers KM, Newman JC, Leng X, Rejeski WJ. Self-monitoring of spontaneous physical activity and sedentary behavior to prevent weight regain in older adults. *Obesity (Silver Spring)*. 2014;22(6):1406-1412.
5. Paschali AA, Goodrick GK, Kalantzi-Azizi A, Papadatou D, Balasubramanyam A. Accelerometer feedback to promote physical activity in adults with type 2 diabetes: a pilot study. *Percept Mot Skills*. 2005;100(1):61-68.
6. Polzien KM, Jakicic JM, Tate DF, Otto AD. The efficacy of a technology-based system in a short-term behavioral weight loss intervention. *Obesity (Silver Spring)*. 2007;15(4):825-830.
7. Reijonsaari K, Vehtari A, Kahilakoski OP, van Mechelen W, Aro T, Taimela S. The effectiveness of physical activity monitoring and distance counseling in an occupational setting - results from a randomized controlled trial (CoAct). *BMC Public Health*. 2012;12:344.
8. Shrestha M, Combest T, Fonda SJ, Alfonso A, Guerrero A. Effect of an accelerometer on body weight and fitness in overweight and obese active duty soldiers. *Mil Med*. 2013;178(1):82-87.
9. Shuger SL, Barry VW, Sui X, et al. Electronic feedback in a diet- and physical activity-based lifestyle intervention for weight loss: a randomized controlled trial. *Int J Behav Nutr Phys Act*. 2011;8:41.
10. Slootmaker SM, Chinapaw MJ, Schuit AJ, Seidell JC, Van Mechelen W. Feasibility and effectiveness of online physical activity advice based on a personal activity monitor: randomized controlled trial. *J Med Internet Res*. 2009;11(3):e27.
11. Tabak M, Brusse-Keizer M, van der Valk P, Hermens H, Vollenbroek-Hutten M. A telehealth program for self-management of COPD exacerbations and promotion of an active lifestyle: a pilot randomized controlled trial. *Int J Chron Obstruct Pulmon Dis*. 2014;9:935-944.
12. Thompson WG, Koeppe GA, Levine JA. Increasing physician activity with treadmill desks. *Work*. 2014;48(1):47-51.
13. Thompson WG, Kuhle CL, Koeppe GA, McCrady-Spitzer SK, Levine JA. "Go4Life" exercise counseling, accelerometer feedback, and activity levels in older people. *Arch Gerontol Geriatr*. 2014;58(3):314-319.
14. Wijsman CA, Westendorp RG, Verhagen EA, et al. Effects of a web-based intervention on physical activity and metabolism in older adults: randomized controlled trial. *J Med Internet Res*. 2013;15(11):e233.

