

## APPENDIX A. SEARCH STRATEGIES

### MEDLINE/EMBASE

- 1 (coronavir\* or corona virus\* or betacoronavir\* or covid19 or covid 19 or nCoV or CoV 2 or CoV2 or sarscov2 **SARS 2** or **SARS-CoV-2** or 2019nCoV or 2019 novel coronavirus\* or 2019 novel CoV or wuhan virus\* or ((wuhan or hubei or huanan) and (severe acute respiratory or pneumonia\*))).ti,ab,kw.
- 2 Coronavirus Infections/ or Coronavirus/ or betacoronavirus/
- 3 1 or 2
- 4 Pulmonary fibrosis.ti,ab,kw. or exp Pulmonary Fibrosis/
- 5 exp Lung Diseases, Obstructive/
- 6 4 or 5
- 7 acute kidney injury.ti,ab,kw. or exp Acute Kidney Injury/
- 8 exp Renal Insufficiency, Chronic/
- 9 (end stage renal disease or ESRD or AKI or CKD).ti,ab,kw.
- 10 7 or 8 or 9
- 11 myocardial infarction.ti,ab,kw. or exp Myocardial Infarction/
- 12 (heart attack or heart failure or MI).ti,ab,kw.
- 13 myocarditis.ti,ab,kw. or exp Myocarditis/
- 14 exp Arrhythmias, Cardiac/
- 15 arrhythmia\*.ti,ab,kw.
- 16 11 or 12 or 14 or 14 or 15
- 17 exp Venous Thrombosis/
- 18 exp Pulmonary Embolism/ or exp Venous Thromboembolism/
- 19 (deep ve\* thrombosis or DVT or pulmonary embolism or PE).ti,ab,kw.
- 20 anemia.ti,ab,kw. or exp Anemia/
- 21 17 or 18 or 19 or 20
- 22 stroke.ti,ab,kw. or exp Stroke/
- 23 exp Cognitive Dysfunction/
- 24 exp Confusion/
- 25 exp Seizures/
- 26 exp Headache/
- 27 (stroke\* or cerebrovascular accident\* or cognitive impairment or cognitive dysfunction or delirium or confusion or seizure\* or headache\*).ti,ab,kw.
- 28 22 or 23 or 24 or 25 or 26 or 27
- 29 exp Diabetes Mellitus/
- 30 diabetes.ti,ab,kw.
- 31 29 or 30
- 32 exp Hepatitis/
- 33 exp Colitis/
- 34 (hepatitis or hepatocellular injur\* or colitis).ti,ab,kw.
- 35 32 or 33 or 34
- 36 "Autoimmune Diseases of the Nervous System"/
- 37 autoimmune disease\*.ti,ab,kw.
- 38 Musculoskeletal Diseases/
- 39 musculoskeletal.ti,ab,kw.
- 40 36 or 37 or 38 or 39
- 41 6 or 10 or 16 or 21 or 28 or 31 or 35 or 40
- 42 exp Hospitalization/ or exp Intensive Care Units/ or Inpatients/ or Subacute Care/
- 43 (hospital or hospitalized or hospitalization or intensive or ICU or care or post?acute or inpatient or inpatients or admit or admitted or admitting).ti,ab,kw.
- 44 42 or 43
- 45 3 and 41 and 44
- 46 limit 45 to english language
- 47 limit 46 to yr="2019 -Current"

## COCHRANE LIBRARY

- 1 MeSH descriptor: [Coronavirus] explode all trees
- 2 (coronavirus):ti,ab,kw
- 3 (betacoronavirus):ti,ab,kw
- 4 (covid19):ti,ab,kw
- 5 (covid 19):ti,ab,kw
- 6 (nCoV):ti,ab,kw
- 7 (CoV2):ti,ab,kw
- 8 (CoV2):ti,ab,kw
- 9 (OR #1-#8)
- 10 ("pulmonary fibrosis"):ti,ab,kw
- 11 MeSH descriptor: [Pulmonary Fibrosis] this term only
- 12 MeSH descriptor: [Lung Diseases, Obstructive] explode all trees
- 13 (acute kidney injury):ti,ab,kw
- 14 MeSH descriptor: [Acute Kidney Injury] this term only
- 15 MeSH descriptor: [Renal Insufficiency, Chronic] this term only
- 16 ("end stage renal disease"):ti,ab,kw
- 17 (ESRD):ti,ab,kw
- 18 (AKI):ti,ab,kw
- 19 (CKD):ti,ab,kw
- 20 ("myocardial infarction"):ti,ab,kw
- 21 MeSH descriptor: [Myocardial Infarction] this term only
- 22 ("heart attack"):ti,ab,kw
- 23 ("heart failure"):ti,ab,kw
- 24 (myocarditis):ti,ab,kw
- 25 MeSH descriptor: [Myocarditis] this term only
- 26 (arrhythmia\*):ti,ab,kw
- 27 MeSH descriptor: [Arrhythmias, Cardiac] this term only
- 28 MeSH descriptor: [Venous Thrombosis] this term only
- 29 MeSH descriptor: [Pulmonary Embolism] this term only
- 30 MeSH descriptor: [Venous Thromboembolism] this term only
- 31 ("deep venous thrombosis"):ti,ab,kw
- 32 ("pulmonary embolism"):ti,ab,kw
- 33 (anemia):ti,ab,kw
- 34 MeSH descriptor: [Anemia] this term only
- 35 MeSH descriptor: [Stroke] this term only
- 36 MeSH descriptor: [Cognitive Dysfunction] this term only
- 37 MeSH descriptor: [Confusion] this term only
- 38 MeSH descriptor: [Seizures] this term only
- 39 MeSH descriptor: [Headache] this term only
- 40 (stroke\*):ti,ab,kw
- 41 ("cerebrovascular accident"):ti,ab,kw
- 42 ("cognitive impairment"):ti,ab,kw
- 43 ("Cognitive dysfunction"):ti,ab,kw
- 44 (delirium):ti,ab,kw
- 45 (confusion):ti,ab,kw
- 46 (seizure\*):ti,ab,kw
- 47 (Headache\*):ti,ab,kw
- 48 (diabetes):ti,ab,kw
- 49 MeSH descriptor: [Diabetes Mellitus] this term only
- 50 MeSH descriptor: [Hepatitis] this term only
- 51 MeSH descriptor: [Colitis] this term only
- 52 (hepatitis):ti,ab,kw
- 53 ("hepatocellular injur\*"):ti,ab,kw

- 54 (colitis):ti,ab,kw
- 55 MeSH descriptor: [Autoimmune Diseases of the Nervous System] this term only
- 56 ("autoimmune disease"):ti,ab,kw
- 57 MeSH descriptor: [Musculoskeletal Diseases] this term only
- 58 (musculoskeletal):ti,ab,kw
- 59 (OR #10-#58)
- 60 (hospitalized):ti,ab,kw
- 61 (hospital):ti,ab,kw
- 62 (hospitalization):ti,ab,kw
- 63 ("intensive care"):ti,ab,kw
- 64 (ICU):ti,ab,kw
- 65 (Post-acute):ti,ab,kw
- 66 (Post acute):ti,ab,kw
- 67 (inpatient\*):ti,ab,kw
- 68 (admit\*):ti,ab,kw
- 69 MeSH descriptor: [Hospitalization] explode all trees
- 70 MeSH descriptor: [Intensive Care Units] this term only
- 71 MeSH descriptor: [Inpatients] this term only
- 72 MeSH descriptor: [Subacute Care] this term only
- 73 (OR #60-#72)
- 74 #9 AND #59 AND #73

## APPENDIX B. PEER REVIEWER COMMENTS AND RESPONSES

Question Text	Comment	Author Responses
Are the objectives, scope, and methods for this review clearly described?	Yes	Thank you
	Yes	
Is there any indication of bias in our synthesis of the evidence?	No	Thank you
	No	
Are there any published or unpublished studies that we may have overlooked?	No	Thank you.
	No	
Additional suggestions or comments can be provided below. If applicable, please indicate the page and line numbers from the draft report.	Well done -- an exhaustive summary of major organ system 'damage' after hospitalization for / with COVID-19. A limitation of available evidence noted near the top of page 46 is that studies did not reliably state what pre-admission / pre-COVID levels of function / dysfunction were present; what organ dysfunction occurred during hospitalization / was present at discharge, and then persisted for what periods of time thereafter. This data may be more useful than comparison groups -- healthcare systems will be interested in what proportion of COVID patients with "x" degree of pulm disability at discharge will have "y" needs that they did not have prior to infection. Depending on what your group thinks about this, you might make a clearer recommendation for future research to include this (say, over comparison groups if authors are thinking about study design and relative importance of different points of comparison).	Thank you.  We agree and have modified the statement on Timing in the future research section to include identification of pre-COVID, discharge, and persistent condition.
	Page 2, Figure 1. Analytic Framework: does not consider vaccination status as an independent variable, a missed opportunity Page 5, line 22: 'September 2021' needs appropriate spacing Page 6, line 32: arrow should point to the 'Hand search k=17' box, not away from it	Pg 2. We added vaccine status to the framework and in other places where modifying factors are discussed. Pg 5. We corrected the spacing. Pg 6. The arrow direction indicates that studies identified by hand-search underwent full-text review.



Question Text	Comment	Author Responses
	<p>In the updated review “COVID19 Post-acute Care Makor Organ Damage: a Living Rapid Review, the authors report existing evidence for the prevalence of post-acute care major organ damage and healthcare or service use needs for patients hospitalized for/with COVID-19.</p> <p>The revision is excellent and appears to have include all of the relevant newer studies that I am aware of (and several that I wasn’t). The authors appropriately point out the limitations in the evidence due to convenience sampling, often low (albeit increasing) sample sizes, variation in measurement definitions, and resulting wide ranges of prevalence reported. I really like the visualization of the studies within figures, as it captures this variation nicely and aids in the reader’s overall impression of observed complications in the different studied populations.</p> <p>I have only 1 minor comment. As this is a scoping review, wherever possible, it would be useful to use the same format to ease the reader. The bubble plots within the pulmonary section are especially helpful as they demonstrate the variation in both prevalence and days to follow-up. Could other figures (ie, cognitive complications, healthcare resource utilization) take on a similar format?</p>	<p>Thank you.</p> <p>We appreciate the suggestion but chose to leave the figures as they appeared in the peer review version.</p>
	<p>Thank you for this comprehensive review!</p> <p>Consider editing as "impaired pulmonary function" for Table 2.</p> <p>Throughout, it would be useful if "study design" could be added to the main tables (e.g., Table 3), just as it is in the appendix tables. This will assist in interpretation.</p> <p>Typo on page 23, "Soc studies..." (line 20)</p> <p>Is this correct? "In a large database study from the US, dementia was newly diagnosed in 0.23% of the COVID19 group and 0.43% of the non-COVID control group (risk difference 0.2% [95%CI 0.7, 0.3], P&lt;001).110"</p> <p>In the Key Findings, I would add a comment on the findings on respiratory endpoints in studies with control groups. This is done systematically for subsequent endpoints.</p> <p>Regarding C4R (c4r-nih.org): Currently, over 45,000 participants have completed COVID-19 questionnaires out of a target population of</p>	<p>Edited as suggested. Thank you..</p> <p>We added a notation on the tables for the studies with a comparator group.</p> <p>Corrected. Thank you.</p> <p>Thank you – this was an error (corrected to read 0.03% of control group).</p> <p>The key respiratory findings section has been edited.</p> <p>Thank you. We now cite the design paper and look forward to the findings!</p>



Question Text	Comment	Author Responses
	approximately 50,000. Major distinguishing features of C4R, which directly address several issues raised in the report, include the availability of research-quality pre-COVID measures of subclinical and subclinical disease, as well as behavioral factors, biomarkers, and multi-'Omics, in hospitalized and non-hospitalized COVID-19 patients and also non-infected comparators. The design paper is pre-printed here: <a href="https://www.medrxiv.org/content/10.1101/2021.03.19.21253986v1.full.pdf">https://www.medrxiv.org/content/10.1101/2021.03.19.21253986v1.full.pdf</a>	

## APPENDIX C. EVIDENCE TABLES

**TABLE 1. STUDY CHARACTERISTICS**

(Shaded rows indicate a newly included study since previous update)

Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Akhtar, 2021 <sup>39</sup> Qatar  Retrospective  Funding: None	Inclusion: Patients with ischemic stroke also diagnosed with COVID-19 (“confirmed” – method not specified)  Exclusion: None report  NOTE: included comparisons with pre-COVID-19 ischemic stroke and negative for COVID-19 ischemic stroke	N=833 (32 COVID-19 positive) Age (years, mean): 54 (COVID-19 group: 49) Gender (% male): 81 (COVID-19 group: 88) Race/ethnicity: NR  Comorbidities: CVD: 12% (COVID-19 group: 13%) CKD: NR COPD: NR DM: 56% (COVID-19 group: 32%) HTN: 72% (COVID-19 group: 41%) Obesity: NR Smoking: 28% (COVID-19 group: 13%)	COVID-19 severity: NR  ICU admission: 44% (COVID-19 group)  Respiratory support Mechanical ventilation or ECMO: 31% (COVID-19 group) NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 11 (COVID-19 group)  Planned/reported time post-hospital (days): 0 (discharge)
Al-Aly, 2021 <sup>104</sup> USA (Veterans)  Retrospective  Funding: VA	Inclusion: Admitted for COVID-19 within 30 days after or 5 days before first positive test and survived at least 30 days after hospital admission; selected from 98,661 patients with positive COVID-19 test between March 01, 2020 and November 30, 2020  Exclusion: None specified  Controls were hospitalized for seasonal influenza and survived 30 days after hospital admission  Propensity scores based on predefined variables were estimated to adjust for potential confounders	N=13,654 (COVID-19 group); N=13,997 (Control group) Age (years, mean): 70 (COVID-19 and Control groups) Gender (% male): 94 (COVID-19 and Control groups) Race/ethnicity: COVID-19 group: White 59%, Black 34%; Control group: White 73%, Black 22%  Comorbidities: NR	COVID-19 severity: NR  ICU admission: 26% (n=3586)  Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay: NR  Planned time post-hospital in patients that survived 30 days after diagnosis (days): 180  Reported time post-hospital (days, median): COVID-19 group: 150, Control group: 157



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Al Kasab, 2020 <sup>23</sup> USA, South America, Europe (28 centers)  Prospective  Funding: None	Inclusion: Consecutive patients undergoing mechanical thrombectomy (MT) for large vessel occlusion; symptomatic patients were tested with RT-PCR methods  Exclusion: None reported	N=13 COVID positive (NOTE: 458 patients underwent MT; 242 were tested for COVID) Age (years, median): 58 Gender (% male): 62 Race: 46% White  Comorbidities: NR	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 39% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, median): 8  Planned/reported time post-hospital (days): 0 (discharge)
Alemanno, 2021 <sup>105</sup>  Italy  Prospective  Funding: none	Inclusion: Positive swab for SARS-CoV-2, stable SatO2 and RR; no need for respiratory assistance or no more than 2 L/min; absence of fever; areas of dependence at the FIM evaluation (FIM score < 100)  Exclusion: Treated for cognitive dysfunctions, under psychotropic drugs prior to recovery, presenting with COVID-19 encephalitis, disease onset <5 days and >20 days  NOTE: separated into 4 groups based on type of respiratory assistance received during acute phase	N=87 (56 at follow-up; Group 1: n=31, Group 2: n=18, Group 3: n=29, Group 4: n=9) Age (years, mean): 67 (Group 1: 60; Group 2: 73; Group 3: 73; Group 4: 63) Gender (% male): 71 Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR  ICU admission: NR  Respiratory Support Mechanical ventilation or ECMO: 36% (Group 1) NIV, HFNC, or CPAP: 21% (Group 2) Other: 33% (Group 3)  Length of hospital stay: NR  Planned time post-hospital (days): 30  Reported time post-hospital: NR
Alharthy, 2021 <sup>24</sup> Saudi Arabia  Prospective  Funding: Hospital	Inclusion: Age >18 years; confirmed serious COVID-19 pneumonia (RT-PCR for SARS-CoV-2); ICU admission  Exclusion: Did not undergo POCUS; 2 consecutive negative RT-PCR results at least 24 hours apart	N=89 Age (years, median): 43 Gender (% male): 84 Race: NR  Comorbidities: NR CVD: CKD: COPD:	COVID-19 severity: 100% severe (dyspnea, respiratory rate ≥30 breaths/min, blood oxygen saturation ≤93%, PaO <sub>2</sub> /FiO <sub>2</sub> <300, development of bilateral pulmonary infiltrates within 24-48 hours, or a combination thereof)  ICU admission: 100%  Respiratory support



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		DM: HTN: Obesity: Smoking:	Mechanical ventilation or ECMO: 84% on ICU admission, 100% within 48 hours (ventilation only); 6% (both) NIV, HFNC, or CPAP: 16% Other: NR  Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)
Alharthy, 2020 <sup>75</sup> Saudi Arabia  Prospective  Funding: Hospital	Inclusion: COVID-19 pneumonia patients admitted to COVID-19 designated ICU; age ≥18, diagnosed with severe COVID-19 (ARF including dyspnea, bilateral pulmonary infiltrates within 24-48 hours)  Exclusion: Did not undergo POCUS for any reason, 2 consecutive negative RT-PCR tests at least 48 hours apart	N=127 survivors (of 171 total patients) Age (years, mean): 45 Gender (% male): 81 Race: NR  Comorbidities: CVD: 12% CKD: 6% COPD: 13% DM: 35% HTN: 49% Obesity: NR Smoking: 12%	COVID-19 severity: 100% severe  ICU admission: 100%  Respiratory support Mechanical ventilation or ECMO: 76% NIV, HFNO or CPAP: 24% Other: 7%  Length of hospital stay (days, mean): 20  Planned time post-hospital (days): 60 and 120  Reported time post-hospital: NR
Anand, 2020 <sup>40</sup> USA  Retrospective  Funding: Foundation	Inclusion: Adults hospitalized with positive PCR testing for SARS-CoV-2 during hospitalization or in the 30 days prior to admission; received either 1) inpatient neurologic or neurocritical care admission or 2) inpatient neurologic or neurocritical care consultation any time during the study period  Exclusion: None reported	N=74 (of 921 adults hospitalized during study period) Age (years, median): 64 Gender (% male): 57 Race/ethnicity: 51% Black or African-American, 30% unknown/declined, 22% Hispanic or Latino, 18% White, 1% Asian  Comorbidities: CAD: 9% CKD: 27% COPD: NR DM: 39%	COVID-19 severity: NR  ICU admission: 46%  Respiratory support Mechanical ventilation or ECMO: 15% NIV or HFNC, or CPAP: NR Other: 38%  Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
		HTN: 58% Obesity: NR Smoking: 32%	
Arab-Zozani, 2020 <sup>76</sup> Iran  Cross-sectional  Funding: Not reported	Inclusion: Discharged from a hospital dedicated to treatment of patients with COVID-19  Exclusion: None reported	N=409 Age (years, mean): 58 Age categories: 7% <=40 years, 26% 41-50 years, 41% 51-60 years, 26% >60 years Gender (% male): 60 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 64% HTN: 60% Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: 18%  Respiratory support: NR  Length of hospital stay (days, mean): 8  Planned time post-hospital: NR  Reported time post-hospital (days, mean): 22
Atalla, 2020 <sup>136</sup> USA  Retrospective  Funding: Not reported	Inclusion: Discharged from hospital; confirmed COVID-19 (RT-PCR for SARS-CoV-2); criteria for hospital admission were individualized – patients with significant comorbidities and moderate to severe COVID-19 (requiring O2 and having abnormal imaging findings) were admitted  Exclusion: None reported  NOTE: 68% of patients (13/19) were admitted for COVID-19; 32% (6/19) admitted for other conditions then developed symptoms and tested positive NOTE: Patients discharged were instructed to seek medical care for relapse of fever, shortness of breath, neurological or thrombotic events, or any change in clinical status; patients received a post-	N=339 (n=19 readmitted, n=320 not readmitted) Age (years, median): 61 Gender (% male): 56 Race: 37% Hispanic, 1% Asian, 16% African American, 43% Caucasian, 3% Other  Comorbidities: CVD: NR CKD: 11%; P=NS between groups COPD: 15% Readmitted: 58%, Not Readmitted: 13%; P<.001 DM: 33% Readmitted: 58%, Not Readmitted: 32%; P=.021 HTN: 45% Readmitted: 68%, Not Readmitted: 44%; P=.038 Obesity: 40%; P=NS between groups Smoking: NR	COVID-19 severity: NR  ICU admission: 33%  Respiratory support Mechanical ventilation or ECMO: 19% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, median): 7 (IQR 4-15)  Planned time post-hospital (days): 30  Reported time post-hospital: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
	discharge follow-up call to monitor recovery process		
Ayoubkhani, 2021 <sup>106</sup>  United Kingdom  Retrospective  Funding: none	Inclusion: Hospitalized for COVID-19, (positive laboratory test or clinical diagnoses) from January 1, 2020 to end of August 2020  Exclusion: Not discharged alive by August 31, 2020 or birth date or gender unknown  Controls were individuals in general population, did not meet inclusion criteria for COVID-19, and had not died before January 1, 2020; 79% had prior hospital admission  Patients and controls matched (1:1) on several confounding variables; all were active patients in National Health Service	N=47,780 (for both COVID-19 group and matched control group) Age (%): COVID-19 group Age <30: 5; 30-49: 16; 50-69: 33; ≥70: 46 Control group <30: 3; 30-49: 19; 50-69: 33; ≥70: 46 Gender (% male): 55 (COVID-19 and Control groups) Race/ethnicity: White 72%, Asian 9%, Black 5% (COVID-19 and Control groups)  Comorbidities: MACE: 24% (COVID-19 and Control groups) CKD: 13% (COVID-19 and Control groups) COPD: COVID-19 group: 14%; Control group: 12% DM: 24% (COVID-19 and Control groups) HTN: 52% (COVID-19 and Control groups) Obesity (BMI ≥30): 32% (COVID-19 and Control groups) Smoking: 8% current, 41% former (COVID-19 and Control groups)	COVID-19 severity: NR  ICU admission: 10% (n=4745)  Respiratory support: NR  Length of hospital stay: NR  Planned time post-hospital: NR  Reported time post-hospital (days, mean): COVID-19 group: 140, Control group: 153
Barbaro, 2020 <sup>41</sup> Multi (International Registry)  Retrospective cohort  Funding: None	Inclusion: Age ≥16, had ECMO support and entered in ELSO registry, laboratory-confirmed COVID-19  Exclusion: No completed COVID-19 addendum; previous ECMO (before COVID-19 diagnosis)	N=1035 (588 discharged alive) Age (years, median): 49 Gender (% male): 74 Race: 34% Black, 33% White (non-Hispanic), 21% Hispanic, 15% Asian, 3% Middle Eastern or North African, 13% Other/Unknown/Multiple  Comorbidities: CVD: 2% CKD: 2% COPD: 3% DM: 24% HTN: NR Obesity: 47%	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 100% (inclusion criteria) NIV, HFNC, or CPAP: 66% (pre-ECMO) Other: NR  Length of hospital stay (days, median): 31  Planned/reported time post-hospital (days): 0 (discharge)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		Smoking: NR	
Bellan, 2021 <sup>107</sup> Italy  Prospective  Funding: Foundation	Inclusion: Patients aged 18 years or older who were discharged after they had been admitted for COVID-19  Exclusion: None reported	N=238 Age (years, median): 61 Gender (% male): 60 Race/ethnicity: NR  Comorbidities: CVD: 4% CKD: 6% COPD: 6% DM: 15% HTN: 41% Obesity: 11% Smoking: 11%	COVID-19 severity: NR  ICU admission: 12%  Respiratory support Mechanical ventilation or ECMO: 9% NIV, HFNC, or CPAP: 21% Other: 43%  Length of hospital stay (days, median): 8.5  Planned time post-hospital (days): 120  Reported time post-hospital (days): 90-120
Benussi, 2020 <sup>25</sup> Italy  Retrospective cohort  Funding: None	Inclusion: Adult (≥18 years) admitted primarily for neurological disease; had outcome of discharge (home or rehabilitation facility) or death; SARS-CoV-2 detected by RT-PCR methods; confirmed COVID-19  Exclusion: None reported  NOTES: reporting data only for patients with <i>cerebrovascular disease</i> on admission; included non-COVID controls	N=111 (43 with COVID-19; 68 non-COVID-19) Age (years, mean): 76 Gender (% male): 56 Race: NR  Comorbidities: CVD: 14% CKD: 5% COPD: NR DM: 22% HTN: 69% Obesity: NR Smoking: 6%	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC, or CPAP: 36% Other: NR  Length of hospital stay (days, median): 5  Planned/reported time post-hospital (days): 0 (discharge)
Bhatt, 2021 <sup>42</sup> USA  Retrospective  Funding: Not reported	Inclusion: History of heart failure (ICD-10 codes) subsequently hospitalized during pandemic period; hospitalizations categorized as related to acute heart failure, COVID-19, or other reasons; COVID-19 determined by ICD-10 code U07.1  Exclusion: None reported	N=8,383 COVID-19 hospitalizations (2,041,855 total hospitalizations during study period) Age (mean, years): 72 Gender (% male): 50 Race/ethnicity: <1% Black Hispanic, 23% Black non-Hispanic, 5% White Hispanic, 41% White non-Hispanic, 32% Other/Unknown  Comorbidities:	COVID-19 severity: NR  ICU admission: 29%  Respiratory support Mechanical ventilation or ECMO: 17% NIV, HFNC, or CPAP: NR Other: NR




Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
	NOTE: included non-COVID-19 hospitalizations group	CVD: 17% CKD: 60% COPD: 42% DM: 61% HTN: 84% Obesity: 29% Smoking: 44%	Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)
Boari, 2021, Boari 2020 (baseline characteristics) <sup>108,162</sup> Italy  Prospective  Funding: University	Inclusion: Confirmed COVID-19 infection (positive RT-PCR assay); bilateral pulmonary interstitial opacities on chest imaging not fully explained by congestive heart failure or other forms of volume overload; acute respiratory distress syndrome showing ≥1 of the following conditions: respiratory rate ≥30 breaths/min; peripheral capillary oxygen saturation (SpO <sub>2</sub> ) ≤93% on ambient air or ratio of partial pressure of oxygen in arterial blood to fractional concentration of oxygen in inspired air (PaO <sub>2</sub> /F <sub>IO</sub> 2) ≤300 mmHg  Exclusion: None reported	N=258 (94 COVID-19 follow-up) Age (years, mean): 71 Gender (% male): 67 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: NR COPD: 14% DM: 26% HTN: 59% Obesity: 22% Smoking: 16%	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay: NR  Planned time post-hospital (days):120  Reported time post-hospital (days): NR
Bowles, 2020 <sup>77</sup> USA  Retrospective cohort  Funding: None	Inclusion: Admitted to home health care after hospitalization for laboratory-confirmed COVID-19; referred from a hospital  Exclusion: None reported	N=1409 Age (years, mean): 67 Gender (% male): 51 Race: 27% Non-Hispanic White, 28% Non-Hispanic Black, 35% Hispanic, 9% Other  Comorbidities: CVD: 1% CKD: NR COPD: 16% DM: NR HTN: NR Obesity: 9% Smoking: NR	COVID-19 severity: NR  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR  Planned time from hospital discharge <u>to first home health visit</u> (days): NR  Reported time from hospital discharge <u>to first home health visit</u> (days): 2.4



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Brendish, 2020 <sup>78</sup> United Kingdom  Prospective cohort  Funding: Foundation, government	Inclusion: Age ≥18, acute respiratory illness or clinically suspected of having COVID-19, molecular point-of-care testing or Rt-PCR  Exclusion: None reported  Patients were part of a clinical trial of molecular point-of-care testing	N=1054 (352 COVID-19 positive, 702 COVID-19 negative) Age (years, median): Positive: 68 Negative: 69 Gender (% male): Positive: 57 Negative: 52 Race: Positive: 74% White, <1% Mixed, 13% Asian, 4% Black Negative: 90% White, 1% Mixed, 3% Asian, 1% Black  Comorbidities- positive, negative: CVD: Positive 36%, negative 40% CKD: Positive 12%, negative 12% COPD: Positive 13%, negative 19% DM: Positive 26%, negative 22% HTN: Positive 41%, negative 40% Obesity: NR Smoking: Positive 6%, negative 17%	COVID-19 severity: NR  ICU admission: Positive: 18% Negative: 6%  Respiratory support Mechanical ventilation or ECMO: COVID Positive: 11%; COVID Negative: 3% NIV, HFNC, or CPAP: COVID Positive: 24%; COVID Negative 6% Other: Oxygen support - COVID Positive 71%; COVID Negative 41%  Length of hospital stay (days, median): COVID Positive: 7.2; COVID Negative 3.7  Planned time post-hospital (days): 30  Reported time post-hospital: NR
Brosnahan, 2020 <sup>79</sup> USA  Retrospective  Funding: Not reported	Inclusion: Confirmed COVID-19, discharged and re-presented with concerns for thrombotic event – included DVT, PE, limb ischemia due to arterial thrombosis, acute coronary syndromes due to coronary thrombosis, acute stroke, rapidly evolving hemodynamic instability with elevated D-dimer  Exclusion: None reported	N=9 (of 1975 patients discharged during study period) Age (years, median): 74 Gender (% male): 67 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: 22% COPD: NR DM: 22% HTN: 33% Obesity: 33% Smoking: NR	COVID-19 severity: NR  ICU admission: NR  Respiratory support: NR  Length of hospital stay (days, mean): 3  Planned time from discharge to re-presenting: Not applicable  Reported time from discharge to re-presenting (days, mean): 5
Casas-Rojo, 2020 <sup>56</sup>	Inclusion: Spanish Society of Internal Medicine registry; age ≥18 years; first	N=15,111 Age (years, median): 69	COVID-19 severity: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Spain  Retrospective cohort  Funding: Foundation/Society	hospital admission; hospital discharge or in-hospital death (consecutive patients with confirmed SARS-CoV-2 (positive RT-PCR or positive result on serological testing and compatible clinical presentation) were eligible  Exclusion: Subsequent admissions of the same patient; denial or withdrawal of consent	Gender (% male): 57 Race: 90% Caucasian, 10% Other  Comorbidities: CVD: 20% CKD (moderate-severe): 6% COPD: 7% DM: 19% HTN: 51% Obesity (BMI $\geq 30$ kg/m <sup>2</sup> ): 21% Smoking: 69% Never, 25% Moderate, 5% Current  NOTE: 4% were healthcare workers	ICU admission: 8%  Respiratory support Mechanical ventilation or ECMO: 7% NIV, HFNC, or CPAP: 13% Other: NR  Length of hospital stay (days, median): 10 (range 1-62, discharged patients)  Planned time post-hospital (days): 30  Reported time post-hospital (days, median): 40 (range 0-102)
Chan, 2021 <sup>102</sup> USA  Retrospective  Funding: Several authors report funding; unclear if related to manuscript	Inclusion: Age $\geq 18$ , laboratory-confirmed SARS-CoV-2 and COVID-19 admitted to 1 of 5 Mount Sinai Health System hospitals 2/27/20-5/30/20  Exclusion: Known end-stage kidney disease prior to admission; hospitalized <48 hours, missing laboratory and vital signs during hospitalization	N=3,993 (demographics for all patients admitted; 3,869 [97%] were discharged) (NOTE: 46% (1,835/3,993) experienced AKI while hospitalized) Age (years, median): 64 Gender (% male): 57 Race: White 24%, Black 36%, Hispanic 26%, Asian 4%, Other or unknown 19%  Comorbidities: CVD: NR CKD: 11% COPD: NR DM: 26% HTN: 38% Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: 24% (976/3993)  Respiratory support Mechanical ventilation or ECMO: 23% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, median): AKI group 10, no AKI group 7 (P<.001, discharged patients only)  Planned time post-hospital: NR  Reported time post-hospital (days, median): 21
Chevinsky 2021 <sup>109</sup> USA  Retrospective  Funding: Not reported	Inclusion: Hospitalized for COVID-19 (ICD-10 code) from March 1 to June 30, 2020  Exclusion: Patients with at least 1 encounter preceding index encounter or who died or were pregnant in index encounter	N=27,284 adults for both COVID-19 and Control groups Age (%): COVID-19 group Age 18-39: 9; 40-49: 10; 50-64: 28; $\geq 65$ : 53 Control group Age 18-39: 11; 40-49: 9; 50-64: 27; $\geq 65$ : 54	COVID-19 severity: NR  ICU admission: both groups 40%  Respiratory support: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
	<p>Controls were hospitalized individuals who did not meet inclusion criteria for COVID-19 and were not diagnosed with COVID-19 during the 4 months after index encounter</p> <p>Patients and controls matched (1:1) based on propensity scores on several confounding variables</p>	<p>Gender (% male): COVID-19 group: 48; Control group: 47 Race/ethnicity: COVID-19 group: White 48%, Black 26%, Asian 2%, Hispanic 13% Control group: White 47%, Black 26%, Asian 2%, Hispanic 14%</p> <p>Comorbidities: NR</p>	<p>Length of hospital stay (days, median): COVID-19 group 6 (range 3, 11); Control group 4 (range 2, 6)</p> <p>Planned time post-hospital (days): 30-120</p> <p>Reported time post-hospital (days): NR</p>
<p>Chopra, 2020<sup>80</sup> USA</p> <p>Retrospective</p> <p>Funding: Health Insurance industry</p>	<p>Inclusion: Hospitalized with COVID-19 (unclear if confirmed), discharged</p> <p>Exclusion: None reported</p>	<p>N=1250 Age (years, median): 62 Gender (% male): 52 Race/ethnicity: 52% Black, 37% White, 11% Unknown</p> <p>Comorbidities: CVD: 24% CKD (moderate/severe): 23% COPD: 10% DM: 35% HTN: 64% Obesity: NR Smoking: NR</p>	<p>COVID-19 severity: NR</p> <p>ICU admission: 13%</p> <p>Respiratory support Mechanical ventilation or ECMO: 6% NIV, HFNC, or CPAP: NR Other: 69%</p> <p>Length of hospital stay (days, median): 5</p> <p>Planned time post-hospital (days): 60</p> <p>Reported time post-hospital (days): NR</p>
<p>Collins, 2020<sup>26</sup> USA</p> <p>Retrospective</p> <p>Funding: University</p>	<p>Inclusion: Persons with HIV admitted with COVID-19 (detection of SARS-CoV-2 via RT-PCR)</p> <p>Exclusion: None reported</p> <p>NOTE: study sites included Atlanta Veterans Affairs Medical Center</p>	<p>N=20 Age (years, median): 57 Gender (% male): 65 Race: 85% Non-Hispanic Black, 5% Non-Hispanic White, 5% Non-Hispanic/Multiracial, 5% Hispanic/Latino</p> <p>Comorbidities: CVD: 30% CKD: 25% Chronic lung disease: 30% DM: 45% HTN: 70% Obesity: 30%</p>	<p>COVID-19 severity: NR</p> <p>ICU admission: 30%</p> <p>Respiratory support Mechanical ventilation or ECMO: 15% NIV, HFNC, or CPAP: 15% Other: 25%</p> <p>Length of hospital stay (days, median): 5</p> <p>Planned/reported time post-hospital (days): 0 (discharge)</p>



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		Smoking: 50% Current, 10% Former, 40% Never	
Curci, 2020 <sup>27</sup> Italy  Cross-sectional  Funding: None	<p>Inclusion: Consecutive referrals to rehabilitation unit; adults (&gt;18); diagnosis of viral interstitial lung disease (CT); positive for SARS-CoV-2 (RT-PCR); previously hospitalized in ICU; clinical stability (able to perform bedside mobilization without reduction in oxygen saturation below 90%); complete weaning from sedative and antipsychotic drugs</p> <p>Exclusion: Respiratory distress signs; cognitive impairment; need of respiratory support (FiO<sub>2</sub> &gt;60%); need of CPAP devices; signs of cardiovascular instability</p>	<p>N=32*</p> <p>Age (years, mean): 73 Gender (% male): 69 Race: NR</p> <p>Comorbidities: CVD: NR CKD: NR COPD: 6% DM: 19% HTN: 63% Obesity: NR Smoking: 28%</p> <p>*Subgroups 1) FiO<sub>2</sub> ≥21% and &lt;40% (n=13); without oxygen support devices or wearing nasal cannula 2) FiO<sub>2</sub> ≥40% and &lt;60% (n=19); wearing non-rebreather mask, Venturi mask, or oxygen mask</p>	<p>COVID-19 severity: NR</p> <p>ICU admission: 100%</p> <p>Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC or CPAP: NR Other: 47%</p> <p>Length of hospital stay (days, mean): 16.4 (ICU admission to rehabilitation unit)</p> <p>Planned/<u>reported time to rehabilitation unit admission</u> (days): 0 (discharge from ICU)</p>
Daher, 2020 <sup>81</sup> Germany  Prospective  Funding: None	<p>Inclusion: Consecutive patients hospitalized due to COVID-19 (confirmed by RT-PCR)</p> <p>Exclusion: Patients with ARDS who needed mechanical ventilation in the ICU</p>	<p>N=33</p> <p>Age (years, mean): 64 Gender (% male): 67 Race/ethnicity: NR</p> <p>Comorbidities: CVD: 19% CKD: 22% COPD: 9% DM: 25% HTN: 59% Obesity: NR Smoking: NR</p>	<p>COVID-19 severity: 100% (criteria NR)</p> <p>Symptom onset to hospitalization: 6 days</p> <p>ICU admission: NR</p> <p>Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC, or CPAP: NR Other: 82%</p> <p>Length of hospital stay (days, mean): 15</p> <p>Planned time post-hospital: NR</p> <p>Reported time post-hospital (days, median): 42 (range 48-71)</p>



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
<p>Daugherty 2021<sup>110</sup> USA</p> <p>Retrospective</p> <p>Funding: Insurance (Research &amp; Development)</p>	<p>Inclusion: Ages 18-65 diagnosed with COVID-19 (SARS-CoV-2); continuous enrollment in the health plan from January 1, 2019 to index date (defined by first of: 1) primary, secondary, or tertiary diagnosis of COVID-19; 2) administrative claims with ICD-10 codes U07.1 or either B34.2 or B97.29 before April 1, 2020; 3) documentation of positive PCR test in outpatient laboratory dataset; or 4) admitted to hospital for COVID-19 (based on diagnostic code))</p> <p>Exclusion: Positive SARS-CoV-2 antibodies but without documented infection; ICD-10 codes B34.2 or B97.29 on or after April 1, 2020; and admitted to hospital for suspected COVID-19 but missing diagnostic codes</p> <p>Controls were ages 18-65 without COVID-19 (SARS-CoV-2) diagnosis with continuous health plan enrollment from January 1 2019 to randomly assigned month and day drawn from the SARS-CoV-2 infection group (2020 comparator group used for analysis of hospitalized patients)</p> <p>Patients and controls matched (1:1) using propensity scores based on 108 variables</p>	<p>N=21,746 hospitalized (N=18,118 for both COVID-19 and control groups in matched analysis after exclusion if less than index date + 21 days of follow-up); demographics and comorbidities NR for hospitalized subgroup</p> <p>Age (years, mean): NR Gender (% male): NR Race/ethnicity: NR</p> <p>Comorbidities: NR</p>	<p>COVID-19 severity: NR</p> <p>ICU admission: 13% (n=2933)</p> <p>Respiratory support: NR</p> <p>Length of hospital stay: NR</p> <p>Planned time <u>post-acute infection</u>* (days): 90-180</p> <p>Reported time <u>post-acute infection</u>* (days, mean): 120</p> <p>NOTE: post-acute infection defined as index date plus 21 days</p>
<p>Dawson, 2020<sup>143</sup> United Kingdom</p> <p>Prospectively collected/ retrospectively analyzed</p>	<p>Inclusion: Admitted with SARS-CoV-2 (RT-PCR confirmed); referred to Speech and Language Therapy team with clinical signs of dysphagia</p> <p>Exclusion: None reported</p>	<p>N=208</p> <p>Age (years, mean): 68 Gender (% male): NR Race/ethnicity: NR</p> <p>Comorbidities: NR</p>	<p>COVID-19 severity: NR</p> <p>ICU admission: 49% (102/208) admitted for ventilatory support</p> <p>Respiratory support Mechanical ventilation or ECMO: NR</p>



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Funding: Not reported			NIV, HFNC, or CPAP: NR Other: 39% (tracheostomy)  Length of hospital stay: NR  Planned/reported time post-hospital: NR
DeBolt, 2020 <sup>43</sup> USA  Retrospective case-control  Funding: Not reported	Inclusion: Ages 18-50; confirmed (RT-PCR) SARS-CoV-2 infection meeting admission criteria for severe and critical COVID-19; pregnant women were admitted for COVID-19 and not obstetrical indications  Exclusion: Inconclusive or negative SARS-CoV-2 laboratory results; comorbidities associated with immunocompromised state (active malignancy, history of transplant or developmental delay, cerebral palsy, trisomy 21 or other aneuploidies)  Controls were non-pregnant women of reproductive age admitted for COVID-19	N=38 pregnant cases; 94 non-pregnant controls Age (mean, years) Cases: 35; Controls: 38 (P<.01) Gender (% male): 0 Race/ethnicity Cases: 24% Non-Hispanic White, 18% Non-Hispanic Black, 40% Hispanic, 8% Asian, 0% American Indian or Alaskan Native, 10% Other or Unknown Controls: 10% Non-Hispanic White, 14% Non-Hispanic Black, 37% Hispanic, 9% Asian, 1% American Indian or Alaskan Native, 30% Other or Unknown  Comorbidities: CVD: NR CKD: NR COPD: Cases 11%, controls 29% DM: Cases 11%, controls 28% HTN: NR Obesity: NR Smoking: Cases 0%, controls 0%	COVID-19 severity (WHO and Chinese CDC criteria) Severe: 76% Cases; 85% Controls Critical: 24% Cases; 15% Controls (P<.01)  ICU admission: 40% Cases; 17% Controls  Respiratory support Mechanical ventilation or ECMO: 26% Cases; 11% Controls NIV, HFNC, or CPAP: 29% Cases; 10% Controls Other: NR  Length of hospital stay (days, mean): 9 (Cases); 7 (Controls)  Planned/reported time post-hospital (days): 0 (discharge)
de Graaf, 2021 <sup>111</sup> the Netherlands  Prospective  Funding: None	Inclusion: All adults (≥18) from geographic region discharged after admission for PCR-confirmed SARS-CoV-2 infection; planned for outpatient clinic after discharge  Exclusion: None reported	N=81 Age (years, mean): 61 Gender (% male): 63 Race/ethnicity: NR  Comorbidities: CVD: 28% CKD: 11% COPD: 4% DM: 23%	COVID-19 severity: NR  ICU admission: 42%  Respiratory support Mechanical ventilation or ECMO: 41% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 17



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		HTN: 34% Obesity: NR Smoking: 11%	Planned time post-hospital (days): 42  Reported time post-hospital (days): NR
de Havenon, 2021 <sup>146</sup> USA  Retrospective  Funding: Government	Inclusion: ICD-10 codes for ischemic stroke in discharge diagnoses, comorbid COVID-19 during same hospitalization (identified by ICD code U07.1 [laboratory testing confirmed])  Exclusion: Elective hospital admissions, patients on hospice prior to admission  NOTE: included historical controls	N=2086 COVID-19 positive cases, 166,586 pneumonia controls Age category: Cases: 12% 18-50 years; 29% 51-64 years; 29% 65-74 years; 11% 75-79 years; 19% ≥80 years Controls: 11% 18-50 years; 26% 51-64 years; 25% 65-74 years; 12% 75-79 years; 26% ≥80 years  Gender (% male): 58 cases, 51 controls Race/ethnicity: Cases: 34% White, 32% Black, 19% Hispanic, 5% Asian, 11% Other/Unknown; Controls: 62% White, 22% Black, 7% Hispanic, 3% Asian, 6% Other/Unknown  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 55% cases, 40% controls HTN: 67% cases, 73% controls Obesity: 25% cases, 17% controls Smoking: 5% cases, 16% controls	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 44% cases, 12% controls NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 18 cases, 8 controls  Planned/reported time post-hospital (days): 0 (discharge)
De Lorenzo, 2020 <sup>82</sup> Italy  Part of COVID-BioB study  Prospective and retrospective	Inclusion: Age ≥18, admitted to emergency department, confirmed (RT-PCR) SARS-CoV-2, follow-up at designated COVID-19 Outpatient Clinic  Exclusion: Admitted for reasons other than COVID-19 who subsequently tested positive for SARS-CoV-2	N=126 (hospitalized patients) Age (years, mean): 61 Gender (% male): 73 Race/ethnicity: 94% European, 6% Hispanic  Comorbidities: CAD: 6% CKD: 2% COPD: 2% DM: 14%	COVID-19 severity: NR  ICU admission: 3% (4/126)  Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC, or CPAP: 25% Other: NR  Length of hospital stay (days, median): 10



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
Funding: None		HTN: 44% Obesity: NR Smoking: NR	Planned time post-hospital: NR  Reported time post-hospital (days, median): 22
De Michieli, 2021 <sup>112</sup> USA  Retrospective  Funding: Government	Inclusion: Consecutive adult patients with confirmed COVID-19 diagnosis (RT-PCF); presented to emergency department and/or were admitted  Exclusion: No Minnesota Research Authorization form or permission to use records, <18 years, indeterminate PCR results, no hs-cTnT during index hospitalization	N=367 Age (years, mean): 61 Gender (% male): 60 Race/ethnicity: White 66%  Comorbidities: CAD: 13% CKD: 21% COPD: 16% DM: 32% HTN: 58% Obesity: 41% Smoking: 36% (current)	COVID-19 severity: NR  ICU admission: 28%  Respiratory support Mechanical ventilation or ECMO: 16% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, median): 7  Planned time post-hospital (days): 30  Reported time post-hospital (days, median): 49
Dennis, 2021 <sup>57</sup> United Kingdom  Prospective  Funding: Government	Inclusion: Positive for SARS-CoV-2 by RT-PCR (n=62), positive antibody test (n=63), or determined to have COVID-19 by 2 independent clinicians based on symptoms (n=73)  Exclusion: Symptoms of active respiratory viral infection; discharged from hospital in last 7 days; contraindications to MRI (metallic implanted devices, claustrophobia)	N=37 (patients hospitalized only) Age (years, mean): 50 Gender (% male): 38 Race/ethnicity: 76% White, 8% South Asian, 5% Black  Comorbidities: Previous heart disease: 3% CKD: NR COPD: NR DM: 0% HTN: 5% Obesity: NR Smoking: 65% never, 35% former, 0% current  NOTE: 35% were health care workers	COVID-19 severity: NR  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR  Planned time <u>post-positive test</u> : NR  Reported time <u>post positive test</u> (days, median): 105  NOTE: Organ function by patient-reported questionnaires, fasting blood investigations, and multi-organ MRI

Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Doher, 2020 <sup>145</sup> Brazil  Retrospective cohort  Funding: None	Inclusion: Age ≥18, confirmed severe COVID-19, admitted to ICU, positive RT-PCR  Excluded: CKD on dialysis	N=201 (101 diagnosed with AKI) Age (years, median): 64 (AKI: 73, non-AKI: 60) Gender (% male): 61 (AKI: 66 non-AKI: 56) Race/ethnicity: NR  Comorbidities: CAD: 8% (13% AKI, 3% non-AKI) CKD: NR COPD: NR DM: 32% (39% AKI, 25% non-AKI) HTN: 49% (58% AKI, 39% non-AKI) Obesity: NR Smoking: 3% (2% AKI, 4% non-AKI)	Time from symptoms to hospital admission: 6 days (median)  COVID-19 severity: NR  ICU admission: 100% (inclusion criteria)  Respiratory support (at ICU admission) Mechanical ventilation or ECMO: 44% (71% AKI, 17% non-AKI) NIV, HFNC, or CPAP: 61% (64% AKI, 57% non-AKI) Other: NR  Length of hospital stay (days, median): 18 AKI, 10 Non-AKI  Planned/reported time post-hospital (days): 0 (discharge)
Egol, 2020 <sup>58</sup> USA  Prospective  Funding: None	Inclusion: Hip fracture; positive RT-PCR test before, during, or after (at rehabilitation) hospitalization  Exclusion: None reported  NOTE: Included comparison data from COVID-19 Suspected and COVID-19 Negative patients	N=17 (COVID-19 positive) Age (years, mean): 82 Gender (% male): 71 Race/ethnicity: 82% White, 0% African American, 12% Hispanic, 6% Asian  Comorbidities: CVD: 47% CKD: 24% (renal failure) COPD: 18% DM: 41% HTN: 65% Obesity: NR Smoking: 53% Never, 35% Former, 12% Current	COVID-19 severity: NR  ICU admission: 29%  Respiratory support Mechanical ventilation or ECMO: 12% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, means): 9.8  Planned time post-hospital (days): 30  Reported time post-hospital (days): NR
El Moheb, 2020 <sup>59</sup> USA  Retrospective	Inclusion: All patients with confirmed SARS-CoV-2 (RT-PCR) who were intubated and admitted to ICU	N=92 (propensity matched subgroup with COVID-19 ARDS) Age (years, median): 62 Gender (% male): 59	COVID-19 severity: NR  ICU admission: 100% (inclusion criteria)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
Funding: Not reported	Exclusion: None reported  NOTE: analysis was limited to patients whose gastrointestinal complications while hospitalized were previously reported; propensity score matching with to identify comparably ill patients with non-COVID-19 ARDS	Race/ethnicity: NR  Comorbidities: CAD: 13% CKD: 20% Chronic lung disease: 29% DM: 37% HTN: 55% Obesity: NR Smoking: 39%	Respiratory support: NR  Length of hospital stay (days, median): 24  Planned/reported time post-hospital: NR
Engelen, 2021 <sup>113</sup> Belgium  Prospective  Funding: None	Inclusion: Age 75 or younger and hospitalized with confirmed COVID-19  Exclusion: Residents of medical care facilities, patients with cognitive impairment, or with a clinical frailty scale greater than 5, patients admitted for non-respiratory reasons with incidental finding of SARS-CoV-2 or patients with hospital stay <2 days	N=146 Age (years, mean): 58 Gender (% male): 62 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: 20% COPD: NR DM: 29% HTN: 45% Obesity: NR Smoking: 44%	COVID-19 severity: NR  ICU admission: 39%  Respiratory support Mechanical ventilation or ECMO: 28% NIV, HFNC, or CPAP: NR Other: 88%  Length of hospital stay (days, median): 11  Planned time post-hospital (days): 42  Reported time post-hospital (days): NR
Eswaran, 2021 <sup>114</sup> USA Retrospective Funding: none	Inclusion: Patients hospitalized with confirmed COVID-19  Exclusion: Patients who died, were discharged to hospice or comfort care, screened positive when hospitalized for an unrelated condition, or were discharged on therapeutic anticoagulation for another indication.	N=447 Age (years, mean): 54 Gender (% male): 51 Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR  ICU admission: 39%  Respiratory support: NR  Length of hospital stay (days, mean): 8  Planned time post-hospital (days): 30  Reported time post-hospital (days): NR
Fisher, 2020 <sup>28</sup> USA	Inclusion: Age >18 years with COVID-19 test performed upon hospitalization;	N=3,345 (positive for COVID-19; total of 4,610 were eligible and tested)	COVID-19 severity: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Retrospective  Funding: None	<p>confirmed case of COVID-19 was a positive RT-PCR result</p> <p>Exclusion: Age &lt;18 years; end stage kidney disease; no creatinine values; unknown sex assignment</p> <p>NOTE: included comparison group of patients hospitalized during same time period in 2019</p>	<p>Age (years, mean): 64 Gender (% male): 53 Race: 8% Non-Hispanic White; 36% Non-Hispanic Black, 37% Hispanic; 19% Other</p> <p>Comorbidities: CVD: NR CKD: 12% COPD: NR DM: 27% HTN: NR Obesity: 43% Smoking: NR</p> <p>NOTE: 16% were nursing home residents</p>	<p>ICU admission: 13%</p> <p>Respiratory support Mechanical ventilation or ECMO: 18% NIV, HFNC, or CPAP: NR Other: NR</p> <p>Length of hospital stay (days, median): 5</p> <p>Planned/reported time post-hospital (days): 0 (discharge)</p>
Frija-Masson, 2020 <sup>60</sup> France  Retrospective  Funding: Not reported	<p>Inclusion: Age &lt;85; confirmed SARS-CoV-2 infection (RT-PCR); discharged from hospital; evaluated with pulmonary function tests 30 days after symptom onset as part of routine care</p> <p>Exclusion: Decline to participate; recurrence; patients with ARDS</p> <p>NOTE: 18% (9/50) treated as outpatients</p>	<p>N=50 Age (years, median): 54 Gender (% male): 56 Race: NR</p> <p>Comorbidities: CVD: NR CKD: NR COPD: NR DM: 16% HTN: 48% Obesity: NR Smoking: 10% active; 18% former</p>	<p>COVID-19 severity: 50% severe (based on CT)</p> <p>ICU admission: 14% (7/50)</p> <p>Respiratory support Mechanical ventilation or ECMO: 2% NIV, HFNC, or CPAP: 8% Other: 50%</p> <p>Length of hospital stay: NR</p> <p>Planned time <u>post-infection</u> (days): 30</p> <p>Reported time <u>post-infection</u>: NR</p>
Fuglebjerg, 2020 <sup>29</sup> Denmark  Case series  Funding: None	<p>Inclusion: Hospitalized with COVID-19 confirmed by PCR testing</p> <p>Exclusion: Chronic lung diseases or New York Heart Association (NYHA) class II or above</p>	<p>N=26 Age (years, median): 63 (range 29-85) Gender (% male): 62 Race: NR</p> <p>Comorbidities: NR (patients had a median of 1 (non-specified) per patient)</p>	<p>COVID-19 severity: NR</p> <p>ICU admission: 31%</p> <p>Respiratory support Mechanical ventilation or ECMO: 15% NIV, HFNC, or CPAP: NR Other: NR</p>



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
			Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)
Garrigues, 2020 <sup>61</sup> France  Prospective, survey  Funding: None	Inclusion: Hospitalized in COVID-19 ward; positive SARS-CoV-2 (RT-PCR) and/or typical abnormalities on chest CT  Exclusion: Directly admitted to ICU without being hospitalized in COVID-19 unit; deceased, unreachable by telephone, demented, bedridden, non-French speaking	N=120 Age (years, mean): 63 Gender (% male): 63 Race: NR  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 22% HTN: 47% Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: 20%  Respiratory support Mechanical ventilation or ECMO: 12% NIV, HFNC, CPAP: 14% Other: NR  Length of hospital stay (days, mean): 13  Planned time post-hospital (days): >100  Reported time post-hospital (days, mean): 111
Goicoechea, 2020 <sup>30</sup> Spain  Retrospective  Funding: Not reported	Inclusion: All patients on maintenance hemodialysis admitted with positive RT-PCR testing for SARS-CoV-2 infection  Exclusion: None reported	N=36 (7 were discharged) Age (years, mean): 71 Gender (% male): 64 Race: NR  Comorbidities: CVD: 22% CKD: 100% COPD: 19% DM: 64% HTN: 97% Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: 3% (1/36)  Respiratory support Mechanical ventilation or ECMO: 3% (NOTE: all required oxygen supplement therapy; severe comorbidities in 11 patients requiring assisted mechanical ventilation limited invasive measures) NIV, HFNC, CPAP: 67% Other: 33%  Length of hospital stay (days, median): 11.4 (discharged patients)  Planned/reported time post-hospital (days): 0 (discharge)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Grewal, 2020 <sup>31</sup> USA  Retrospective  Funding: Not reported	Inclusion: Diagnosis of acute ischemic stroke (AIS) (confirmed with MRI or CT); positive for COVID-19 (RT-PCR); divided patients into “COVID” group (initially with COVID-19 symptoms who developed AIS) and “neuro” group (admitted for AIS and tested positive for COVID-19) (NOTE: included control groups of non-COVID-19 AIS patients hospitalized during study time frame and in 2019)  Exclusion: None reported	N=13 (6 in “COVID” group, 7 in “neuro” group) Age (years, mean): 62 Gender (% male): 46 Race: 46% Latino, 31% African-American  Comorbidities: CAD: 15% CKD: NR COPD: NR DM: 69% HTN: 69% Obesity: 15% Smoking: NR	COVID-19 severity: 8 (62%) severe or critical; 5 (38%) mild or regular  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)
Gupta, 2021 <sup>139</sup> USA  STOP-COVID study  Cohort  Funding: None	Inclusion: Age ≥18; consecutively admitted to ICUs at 67 hospitals, laboratory confirmed diagnosis of COVID-19  Exclusion: History of ESKD  NOTE: authors report that 153 of the patients in this study were included in Chan 2020 (above) 7	N=3099 (637 with AKI-RRT, 216 discharged) Age (years, median): 62 Gender (% male): 65 Race/ethnicity: 37% White, 31% Black, 6% Asian, 34% Hispanic, 26% Other/Unknown  Comorbidities: CAD: 13% CKD: 67% eGFR <90 ml/min per 1.73 m <sup>2</sup> COPD: 8% DM: 14% insulin dependent, 26% noninsulin dependent HTN: 60% Obesity: NR Smoking: 30% current or former	COVID-19 severity: NR  ICU admission: 100%  Respiratory support Mechanical ventilation or ECMO: 66% (on ICU admission) NIV, HFNC, or CPAP: 22% Other: NR  Length of hospital stay: NR  Planned time post-hospital (days): 0 (discharge)  Reported time post-hospital (days): 60 after ICU admission
Hall, 2021 <sup>115</sup> United Kingdom  Retrospective  Funding: None	Inclusion: All patients attending the hospital with either a positive PCR for COVID-19 or a clinical radiological diagnosis and had persistent symptoms or required ICU admission	N=200 (179 received inpatient care, 21 discharged directly from the emergency department) Age (years, mean): 55 Gender (% male): 62 Race/ethnicity: NR  Comorbidities:	COVID-19 severity: NR, majority were not reported as “critically unwell”  ICU admission: 39%  Respiratory support Mechanical ventilation or ECMO: 28% NIV, HFNC, or CPAP: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
	Exclusion: Patients who died or were not yet discharged. Patients unable to attend clinic for follow-up due to frailty	CVD: NR CKD: NR COPD: 2% DM: 28% HTN: 36% Obesity: 36% Smoking: 15%	Other: 70%  Length of hospital stay (days, median): 9  Planned time post-hospital (days): 28-42  Reported time post-hospital: NR
Hamilton, 2020 <sup>83</sup> United Kingdom  Retrospective  Funding: None	Inclusion: Age ≥18, tested positive (RT-PCR) for COVID-19, 4 hospitals  Exclusion: Day cases and known hemodialysis patients	N=1032 (210 with AKI) Age (years, median): 71 Gender (% male): 55 Race/ethnicity: 70% White, 11% Asian, 9% Black, 9.3% Mixed/Other/Unknown  Comorbidities: CVD: 4% CKD: 14% COPD: 25% DM: 23% without complications, 4% with complications HTN: NR Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission (critical care): 16% (165/1032)  Respiratory support Mechanical ventilation or ECMO: 75% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 10  Planned time <u>post-COVID positive admission</u> (days): 30  Reported time <u>post-COVID positive admission</u> (days): NR
Han, 2021 <sup>116</sup> China  Prospective  Funding: Government, university	Inclusion: Severe COVID-19 patients discharged from the hospital  Exclusion: NR	N=114 Age (years, mean): 54 Gender (% male): 70 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: NR COPD: 14% DM: 11% HTN: 28% Obesity: 22% Smoking: 14% (history)	COVID-19 severity: 100% severe  ICU admission: 9%  Respiratory support Mechanical ventilation or ECMO: 21% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 17  Planned time <u>post-disease onset</u> (days): 180  Reported time <u>post-disease onset</u> (days, mean): 175




Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
Hegde, 2020 <sup>44</sup> USA  Retrospective case series  Funding: None	Inclusion: Age ≥18, laboratory-confirmed COVID-19 (positive for SARS-CoV-2 by RT-PCR), transthoracic echocardiogram performed during hospital stay with features consistent with Takotsubu cardiomyopathy  Exclusion: None reported	N=7 Age (years, mean): 71 Gender (% male): 57 Race/ethnicity: NR  Comorbidities: CAD: 14% CKD: 14% COPD: NR DM: 71% HTN: 86% Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: 86%  Respiratory support Mechanical ventilation or ECMO: 86% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 18  Planned/reported time post-hospital (days): 0 (discharge)
Hill, 2020 <sup>84</sup> USA  Retrospective  Funding: Not reported	Inclusion: Positive for SARS-CoV-2 (RT-PCR)  Exclusion: Indication for heparin bolus was thrombosis of an extracorporeal circuit	N=2748 (2075 survived to discharge) Age (years, mean): NR Gender (% male): NR Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 23% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay: NR  Planned time post-hospital : NR  Reported time post-hospital (days, mean): 21
Hittesdorf, 2020 <sup>140</sup> USA  Retrospective  Funding: Institution/ Department	Inclusion: Admitted to a provisional ICU, severe COVID-19 (not defined)  Exclusion: Incomplete records or follow-up data	N=116 Age (years, mean): 62 Gender (% male): 65 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 43% HTN: 53%	COVID-19 severity: 100% (criteria NR)  ICU admission: 100%  Respiratory support: Mechanical ventilation: 100% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 60 (survivors)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
		Obesity: 44% Smoking: NR	Planned time post-hospital (days): 0 (discharge) and 90  Reported time post-hospital: NR
Hu, 2020 <sup>45</sup> China  Cross-sectional  Funding: Government	Inclusion: Patients admitted to a hospital designated to treat patients with SARS-CoV-2, diagnosis of COVID-19 based on WHO guidelines and RT-PCR methods  Exclusion: Lack of available blood samples, no CT examination, death	N=76 Age (years, mean): 51 Gender (% male): 45 Race/ethnicity: NR  Comorbidities: CVD: 12% CKD: NR COPD: NR DM: 11% HTN: 22% Obesity: NR Smoking: NR	COVID-19 severity: 17% severe, 83% non-severe  ICU admission: NR  Respiratory support: NR  Length of hospital stay (days, median): 14 (18 [severe], 13 [non-severe])  Planned/reported time post-hospital (days): 0 (discharge)
Huang C, 2021 <sup>85</sup> China  Retrospective and Prospective  Funding: Government, University, Foundation	Inclusion: Adults, laboratory confirmed COVID-19, discharged  Exclusion: Died before follow-up visit; follow-up would be difficult owing to psychotic disorder, dementia, or readmission to hospital attributed to underlying disease; unable to move freely or immobile before or after discharge; declined participation; unable to be contacted; living outside Wuhan (study city) or in nursing or welfare homes	N=1733 Age (years, median): 57 Gender (% male): 52 Race/ethnicity: NR  Comorbidities: CVD: 7% CKD: 2% COPD: 2% DM: 12% HTN: 29% Obesity: NR Smoking: 6% current, 3% former	Time from symptom onset to admission (days, median): 15  COVID-19 severity: NR  ICU admission: 4%  Respiratory support Mechanical ventilation or ECMO: 1% NIV, HFNC, or CPAP: 6% Other: 68%  Length of hospital stay (days, median): 14  Planned time post-hospital (days): 180  Reported time post-hospital (days, median): 153

Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Huang L, 2020 <sup>62</sup> China  Retrospective  Funding: Foundation, Government	Inclusion: Consecutive patients referred for CMR due to cardiac symptoms after discharge; previously confirmed with SARS-CoV-2 (RT-PCR); considered recovered by hospital discharge criteria  Exclusion: History of CAD or myocarditis; contraindication to gadolinium contrast; CMR image quality not sufficient for analysis	N=26 Age (years, median): 38 Gender (% male): 38 Race: NR  Comorbidities: CAD: 0% CKD: 0% COPD: 0% DM: 0% HTN: 8% Obesity: NR Smoking: NR  NOTE: also included data from healthy controls (similar age and gender distribution, no CVD or systemic inflammation) who underwent CMR at same hospital	COVID-19 severity: 0 critical, 4 severe, 22 moderate  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 0% NIV, HFNC, or CPAP: 81% Other: NR  Length of hospital stay: NR  Planned/reported time post-hospital: NR  NOTE: Median time from cardiac symptom onset to CMR was 47 days
Huang Y, 2020 <sup>63</sup> China  Retrospective  Funding: Not reported	Inclusion: Age over 18 years; released from hospital over 1 month; confirmed SARS-CoV-2 infection (RT-PCR)  Exclusion: Previous history of pulmonary resection, neurological disease, or mental illness; could not be contacted or declined participation	N=57 Age (years, mean): Severe: 53; Non-severe: 44; P=.03 Gender (% male): Severe 71%; Non-severe: 35%; P=.01 Race: NR  Comorbidities: CVD: 5% CKD: NR COPD: 0% DM: 7% HTN: 19% Obesity: NR Smoking: 16%	COVID-19 severity: 30% severe, 70% non-severe  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC, or CPAP: NR Other: 30%  Length of hospital stay (days, mean): 21  Planned time post-hospital (days): 30  Reported time post-hospital: NR
Jacobs, 2020 <sup>17</sup> USA  Prospective	Inclusion: Aged 18 or older; diagnosis of viral RNA PCR-confirmed COVID-19 infection during hospitalization; hospitalization duration of at least three days	N=183 Age (years, median): 57 Gender (% male): 62	COVID-19 severity: 95% mild  ICU admission: NR  Respiratory support



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Funding: None	Exclusion: Individuals who had expired during or after hospitalization; non-English speakers; and individuals with a documented diagnosis of dementia or delirium	Race/ethnicity: 54% White, 9% Black, 9% Asian, 1% American Indian/Alaskan, 27% other/non-white  Comorbidities: CVD: 12% CKD: NR COPD: 4% DM: 28% HTN: 48% Obesity: 49% Smoking: NR	Mechanical ventilation or ECMO: 5% NIV, HFNC, or CPAP: 81% Other: 13%  Length of hospital stay (days, median): 7  Planned time post-hospital (days): 35 (±5)  Reported time post-hospital: NR
Karaarslan, 2021 <sup>118</sup> Turkey  Prospective  Funding: None	Inclusion: Aged between 18 and 70 years or older and discharged following hospitalization for COVID-19  Exclusion: Individuals who received intensive care unit care during hospitalization	N=300 Age (years, mean): 53 Gender (% male): 60 Race/ethnicity: NR  Comorbidities: CVD: 15% CKD: 2% COPD: 2% DM: 28% HTN: 32% Obesity: NR Smoking: 7%	COVID-19 severity: NR  ICU admission: None  Respiratory support: NR  Length of hospital stay (days, mean): 8  Planned time post-hospital (days): 14 and 30  Reported time post-hospital: NR
Katz, 2020 <sup>32</sup> USA  Retrospective  Funding: None	Inclusion: Confirmed SARS-CoV-2 infection (RT-PCR); concurrent stroke diagnosis (stroke symptom onset during COVID-19 illness or onset of COVID-19 symptoms or SARS-CoV-2 positivity within 14 days of stroke symptom onset) confirmed by imaging  Exclusion: None reported  NOTE: included control group of all stroke patients admitted 1 year earlier between same dates to same hospitals	N=86 Age (years, mean): 67 Gender (% male): 56 Race: 30% White, 31% Black, 12% Asian, 27% Multiracial/other  Comorbidities: CVD: NR CKD: NR COPD: NR DM: NR HTN: NR Obesity (BMI ≥30 kg/m <sup>2</sup> ): 31%	COVID-19 severity: among n=45 testing positive for COVID-19 after stroke symptoms 51% (23/45) had mild COVID-19 symptoms and 29% (13/45) were asymptomatic  ICU admission: 51% (critical care admission)  Respiratory support Mechanical ventilation or ECMO: 44% NIV, HFNC, or CPAP: 9% Other: 72%  Length of hospital stay: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		Smoking: NR	Planned/reported time post-hospital (days): 0 (discharge)  NOTE: 48% (41/86) had stroke onset during hospitalization for COVID-19
Khalili, 2020 <sup>86</sup> Iran  Prospective cohort  Funding: Not reported	Inclusion: Hospitalized with COVID-19 (confirmed by RT-PCR [76%] or CT findings), included patient with and without diabetes  Exclusion: None reported	N=254 (127 with diabetes) Age (years, mean): 66 Gender (% male): 56 Race/ethnicity: NR  Comorbidities: CVD: 8% CKD: 8% COPD: 3% DM: 50% HTN: 43% Obesity: NR Smoking: 4% current, 2% former	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 11% NIV, HFNC, or CPAP: 89% Other: NR  Length of hospital stay (days, mean): 6 (range 1-60)  Planned time <u>post-admission</u> (days): 90  Reported time <u>post-admission</u> : NR
Knights, 2020 <sup>137</sup> United Kingdom  Retrospective  Funding: Not reported	Inclusion: Admitted to hospital with positive COVID-19 test  Exclusion: None reported	N=108 Age (years, mean): 69 Gender (% male): 58 Race: White British: 76%  Comorbidities: CVD: NR CKD: 6% COPD: 15% DM: 23% HTN: 45% Obesity: 31% Smoking: 44% Ex or current, 56% never  NOTE: 10% were care home residents; 21% had a “package of care”	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 8% NIV, HFNC, or CPAP: 24% Other: 55%  Length of hospital stay (days, median): 8 [IQR 4-13]  Planned time <u>post-admission to study end point</u> (days): NR  Reported time <u>post-admission to study end point</u> (days, median): 26



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Li, 2021 <sup>119</sup> China  Prospective  Funding: Government	Inclusion: Patients admitted to the hospital for COVID-19  Exclusion: NR	N=107 Age (years, mean): 66 Gender (% male): 50 Race/ethnicity: NR  Comorbidities: CVD: 13% CKD: NR COPD: NR DM: 11% HTN: 41% Obesity: NR Smoking: NR	COVID-19 severity: 55% mild, 45% critical/severe  ICU admission: None  Respiratory support: NR  Length of hospital stay: NR  Planned time post-hospital (days): 90-180  Reported time post-hospital (days, median): 93 (mild), 101 (critical severe)
Liotta, 2020 <sup>33</sup> USA  Retrospective  Funding: Not reported	Inclusion: Admitted with COVID-19; diagnosis confirmed by SARS-CoV-2 RT-PCR  Exclusion: None reported	N=509 Age (years, mean): 59 Gender (% male): 55 Race: 53% White, 30% Black or African American, 4% Asian, 13% Other/Unknown/Declined  Comorbidities: CVD: NR CKD: 11% COPD: NR DM: 30% HTN: 54% Obesity (BMI >30 kg/m <sup>2</sup> ): 52% Smoking: 28% Current	COVID-19 severity: 26% severe  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 27% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, median): 7  Planned/reported time post-hospital (days): 0 (discharge)
Liu, 2020 <sup>141</sup> China  Retrospective  Funding: Government	Inclusion: Discharged following hospitalization; infected with SARS-CoV-2 (testing details NR)  Exclusion: None reported	N=51 Age (years, mean): 47 Gender (% male): 42 Race/ethnicity: NR  Comorbidities: CVD: 2% CKD: NR COPD: NR DM: 8%	COVID-19 severity: All 'common' COVID-19 (fever, some respiratory-infection symptoms, pneumonia on radiographic images)  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		HTN: 14% Obesity: NR Smoking: 6%	Planned time post-hospital (days): 0 (last day before discharge), 14, and 28  Reported time post-hospital (days, median): 19 and 31
Loerinc, 2021 <sup>87</sup> USA  Retrospective  Funding: None	Inclusion: Confirmed SARS-CoV-2 by RT-PCR or ICD-10 code for COVID-19; discharged from hospital  Exclusion: Died during index hospital stay, admitted for unrelated reasons and incidentally tested (provider discretion) for COVID-19, discharged to home for end-of-life care with no additional post-discharge needs, transferred from study hospital to outside facility for continued hospitalization	N=310 Age (years, median): 58 Gender (% male): 49 Race/ethnicity: 69% African American, 18% White, 4% Hispanic, 9% Other  Comorbidities: CAD: 8% CKD: 19% COPD: 5% DM: 36% HTN: 65% Obesity (BMI ≥30): 45% Smoking: NR	COVID-19 severity: NR  ICU admission: 22%  Respiratory support Mechanical ventilation or ECMO: 14% NIV, HFNC, or CPAP: NR Other: 70%  Length of hospital stay (days, median): 5  Planned time post-hospital (days): 30  Reported time post-hospital: NR
Lovinsky-Desir, 2020 <sup>64</sup> USA  Retrospective  Funding: Government, Foundation	Inclusion: Sequential patients 65 years or younger; positive for severe SARS-CoV-2 (RT-PCR); hospitalized or died in the emergency department  Exclusion: None reported	N=1243 (age 21-29 [n=300] and 40-65 [n=943] groups only) Age (years, median): Age 21-39: 31-32 years Age 40-65: 56-58 years Gender (% male): 59 Race: 22% Black, 19% White, 1% Asian, 35% Other (NOTE: race identification declined for some patients)  Comorbidities: CVD: NR CKD: NR COPD: 0% (excluded from analysis) DM: NR HTN: NR Obesity: 42% Smoking: 59% Never, 4% Current, 11% Former	COVID-19 severity: 100% severe defined as hospitalization with confirmed positive SARS-CoV-2 PCR result or death in emergency department  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 21% NIV, HFNC, or CPAP: NR Other: 6% (tracheostomy)  Length of hospital stay (days, median): 4-6 days  Planned/reported time post-hospital: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		(NOTE: smoking status missing for some patients)	
Lv, 2020 <sup>88</sup> China  Retrospective  Funding: University and Government	Inclusion: Patients with laboratory (nucleic acid) confirmed infection with COVID-19 and lung involvement confirmed by chest imaging, improved and discharged  Exclusion: Pregnant	N=137 Age (years, mean): 47 Gender (% male): 52 Race/ethnicity: NR Comorbidities: CVD: NR CKD: NR COPD: 2% DM: NR HTN: NR Obesity: NR Smoking: 4%	COVID-19 severity: 20% (27/137) severe, 80% (110/137) non-severe  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 0% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay: NR  Planned time post-hospital (days): 14  Reported time post-hospital: NR
Mathew, 2020 <sup>46</sup> India  Retrospective  Funding: None	Inclusion: COVID-related stroke ( <i>ie</i> , stroke and positive RT-PCR for SARS-CoV-2), stroke confirmed by CT or MRI of the brain  Exclusion: None reported	N=62 Age (years, mean): 56 Gender (% male): 77 Race/ethnicity: NR  Comorbidities: CAD: 8% CKD: NR COPD: NR DM: 55% HTN: 62% Obesity: NR Smoking: NR	COVID-19 symptom onset to stroke symptom onset (days, mean): 12.5  COVID-19 severity: NR  ICU admission: NR  Respiratory support: NR  Length of hospital stay (days, mean): 16  Planned/reported time post-hospital (days): 0 (discharge)
Matsunaga, 2020 <sup>47</sup> Japan  COVIREGI-JP  Registry	Inclusion: Positive SARS-CoV-2 test, inpatient treatment  Exclusion: Participants in other clinical studies and inclusion in registry deemed inappropriate, refused to participate	N=2638 (number of cases for each parameter varied due to missing data) Age (years, median): 56 (non-severe=52, severe=57) Gender (% male): 59 (non-severe=55, severe=68) Race/ethnicity: 96% Japanese	COVID-19 severity: 68% non-severe; 32% severe  ICU admission: 11% (282/2638) (3% of non-severe cases, 26% of severe cases)  Respiratory support



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Funding: Government		Comorbidities: CVD: 1% CKD: NR COPD: 2% DM: 3% with complications, 14% without complications HTN: 15% Obesity: 6% Smoking: 14% Current	Mechanical ventilation or ECMO: 8% (2% of non-severe cases; 23% of severe) NIV, HFNC, or CPAP: 4% (2% of non-severe, 11% of severe) Other: NR  Symptom onset to hospitalization (days, median): 7  Length of hospital stay (days, median): 15  Planned/reported time post-hospital (days): 0 (discharge)
Mo, 2020 <sup>34</sup> China  Cross-sectional  Funding: Government	Inclusion: Hospital admitted; laboratory confirmed noncritical COVID-19  Exclusion: Critical cases	N=110 Age (years, mean): 49 Gender (% male): 50 Race: NR  Comorbidities: CVD: 3% Kidney disease: 2% Lung disease: 3% DM: 8% HTN: 24% Obesity: NR Smoking: 12%	COVID-19 severity: 22% mild, 61% pneumonia, 17% severe pneumonia  ICU admission: 0%  Respiratory support Mechanical ventilation or ECMO: 0% (critical cases excluded) NIV, HFNC, or CPAP Other: NR  Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)
Monday, 2020 <sup>89</sup> USA (Veterans)  Retrospective  Funding: None	Inclusion: Veterans with laboratory confirmed COVID-19 infection (RT-PCR)  Excluded: Civilians admitted to aid local health systems	N=79 Age (years, median): 69 Gender (% male): 94 Race/ethnicity: 90% African American, 9% White, 1% Other  Comorbidities: CAD: 35% CKD: 25% COPD: 38%	COVID-19 severity: NR  ICU admission: 11% (9/79) directly to ICU; 26% (18/70) were transferred to ICU  Respiratory support (highest required) Mechanical ventilation or ECMO: 30% NIV, HFNC, or CPAP: 3% Other: 51%



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		DM: 61% HTN: 92% Obesity: 53% Smoking: 66% (current or former)	Symptoms onset to admission (days, median): 7  Length of hospital stay (days, median): 6  Planned time post-hospital (days): 30  Reported time post-hospital: NR
Morin, 2021 <sup>120</sup> (COMEBAC Study Group) France  Prospective  Funding: Hospital/University	Inclusion: Adults (>18 years) admitted to a university hospital; survived 4 months after hospital discharge or after ICU discharge; had been hospitalized for >24 hours primarily because of COVID-19; diagnosis of SARS-CoV-2 (RT-PCR) or CT lung scan (or both)  Exclusion: Death within 4 months after discharge, persistent hospitalization, end-stage cancer, dementia, nosocomial COVID-19 infection, incidental positive SARS-CoV-2 results during hospitalization for a different medical indication	N=478 Age (years, mean): 61 Gender (% male): 58 Race/ethnicity: NR  Comorbidities: CVD: 16% CKD: 11% COPD: 4% DM: 27% HTN: 47% Obesity: 37% Smoking: No 76%, Former 18%, Current 6%	COVID-19 severity: NR  ICU admission: 30%  Respiratory support Mechanical ventilation or ECMO: 15% (51% of patients in ICU) NIV, HFNC, or CPAP: NR Other: 43  Length of hospital stay (median): 9 days  Planned time post-hospital (days): 120  Reported time post-hospital (days, median): 113 for telephone assessment; 125 for ambulatory assessment
Mowla, 2020 <sup>48</sup> Multinational  Retrospective  Funding: None	Inclusion: Adult hospitalized patients with definitive diagnosis of cerebral venous sinus thrombosis (CVST) and confirmed diagnosis of SARS-CoV-2 infection (RT-PCR) or typical symptoms and corresponding findings on chest CT if RT-PCR wasn't available (n=1)  Exclusion: None reported  NOTE: CVST was attributed to SARS-CoV-2 infection if infection symptoms or diagnosis (whichever came first) occurred	N=13 (COVID-19 group) Age (years, mean): 51 Gender (% male): 39 Race/ethnicity: NR  Comorbidities: NR	SARS-CoV-2 detected on same day as CVST presentation (n=2); COVID-19 associated symptoms prior to CVST symptom onset (n=11)  COVID-19 severity: 1 with no symptoms, 9 with mild to moderate symptoms, 1 with severe (no severity data for 2 patients)  ICU admission: NR  Respiratory support: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
	between 2 weeks prior to the onset of CVST symptoms and 2 days after hospital admission  NOTE: Included historical control group		Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)
Naar, 2020 <sup>49</sup> USA  Prospective  Funding: Not reported	Inclusion: Severe COVID-19 (RT-PCR confirmed) admitted to ICU  Exclusion: None reported	N=206 Age (years, median): 60 Gender (% male): 65 Race/ethnicity: 43% Hispanic or Latino, 28% White (Non-Hispanic), 11% Black or African American, 16% Other  Comorbidities: CHD: 9% CKD: 13% COPD: 8% DM: 43% HTN: 50% Obesity: NR Smoking: 23%	COVID-19 severity: NR  ICU admission: 100% (inclusion criteria)  Respiratory support Mechanical ventilation or ECMO: 87% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 19  Planned/reported time post-hospital (days): 0 (discharge)
Nachegea, 2020 <sup>50</sup> Democratic Republic of the Congo  Retrospective cohort  Funding: Not reported	Inclusion: All COVID-19 patients admitted; RT-PCR testing was done  Exclusion: None reported	N=766 (includes 34 children) Age (years, median): 46 Gender (% male): 66 Race/ethnicity: NR  Comorbidities: CAD: NR CKD: 1% Asthma/COPD: 3% DM: 14% HTN: 25% Obesity (self-report): 5% Smoking: NR	COVID-19 severity: 61% Mild, 14% Moderate, 21% Severe, 4% Critical (WHO criteria)  ICU admission: 25% (all severe and critical patients)  Respiratory support: NR  Length of hospital stay (days, median): 13 (for 645 recovered patients)  Planned/reported time post-hospital (days): 0 (discharge)
Nemer, 2021 <sup>51</sup> USA  Retrospective	Inclusion: Positive for SARS-CoV-2 (by RT-PCR), admitted from emergency department to non-ICU bed and subsequently discharged	N=350 Age (years, mean): 64 Gender (% male): 55 Race/ethnicity: NR	COVID-19 severity: NR  ICU admission: 14% (48/350)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Funding: None	Exclusion: None reported	Comorbidities: CAD: 16% CKD: 17% COPD: 8% DM: 32% HTN: 66% Obesity: NR Smoking: 28% (current or prior)	Respiratory support Mechanical ventilation or ECMO: 6% NIV, HFNC, or CPAP: 5% Other: 86%  Length of hospital stay (days, mean): 6  Planned/reported time post-hospital (days): 0 (discharge)
Nersesjan, 2021 <sup>142</sup> Denmark  Prospective  Funding: Foundation	Inclusion: Age ≥18, admitted to tertiary referral center from the hospital's COVID-19 ICU, COVID-19 intermediate care unit, and the neurological department; positive for SARS-CoV-2 (RT-PCR) except 1 patient with clinical suspicion (later confirmed)  Exclusion: None reported  NOTE: patients from ICU and intermediate care unit were screened for neurological and psychiatric symptoms during those admissions and evaluated for cognitive status and neurological deficits at discharge; patients from neurological department were screened for suspected neurological complications during/after a COVID-19 infection	N=61 Age (years, mean): 63 Gender (% male): 63 Race/ethnicity: 80% Caucasian, 0% Asian, 1% Middle Eastern, 0% African, 4% Inuit  Comorbidities: CAD: 13% CKD: NR COPD: 8% DM: 15% HTN: 15% Obesity: Smoking:	COVID-19 severity: NR  ICU admission: 57%  Respiratory support Mechanical ventilation or ECMO: 64% NIV, HFNC, or CPAP: 5% Other: 86%  Symptom onset to admission (days, mean): 7  Length of hospital stay (days, mean): 30  Planned time post-hospital (days): 90  Reported time post-hospital: NR
Ng, 2020 <sup>35</sup> USA  Retrospective  Funding: Not reported	Inclusion: All adult (age ≥18 years) patients who tested positive for COVID-19 (RT-PCR); hospitalized in 1 of 13 hospitals in a large health system  Exclusion: Transferred to hospitals outside the health system; admitted to inpatient obstetric service; end stage kidney disease; prior kidney transplant; <2 serum creatinine levels during admission	N=9,657 (demographic data for 40% [3,854/9,657] who developed AKI while hospitalized; 638 [17%] required KRT) Age (years, medians): KRT: 64 Non-KRT: 71 (P<.001) Gender (% male): KRT: 79 Non-KRT: 58 (P<.001) Race/Ethnicity:	COVID-19 severity: NR  ICU admission: KRT: 92%, Non-KRT: 45% (P<.001)  Respiratory support Mechanical ventilation or ECMO: KRT: 92%, Non-KRT 41% (P<.001) NIV, HFNC, or CPAP: NR Other: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		KRT: 25% Non-Hispanic White, 22% Non-Hispanic Black, 22% Hispanic Non-KRT: 38% Non-Hispanic White, 21% Non-Hispanic Black, 19% Hispanic (P<.001)  Comorbidities (all P<.001): CAD: KRT: 13%, Non-KRT: 18% CKD: KRT: 7%, Non-KRT: 9% COPD: KRT: 6%, Non-KRT: 8% DM: KRT: 48%, Non-KRT: 43% HTN: KRT: 64%, Non-KRT: 69% Obesity: BMI 30 or higher KRT: 45%, Non-KRT: 32% Smoking: KRT: 64% never, 22% current, 14% unknown Non-KRT: 67% never, 23% current, 10% unknown (P<.001)	Length of hospital stay (days, median): KRT: 29; Non-KRT: 12  Planned/reported time post-hospital (days): 0 days (discharge)
Ntaios, 2020 <sup>36</sup> Multi-national (Global COVID-19 Stroke Registry)  Retrospective  Funding: None	Inclusion: Hospitalized with laboratory-confirmed COVID-19 (96% by PCR, 4% by serology) and acute ischemic stroke  (NOTE: median delay between initiation of COVID-19 symptoms and stroke onset=7 days [IQR 2-15])  Exclusion: Infected after onset of stroke  (NOTE: also included propensity matched group of non-COVID-19 patients from other registries)	N=174 Age (years, median): 71 Gender (% male): 62 Race: NR  Comorbidities: CAD: 17% CVD: NR CKD: NR COPD: NR DM: 31% HTN: 68% Obesity: 37% Smoking: 28%	COVID-19 severity: NR  ICU admission: 23% (40/174)  Respiratory support Mechanical ventilation or ECMO: 16% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay: NR  Planned/reported time post-hospital (days): 0 (discharge)
Nugent, 2021 <sup>121</sup>  USA  Retrospective	Inclusion: Tested for COVID-19 by RT-PCR, developed AKI during hospitalization, survived past discharge, did not require dialysis within 3 days of discharge, had ≥1 measurement of serum creatinine as an outpatient post-discharge	N=1612 (182 COVID-19) Age (years, median): 70 (67 COVID-19 group) Gender (% male): 50 (53 COVID-19 group) Race/ethnicity: 40% Black, 41% White, 3% Asian, 17% Other; 22% Hispanic (COVID-19 group)	COVID-19 severity: NR  ICU admission: 37% (COVID-19 group)  Respiratory support Mechanical ventilation or ECMO: 29% (COVID-19 group)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Funding: Foundation	Exclusion: Age <18 years, determined to have ESKD, received prior kidney transplant, initial creatinine level ≥4 mg/dL	Comorbidities: CVD: NR CKD: 35% (33% COVID-19 group) COPD: 47% (45% COVID-19 group) DM: 52% (64% COVID-19 group) HTN: 89% Obesity: NR Smoking: NR	NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 14 (COVID-19 group)  Planned time post-hospital: NR  Reported time post-hospital (days, median): 92 (COVID-19 group)
Osikomaiya, 2021 <sup>122</sup>  Nigeria  Retrospective  Funding: Government	Inclusion: Discharged from COVID-19 facility based on resolution of symptoms and/or 2 negative RT-PCR SARS-CoV-2 results (at least 24 hrs apart)  Exclusion: Potential confounding comorbidities or concurrent infections	N=274 Age (years, mean): 42 Gender (% male): 66 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 3% HTN: 16% Obesity: NR Smoking: NR	COVID-19 severity (WHO): 7% asymptomatic, 51% mild, 39% moderate, 3% severe  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR  Planned time post-hospital (days): 14  Reported time post-hospital (days, median): 15
Overstad, 2020 <sup>52</sup> Norway  Retrospective  Funding: Not reported	Inclusion: Confirmed (RT-PCR) SARS-CoV-2, included in local quality register for COVID-19  Exclusion: None reported	N=70 Age (years, median): 59 Gender (% male): 67 Race/ethnicity: 60% Norwegian; 40% immigrants  Comorbidities: CAD: 30% CKD: 10% COPD/asthma: 16% DM: 24% HTN: NR Obesity: 31% Smoking: NR	Duration of symptoms prior to admission (days, median): 7  COVID-19 severity: 19% critically ill  ICU admission: 19%  Respiratory support Mechanical ventilation or ECMO: 19% NIV, HFNC, or CPAP: 6% Other: NR  Length of hospital stay (days, median): 6



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
			Planned/reported time post-hospital (days): 0 (discharge)
Ozer, 2021 <sup>123</sup> Turkey  Prospective  Funding: NR	Inclusion: Hospitalized with diagnosis of COVID-19 (positive PCR test); admitted to cardiology clinic for 'routine control'  Exclusion: Coronary artery disease, heart failure, atrial fibrillation, previous cerebrovascular disease, renal failure, severe COPD, malignancy, poor echogenicity, <18 years of age	N=74 Age (years, mean): 60 Gender (% male): 39 Race/ethnicity: NR  Comorbidities: CVD: NR (excluded) CKD: NR (excluded) COPD: NR (severe excluded) DM: 11% HTN: 43% Obesity: NR Smoking: 8%	COVID-19 severity: NR  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR  Planned time post-hospital: NR  Reported time post-hospital (days, mean): 30
Parra, 2020 <sup>90</sup> Spain  Case-control  Funding: None	Inclusion: Laboratory confirmed SARS-CoV-2 infection, admitted and discharged alive  Exclusion: Death during 3 weeks following discharge for controls  NOTE: Case patients were readmitted within 3 weeks of discharge with clinical presentation related to the infection or its treatment; controls were discharged but not readmitted	N=61 readmitted patients Age (years, median): 67 Gender (% male): 45 Race/ethnicity: NR  Comorbidities: CVD: 26% CKD: NR COPD: 20% DM: 23% HTN: 55% Obesity: 10% Smoking: NR	COVID-19 severity: NR  ICU admission: 5%  Respiratory support: NR  Length of hospital stay (days, median): 6  Planned time post-hospital (days): 21  Reported time post-hospital (days, median): 6
Patell, 2020 <sup>65</sup> USA  Retrospective  Funding: Not reported	Inclusion: Age ≥18 years; positive for SARS-CoV-2 (RT-PCR)  Exclusion: Hospitalized at time of analysis; discharged without any form of post-discharge contact in hospital medical records; discharged on therapeutic anticoagulation (separate reporting for	N=163 Age (years, median): 58 Gender (% male): 48 Race: 37% White  Comorbidities: Heart disease: 12% CKD: 10% Chronic respiratory disease: 22%	COVID-19 severity: NR  ICU admission: 26%  Respiratory support: NR  Length of hospital stay (days, median): 6  Planned time post-hospital (days): 30



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
	patients discharged on prophylactic-dose anticoagulation)	DM: 31% HTN: 53% Obesity: NR Smoking: NR	Reported time post-hospital: NR
Paterson, 2020 <sup>37</sup> United Kingdom  Retrospective case series  Funding: Several authors receive funding; not specified if related to this project	Inclusion: Patients referred to COVID-19 neurology/encephalitis and neurovascular multi-disciplinary team meetings; “definite” cases determined with RT-PCR  Exclusion: None reported	N=43 (demographic data for 16 with definite COVID-19 diagnosis and discharged) Age (years, mean): 57 Gender (% male): 56 Race: 63% White, 23% Black, 13% Asian  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 6% HTN: 38% Obesity: NR Smoking: NR	COVID-19 severity: 19% critical, 19% severe, 63% mild  ICU admission: 25%  Respiratory support Mechanical ventilation or ECMO: 19% (critical cases, by definition) NIV, HFNC, or CPAP: NR Other: 19% (severe cases, by definition)  Length of hospital stay (days, mean): 16.6 (reported for 12 patients)  Planned/reported time post-hospital (days): 0 (discharge)
Perry, 2020 <sup>53</sup> United Kingdom  Retrospective case-control  Funding: None	Inclusion: Clinical diagnosis of stroke and tested positive for SARS-CoV-2 within 4 days of admission (or within 4 days of stroke for inpatient strokes) or stroke in patients with clinical suspicion of COVID-19 at time of admission and were found to be SARS-CoV-2 positive at any point during first 10 days of admission  Exclusion: Subarachnoid hemorrhage, acquired COVID-19 after stroke, date of stroke onset could not be estimated NOTE: included concurrent control group of patients who were either consistently SARS-CoV-2 negative or never tested because no symptoms or signs of COVID-19	N=86 (COVID-19 group) Age (years, median): 75 Gender (% male): 55 Race/ethnicity: 72% White, 10% Black, 18% Asian, 1% Mixed/Other  Comorbidities: NR	For 44 cases with ischemic stroke and known dates of onset of stroke and COVID-19, onset of COVID-19 symptoms occurred a median of 6 days before stroke onset; for 3 patients with intracerebral hemorrhage and known dates, COVID-19 symptoms occurred a median of 4 days after stroke onset  COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 8% NIV, HFNC, or CPAP: 1% Other: 43%  Length of hospital stay (days, median): 7



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
			Planned/reported time post-hospital (days): 0 (discharge)
Puntmann, 2020 <sup>66</sup> Germany  Prospective  Funding: Government, Industry, Institution	Inclusion: Minimum of 2 weeks post-diagnosis of SARS-CoV-2 by RT-PCR; resolution of respiratory symptoms; negative results on swab test at end of isolation period  Exclusion: Recently recovered from COVID-19 and referred for clinical CMR imaging; unwilling to participate; absolute contraindications for contrast-enhanced magnetic resonance study  NOTE: included healthy and risk-factor matched control groups	N=100 Age (years, mean): 49 Gender (% male): 53 Race: NR  Comorbidities: CVD: 13% CKD: NR COPD: 21% DM: 18% HTN: 22% Obesity: NR Smoking: 22%	COVID-19 severity: 18% asymptomatic, 49% mild/moderate (both recovered at home), 33% severe (required hospitalization)  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 2%, 6% (hospitalized group) NIV, HFNC, or CPAP: 17%, 52% (hospitalized group) Other: 28% (NR for hospitalized group)  Length of hospital stay: NR  Planned/reported time post-hospital: NR  NOTE: median time from diagnosis to CMR was 71 [IQR 64-92] days)
Qin 2021 <sup>103</sup> China  Prospective  Funding: None	Inclusion: Patients hospitalized with PCR-confirmed COVID-19  Exclusion: Patients who died or were lost to follow-up.	N=647 (81 w pulmonary outcomes) Age (years, mean): 58 Gender (% male): 44 Race/ethnicity: NR  Comorbidities: CVD: 5% CKD: NR COPD: 6% (listed as chronic respiratory disease) DM: 11% HTN: 30% Obesity: NR Smoking: NR	COVID-19 severity: 51% non-severe, 49% severe  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 40% NIV, HFNC, or CPAP: 16% Other: NR  Length of hospital stay (days, mean): 18 (for 81 pulmonary patient subset)  Planned time post-hospital (days): 90  Reported time post-hospital: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Raman, 2021 <sup>124</sup> United Kingdom  Prospective  Funding: Government, Foundation	Inclusion: All patients with moderate to severe laboratory COVID-19 (positive SARS-CoV-2)  Exclusion: Severe comorbidities (end-stage renal, cardiac, liver, or neurological disease), contradictions to MRI  NOTE: included uninfected (negative for SARS-CoV-2 and asymptomatic) controls from the community (not hospitalized), group-matched for age, sex, body mass index, and risk factors	N=58 COVID-19 Age (years, mean): 55 Gender (% male): 59 Race/ethnicity: 22% Black/Asian and minority ethnic groups; 78% White  Comorbidities: CAD: 3% CKD: NR COPD: 5% DM: 16% (Type 1 and 2) HTN: 38% Obesity: NR Smoking: 35% Current or ex-smoker	COVID-19 severity: Moderate to severe (inclusion criteria)  ICU admission: 36% (21/58)  Respiratory support Mechanical ventilation or ECMO: 21% NIV, HFNC, CPAP: 26% Other: 46%  Length of hospital stay (days, median): 8.5  Planned time post-hospital (days): 30-180  Reported time post-hospital (days, median): 48
Ramani, 2021 <sup>91</sup> USA  Retrospective  Funding: Not reported	Inclusion: Admitted with COVID-19 (not confirmed), follow-up in post-COVID-19 ICU clinic approximately 6 weeks after discharge  Exclusion: None reported	N=28 (attended clinic) Age (years, median): 56 Gender (% male): 61 Race/ethnicity: 25% African American, 57% Hispanic, 14% White, 4% Asian Indian  Comorbidities: NR	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 86% NIV, HFNC, CPAP: NR Other: NR  Length of hospital stay (days, median): 22  Planned time post-hospital: NR  Reported time post-hospital (days, median): 40
Rashidi, 2020 <sup>92</sup> Iran  Retrospective  Funding: Not reported	Inclusion: Hospitalized patients with diagnosis of COVID-19 (RT-PCR [64%] or chest CT scan [36%])  Exclusion: None reported	N=1529 Age (years, median): 56 Gender (% male): 54 Race/ethnicity: NR  Comorbidities: CAD: 10%	COVID-19 severity: NR  ICU admission: 8%  Respiratory support: NR  Length of hospital stay: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		CKD: 5% COPD: 10% DM: 18% HTN: 29% Obesity: 14% Smoking: 15%	Planned time post-hospital (days): ≥45  Reported time post-hospital (days, range): 45-55
Rass, 2021 <sup>125</sup>  Austria  Prospective  Funding: NR	Inclusion: positive RT-PCR test, hospitalization or outpatient management, age ≥18y  Exclusion: age <18 years and patients who died during acute phase of COVID-19	N=135 (103 hospitalized) Age (years, median): 56 Gender (% male): 61 Race/ethnicity: NR  Comorbidities: CVD: 40% CKD: 7% COPD: NR DM: 18% HTN: 30% Obesity: NR Smoking: 3% (40% former)	COVID-19 severity (based on type of hospitalization): 23% severe (ICU), 53% moderate (non-ICU), 24% mild (outpatient)  ICU admission: 23%  Respiratory support Mechanical ventilation or ECMO: 22% NIV, HFNC, or CPAP: NR Other: 50%  Length of hospital stay (days, mean): 8  Planned time post-hospital (days): 90 Reported time post-hospital: NR
Remy-Jardin, 2021 <sup>126</sup>  France  Retrospective  Funding: None	Inclusion: patients hospitalized with confirmed COVID-19 who had residual respiratory symptoms ( <i>ie</i> , dyspnea, dry cough) and/or concern on chest radiographic abnormalities and were referred to the department of pulmonology for specialized follow-up. Underwent dual-energy CT angiography (DECTA)  Exclusion: Patients with no need for further investigation, received non-contrast CT or single-energy CTA	N=55 Age (years, mean): 60 Gender (% male): 76 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: 1% COPD: 5% DM: 16% HTN: 40% Obesity: 35% Smoking: 44%	COVID-19 severity: NR but 42% had “critical respiratory status” (admitted to ICU)  ICU admission: 42%  Respiratory support Mechanical ventilation or ECMO: 29% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay: NR  Planned time post-hospital (days): 90  Reported time post-hospital (days, median): 144 to CT exam



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Richardson, 2020 <sup>67</sup> USA  Case Series  Funding: Government	Inclusion: Consecutive patients at 12 hospitals in an academic health system requiring hospital admission with confirmed SARS-CoV-2 infection (RT-PCR)  Exclusion: None reported	N=5,700 (2,081 were discharged) Age (years, median): 63 Gender (% male): 60 Race: African American 23%, Asian 9%, White 40%, Multiracial 29%  Comorbidities: CAD: 11% CKD: 5% COPD: 5% DM: 34% HTN: 57% Obesity: 42% Smoking: 16%	COVID-19 severity: NR  ICU admission (discharged): 4% (82/2081)  Respiratory support Mechanical ventilation or ECMO (discharged): 2% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (medians, discharged patients): <18 years (n=32): 2.0 days 18-65 years (n=1,373): 3.8 days >65 years (n=676): 4.5 days  Planned time post-hospital: NR  Reported time post-hospital (days, median): 4.4
Roberts, 2020 <sup>68</sup> United Kingdom  Prospective  Funding: Not reported	Inclusion: Patients discharged following admission for COVID-19; 6-week follow-up for hospital-associated VTE (HA-VTE) events  Exclusion: None reported  NOTES: 1) patients received thromboprophylaxis while hospitalized 2) comparison cohort of post-discharge HA-VTE following medical admission in 2019	N=1877 Age (years, mean): NR Gender (% male): NR Race: NR  Comorbidities: NR	COVID-19 severity: NR  ICU admission: NR (11% [208/1877] admitted to critical care)  Respiratory support: NR  Length of hospital stay: NR  Planned/reported time post-hospital (days): 90
Rodriguez, 2020 <sup>54</sup> USA  American Heart Association	Inclusion: Age ≥18 years, hospitalized with COVID-19 as diagnosis, discharged with complete data on admission and discharge dates, age, sex, and medical history	N=7,868 Age (years, mean): 61 Gender (% male): 55 Race/ethnicity: 33% Hispanic, 26% non-Hispanic Black, 6% Asian, 35% non-Hispanic White	Time from symptom onset to hospital arrival (days, weighted mean): 5  COVID-19 severity: NR  ICU admission: 29%



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
COVID-19 CVD Registry  Funding: Industry, Foundation	Exclusion: None reported	Comorbidities: CAD: 9% CKD on dialysis: 4% Pulmonary disease: 19% DM: 37% HTN: 60% Obesity: 39% Smoking: 6%	Respiratory support Mechanical ventilation or ECMO: 22% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, weighted mean): 6  Planned/reported time post-hospital (days): 0 (discharge)
Sachdeva, 2020 <sup>69</sup> USA  Retrospective  Funding: None	Inclusion: Age ≥18 years; end stage kidney disease on chronic peritoneal dialysis; hospitalized with COVID-19 (positive by PCR testing)	N=11 Age (years, median): 54 (<50: 36%; 50-59: 27%; 60-69: 27%; 70-79: 9%) Gender (% male): 27 Race: 9% Hispanic, 45% Non-Hispanic Black; 9% Non-Hispanic White; 36% Other or Unknown  Comorbidities: CAD: 9% CKD: 100% COPD: 0% DM: 45% HTN: 91% Obesity (BMI ≥30 kg/m <sup>2</sup> ): 36% Smoking: 82% Never, 18% Former	COVID-19 severity: NR  ICU admission: 27%  Respiratory support Mechanical ventilation or ECMO: 27% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, mean): 9 (range 2-23)  Planned/reported time post-hospital: NR
Salisbury, 2020 <sup>93</sup> United Kingdom  Retrospective  Funding: Government	Inclusion: Age ≥18 years, SARS-CoV-2 (RT-PCR)  Exclusion: Admission <24 hours, VTE diagnosed within first 24 hours after presentation, diagnosed with COVID-19 during hospital stay for other medical conditions	N=152 (subgroup discharged without an indication for therapeutic anticoagulation) Age (years, median): 62 Gender (% male): NR Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR  ICU or high dependency unit admission: 16%  Respiratory support: NR for subgroup  Length of hospital stay (days, mean): 7  Planned time post-hospital (days): 90  Reported time post-hospital (days): 42



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Sami, 2020 <sup>94</sup> Iran  Prospective  Funding: Government, University	Inclusion: Admitted for COVID-19 (RT-PCR or chest CT), discharged  Exclusion: Minors	N=490 Age (years, mean): 57 Gender (% male): 61% Race/ethnicity: NR  Comorbidities: HTN: 35% CVD/CAD: 14% DM: 28% CKD: 3% COPD: 2% Smoking: 14%	COVID-19 severity: 12% of patients discharged had composite events of mechanical ventilation or ICU admission  ICU admission: 8% (overall)  Respiratory support: NR  Length of hospital stay (days, median): 5  Planned time post-hospital (days): 7, 30, 84, and 365  Reported time post-hospital (days): 7 and 28
Shah, 2020 <sup>95</sup> Canada  Prospective  Funding: Foundation, University	Inclusion: Adults hospitalized with laboratory-confirmed SARS-CoV-2 infection	N=60 Age (years, median): 67 Gender (% male): 68 Race/ethnicity: NR  Comorbidities: HTN: 35% DM: 22% CPOD: 13% CAD: 10% CKD: 7%	COVID-19 severity: NR  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 20% NIV, HFNC, or CPAP: NR Other: 78%  Length of hospital stay (days, mean): 10  Planned time <u>post-symptom onset</u> (days): 90  Reported time <u>post-symptom onset</u> : NR
Sibilia, 2021 <sup>144</sup> Spain  Retrospective  Funding: None	Inclusion: All patients discharged after hospitalization with COVID-19  Exclusion: NR	N=172 Age (years, mean): 56 Gender (% male): 57 Race/ethnicity: NR  Comorbidities: CVD: 14% CKD: NR COPD: 4% DM: 15%	COVID-19 severity: 29% moderate, 71% severe  ICU admission: 43%  Respiratory support Mechanical ventilation or ECMO: 18% NIV, HFNC, or CPAP: NR Other: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		HTN: 33% Obesity: NR Smoking: 5%	Length of hospital stay (days, mean): 20  Planned time post-hospital (days): 90  Reported time post-hospital (mean, days): 101
Somani, 2020 <sup>70</sup> USA  Retrospective  Funding: Government	Inclusion: Age ≥18 years; laboratory confirmed SARS-CoV-2; admitted and subsequently discharged alive from 5 health system hospitals  Excluded: Discharge before April 12, 2020 (all patients had ≥14 day observation for possible readmission); returned <12 hours after discharge; died during index admission	N=2,864 (n=103 returned to hospital; 2,761 did not) Age (years, median): 66 Gender (% male): 58 Race: 4% Asian, 28% Black, 27% Hispanic, 24% White; 17% Unknown/Other NOTE: no differences between groups for Age, Gender, or Race  Comorbidities: CAD: 8% CKD: 5% COPD: Returned: 7%; No Return: 3%; P=.035 DM: 15% HTN: Returned: 35%; No Return: 22%; P=.003 Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: Returned: 6%; No Return: 19%; P=.001  Respiratory support Mechanical ventilation or ECMO: Returned: 1%; No Return: 11%; P=.003 NIV, HFNC, or CPAP: Returned 42%; No Return: 49% Other: NR  Length of hospital stay (days, median): Returned: 4.7; No Return: 7; P=.006  Planned time post-hospital (days): <14  Reported time post-hospital: NR
Sonnweber 2020 <sup>96</sup> Austria  Prospective  Funding: Foundation, Industry, University	Inclusion: Diagnosis of COVID-19 (RT-PCR and typical presentation), hospitalized or outpatient care with persisting symptoms  Exclusion: Unable to attend regular follow-up, rejection of study participation	N=145 (n=133 for second follow-up) Age (years, mean): 57 Gender (% male): 55 Race/ethnicity: NR  Comorbidities: CVD: 40% HTN: 30% COPD: 6% DM: 17% CKD: 7% Obesity: NR Smoking: 3%	COVID-19 severity: 75% hospitalized  ICU admission: 22%  Respiratory support Mechanical ventilation or ECMO: 27% NIV, HFNC, or CPAP: 3% Other: 66%  Length of hospital stay: NR  Planned time <u>post-diagnosis</u> (days): 60 and 100



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
			Reported time <u>post-diagnosis</u> (days, mean): 63 and 103
Spinicci, 2021 <sup>127</sup>  Italy  Retrospective  Funding: University	Inclusion: Patients hospitalized with COVID-19  Exclusion: Discharged >10 weeks, unable to attend clinical visit due to hospitalization or residents in care facilities, patient refusal	N=100 Age (years, median): 68 Gender (% male): 59 Race/ethnicity: NR  Comorbidities: CVD: 12% CKD: 7% COPD: 12% DM: 21% HTN: 50% Obesity: 25% Smoking: NR	COVID-19 severity (WHO): 9% mild, 32% moderate, 12% severe, 47% critical  ICU admission: 31%  Respiratory support Mechanical ventilation or ECMO: 21% NIV, HFNC, or CPAP: 33% Other: 33%  Length of hospital stay (days, median): 16  Planned time post-hospital (days): 56  Reported time post-hospital (days, median): 60
Stevens, 2020 <sup>97</sup>  USA  Retrospective  Funding: Not reported	Inclusion: Positive for SARS-CoV-2 (RT-PCR), admitted to an ICU, received RRT for AKI  Exclusion: ESRD on KRT prior to admission	N=115 Age (years, median): 63 Gender (% male): 73 Race/ethnicity: 32% Black, 23% White, 4% Asian, 23% Multi-racial, 17% NR  Comorbidities: CAD: 10% CKD: 28% COPD: 7% DM: 50% HTN: 70% Obesity: 54% Smoking: 8%	Time from symptom onset to presentation (days, median): 6  COVID-19 severity: NR  ICU admission: 100% (inclusion criteria)  Respiratory support Mechanical ventilation or ECMO: 99% NIV, HFNC, or CPAP: NR Other: NR  Length of hospital stay (days, median): 36  Planned time post-hospital: NR  Reported time post-hospital (days, median): 29 (from RRT initiation)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
Suarez-Robles, 2021 <sup>128</sup> France  Retrospective  Funding: None	Inclusion: Patients discharged for PCR-confirmed COVID-19  Exclusion: subjects who did not consent	N=134 Age (years, mean): 59 Gender (% male): 46 Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR  ICU admission: 1%  Respiratory support Mechanical ventilation or ECMO: NR NIV, HFNC, or CPAP: NR Other: 3%  Length of hospital stay: NR  Planned time post-hospital (days): 90  Reported time post-hospital: NR
Suleyman, 2020 <sup>138</sup> USA  Retrospective case series  Funding: Not reported	Inclusion: Consecutive adult patients evaluated at 5 hospitals and 9 emergency departments in a health system; confirmed SARS-CoV-2 infection (RT-PCR)  Exclusion: Lack of demographic and baseline data	N=355 (hospitalized) (108 discharged home after initial evaluation not reported here) Age (years, mean): 61 Gender (% male): 47 Race: 73% African American  Comorbidities: CAD: 16% CKD: 45% COPD: 12% DM: 43% HTN: 73% Obesity: 59% Smoking: 39%	COVID-19 severity: NR  ICU admission: 40%  Respiratory support Mechanical ventilation or ECMO: General practice unit: 0% (0/234); ICU: 81% (114/141) P<.001 NIV, HFNC or CPAP: NR Other: NR  Length of hospital stay (days, median): General practice unit: 5 [3-7] ICU: 15 [9-23] P<.001  Planned time post-hospital (days): 30  Reported time post-hospital: NR
Taquet, 2021 <sup>129</sup> USA  Retrospective	Inclusion: Age >10 years, confirmed diagnosis of COVID-19  Exclusion: NR	N=236379 (COVID-19 group) Age (years, mean): 46 Gender (% male): 44	COVID-19 severity: NR  ICU admission: 4% (8945/236379)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
Funding: University	NOTE: Participants organized into one primary cohort (confirmed diagnosis of COVID-19) and 2 matched control cohorts (diagnosed with influenza and diagnosed with any respiratory tract infection including influenza)	Race/ethnicity: 57% White, 19% Black or African American, 3%Asian, 20% Unknown; 16% Hispanic or Latino  Comorbidities: CVD: NR CKD: 7% COPD: NR DM: 16% HTN: 30% Obesity: 18% Smoking: NR	Respiratory support: NR  Length of hospital stay: NR  Planned time post-hospital (days): 180  Reported time post-hospital: NR
Tomasoni, 2021 <sup>130</sup> Italy  Cross-sectional  Funding: Foundation	Inclusion: Patients with documented clinical recovery and virological clearance after hospitalization for COVID-19 disease (clinical recovery was defined as absence of fever for 48-72 hours and normal oxygen saturation on ambient air with concomitant hospital discharge and virological clearance was defined as presence of 2 consecutive negative nasopharyngeal swabs taken 24-48 hours apart, at least 14 days after clinical recovery)  Exclusion: NR	N=105 Age (years, median): 55 Gender (% male): 73% Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR (all patients had interstitial pneumonia at admission)  ICU admission: NR  Respiratory support Mechanical ventilation or ECMO: 28% (non-invasive or orotracheal intubation) NIV, HFNC, or CPAP: NR Other: 72% (low flow oxygen or none)  Length of hospital stay (days, mean): 8  Planned time post-hospital (days): 30-90  Reported time post-hospital (days, median): 46
Tudoran, 2021 <sup>131</sup> Romania  Prospective  Funding: None	Inclusion: Hospitalized for SARS-CoV-2 (RT-PCR) and discharged alive; mild/moderate disease; age <55 years; no history of significant associated cardiovascular pathology or diabetes; cardiologic exam with routine transthoracic echocardiography (TTE) at/near time of admission	N=125 Age (years, median): 47 Gender (% male): 50 Race/ethnicity: NR  Comorbidities: CVD: NR CKD: NR COPD: NR	COVID-19 severity: 100% mild/moderate  ICU admission: 0% (exclusion criteria)  Respiratory support Mechanical ventilation or ECMO: 0% (exclusion criteria) NIV, HFNC, or CPAP: NR Other: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
	Exclusion: Age >55; severe or critical. COVID-19; respiratory insufficiency requiring mechanical ventilation and/or ICU stay during hospitalization; associated medical conditions on TTE as determined by cardiologist	DM: 0% (inclusion criteria) HTN: NR Obesity: 29% Smoking: NR	Length of hospital stay: NR Planned time post-hospital (days): 42-70 Reported time post-hospital: NR
Venturelli, 2021 <sup>132</sup> Italy  Prospective  Funding: Public	Inclusion: Patients aged >18 discharged from the emergency department or admitted wards of the hospital with any condition possibly related to SARS-CoV-2  Exclusion: Asymptomatic pregnant women admitted for delivery and asymptomatic patients found positive to the molecular test admitted for planned procedure for other conditions	N=767 (NOTE: 510 felt recovered) Age (years, mean): 63 Gender (% male): 67 Race/ethnicity: NR  Comorbidities: CVD: 10% CKD: NR COPD: 5% DM: 7% HTN: 22% Obesity: 22% Smoking: 4%	COVID-19 severity: NR  ICU admission: 9%  Respiratory support Mechanical ventilation: 8% NIV, HFNC, or CPAP: 18% Other: 73  Length of hospital stay (days, median): 18 (without ICU admission), 58 (with ICU admission)  Planned time post-hospital: NR  Reported time post-hospital (days, median): 81
Vizcaychipi, 2020 <sup>38</sup> United Kingdom  Prospective  Funding: None	Inclusion: Admitted to Emergency Department; completed hospital encounter (discharged alive or died)  Exclusion: Remained in admitting hospital; transferred to another hospital  NOTE: study was designed to evaluate the effect of an electronic medical record alert system on early mortality related to COVID-19	N=1,039 admitted; data for n=939 who completed hospital encounter Age (years, median): 67 Gender (% male): 60 Race: 62% White  Comorbidities: CVD: NR CKD: NR COPD: 10% DM: 38% HTN: 53% Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: 14.4% (150/1039)  Respiratory support: NR  Length of hospital stay (days, median): 7  Planned/reported time post-hospital (days): 0 (discharge)



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
Vlachou, 2021 <sup>98</sup> United Kingdom  Retrospective  Funding: Not reported	Inclusion: Admitted with positive test for SARS-CoV-2, underwent CT pulmonary angiography (CTPA) because of increasing oxygen requirements or refractive hypoxia, not improving on oxygen, very elevated D-dimer, or tachycardia disproportionate to clinical condition  Exclusion: None reported	N=39 Age (years, mean): 62 Gender (% male): 56 Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR (100% 'severe' – not defined)  ICU admission: NR  Respiratory support: NR  Length of hospital stay (days, median): 7  Planned time post-hospital (days): 28  Reported time post-hospital: NR
Wang, 2020 <sup>71</sup> China  Prospective cohort  Funding: Government	Inclusion: Confirmed COVID-19 patients discharged from hospital  Exclusion: Could not be contacted or refused to participate	N=131 Age (years, median): Non-severe: 38; Severe: 60 (P<.0001) Gender (% male): 45 Race: NR  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 2% HTN: 3% Obesity: NR Smoking: NR	COVID-19 severity: 53% (69/131) severe  ICU admission: NR (NOTE: "severe" category did not require ICU admission)  Respiratory support: NR  Length of hospital stay (days, median): 15  Planned time post-hospital (days): 28  Reported time post-hospital (days, median): 25
Wu, 2021 <sup>133</sup> China  Prospective  Funding: Government	Inclusion: ≥18 years, severe COVID-19 (confirmed with RT-PCR), discharged from hospital  Exclusion: History of hypertension, diabetes, cardiovascular disease, cancer, or chronic lung disease (including asthma and COPD) or history of smoking documented at time of hospital admission; required intubation and mechanical ventilation	N=83 Age (years, mean): 60 Gender (% male): 57 Race/ethnicity: NR  Comorbidities: see Exclusion CVD: 0% CKD: NR COPD: 0% DM: 0% HTN: 0% Obesity: NR	COVID-19 severity: Severe (inclusion criteria)  ICU admission: 0% (exclusion criteria)  Respiratory support Mechanical ventilation or ECMO: 0% (exclusion criteria) NIV, HFNC, or CPAP: 55% Other: 45%  Length of hospital stay (days, median): 29



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		Smoking: 100% never smoked	Planned time post-hospital (days): 90, 180, 270, and 365  Reported time post-hospital (days, median): 98, 189, 275, and 348
Xia, 2020 <sup>55</sup> China  Retrospective  Funding: Not reported	Inclusion: Confirmed COVID-19, hospitalized in regular care inpatient ward	N=282 (mild and moderate COVID-19) Age (years, mean): 48 Gender (% male): 50 Race/ethnicity: NR  Comorbidities: CVD: 22% CKD: NR COPD: 2% DM: 11% HTN: NR Obesity: NR Smoking: NR	Time from symptom onset to hospitalization (days, mean): 5  COVID-19 severity: 2% mild, 98% moderate  ICU admission: 0% (inclusion criteria)  Respiratory support: NR  Length of hospital stay (days, mean): 16  Planned/reported time post-hospital (days): 0 (discharge)
Xiong, 2021 <sup>99</sup> China  Prospective  Funding: Not reported	Inclusion: Ages 20-80 years, diagnosed with COVID-19, cured and discharged  Exclusion: Severe and complex underlying diseases, receiving invasive treatment, women who were pregnant or breastfeeding  NOTE: included control group free of COVID-19, similar demographics, completely quarantined at home for >3 months with little physical work	N=538 (those who completed telephone follow-up from group of 891 discharged) Age (years, median): 52 Gender (% male): 46 Race/ethnicity: NR  Comorbidities: CHD: 3% CKD: 2% COPD: 4% DM: 7% HTN: 15% Obesity: NR Smoking: NR	COVID-19 severity: 5% critical, 34% severe, 62% "general"  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR  Planned time post-hospital: NR  Reported time post-hospital (days, median): 97



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
Xu, 2020 <sup>72</sup> China  Retrospective case series  Funding: Government	Inclusion: Adults with confirmed SARS-CoV-2 infections (RT-PCR); critically ill (admitted to ICU, requiring mechanical ventilation or fraction of inspired oxygen concentration ≥60%)  Exclusion: Deceased within 48 hours after ICU admission	N=92 (survivors; data from 147 non-survivors not reported here) Age (years, mean): 58 Gender (% male): 58 Race: NR  Comorbidities: CVD: 15% CKD: NR COPD: 3% DM: 20% HTN: 45% Obesity: NR Smoking: NR	COVID-19 severity: 100% critically ill  ICU admission: 100%  Respiratory support Mechanical ventilation or ECMO: 9% NIV, HFNC, or CPAP: 44% Other: INR  Length of hospital stay: NR  Planned time post-hospital (days): 60  Reported time post-hospital: NR
Yasin, 2021 <sup>134</sup> Egypt  Retrospective  Funding: None	Inclusion: Patients hospitalized with confirmed COVID-19 by nasopharyngeal swab RT-PCR testing who underwent initial CT during hospitalization and follow-up CT after discharge  Exclusion: NR	N=210 Age (years, mean): 54 Gender (% male): 71 Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: NR  ICU admission: 25%  Respiratory support: NR  Length of hospital stay (days, mean): 16  Planned time post-hospital: NR  Reported time post-hospital (days, mean): 42
You, 2020 <sup>73</sup> China  Case series  Funding: None	Inclusion: Laboratory-confirmed COVID-19; pulmonary function test after discharge from hospital  Exclusion: None reported	N=18 Age (years, mean): 51 Gender (% male): 56 Race: NR  Comorbidities: CVD: NR CKD: NR COPD: NR DM: 6% HTN: 17% Obesity: NR	COVID-19 severity: 67% (12/18) moderate, 28% (5/18) severe, 6% (1/18) critical  ICU admission: NR  Respiratory support: NR  Length of hospital stay (days, mean): 28  Planned time post-hospital: NR



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics Time of Post-hospital Follow-up
		Smoking: NR	Reported time post-hospital (days, mean): 40 (to pulmonary function test)
Yu, 2020 <sup>74</sup> China  Retrospective case series  Funding: Government	Inclusion: COVID-19 positive confirmed by pharyngeal swab nucleic acid testing; hospitalized then discharged after treatment; underwent thin-section chest CT scans at least twice while hospitalized and at least once following discharge  Exclusion: None reported	N=32 (n=14 fibrosis group, n=18; non-fibrosis group) Age (years, medians): fibrosis group 54; non-fibrosis group 37; P=.008 Gender (% male): 69 Race: NR  Comorbidities: CVD: 6% CKD: NR COPD: 3% DM: 6% HTN: 13% Obesity: NR Smoking: NR	COVID-19 severity: NR  ICU admission: fibrosis group 35.7% (5/14); non-fibrosis group 0% (0/18); P=.01  Respiratory support: NR  Length of hospital stay (days, median): fibrosis group 19.5 [11.5-21.8]; non-fibrosis group 10.0 [6.0-15.3]; P=.001  Planned time post-hospital: NR  Reported time post-hospital (days, median): fibrosis group 9 [7-11]; non-fibrosis group 9 [7.8-11.3]
Zhang, 2020 <sup>100</sup> China  Retrospective  Funding: Government	Inclusion: Convalescent patients (recovered from COVID-19)  Exclusion: None reported	N=112 (adult group) Age (years, median): 37 Gender (% male): 56 Race/ethnicity: NR  Comorbidities: NR	COVID-19 severity: 17% severe, 83% non-severe  ICU admission: NR  Respiratory support: NR  Length of hospital stay: NR  Planned time post-hospital (days): 14  Reported time post-hospital: NR
Zhao, 2020 <sup>101</sup> China  Retrospective  Funding: Government	Inclusion: Adults with confirmed SARS-CoV-2 (RT-PCR)  Exclusion: Critical COVID-19	N=55 Age (years, mean): 48 Gender (% male): 58 Race/ethnicity: NR  Comorbidities: CVD: 4%	COVID-19 severity: 7% mild, 86% moderate, 7% severe  ICU admission: NR  Respiratory support: Mechanical ventilation or ECMO: 0%



Author, Year Country Study Design Funding	Inclusion/Exclusion Criteria	Baseline Demographic Data	Hospitalization Characteristics  Time of Post-hospital Follow-up
		CKD: NR COPD: NR DM: 4% HTN: 11% Obesity: NR Smoking: NR	NIV, HFNC, or CPAP: NR Oxygen: 26%  Length of hospital stay (days, mean): 15  Planned time post-hospital (days): 90  Reported time post-hospital: NR
Zhou, 2021 <sup>135</sup> China  Prospective  Funding: None	Inclusion: PCR-confirmed COVID-19  Exclusion: Age <18 years; classified as severe community acquired pneumonia; required invasive or non-invasive ventilatory support; admitted to ICU; history of heart failure	N=97 Age (years, mean): 47 Gender (% male): 54 Race/ethnicity: NR  Comorbidities: CAD: 6% CKD: 1% COPD: NR DM: 11% HTN: 25% Obesity: NR Smoking: NR	COVID-19 severity: 100% non-severe  ICU admission: 0% (exclusion criteria)  Respiratory support Mechanical ventilation or ECMO: 0% (exclusion criteria) NIV, HFNC, or CPAP: 0% (exclusion criteria) Other: NR  Length of hospital stay (days, median): 17  Planned time post-hospital: NR  Reported time post-hospital (days, median): 11

*Abbreviations:* AKI=acute kidney injury; ARDS=acute respiratory distress syndrome; ARF=acute respiratory failure; CAD=coronary artery disease; CMR=cardiovascular magnetic resonance; COVID-19, SARS-CoV-2: 2019 novel coronavirus; CPAP=continuous positive airway pressure; CT=computed tomography; CVD=cardiovascular disease; CKD=chronic kidney disease; COPD=chronic obstructive pulmonary disease; DM=diabetes mellitus; ECMO=extracorporeal membrane oxygenation; ELSO=Extracorporeal Life Support Organization; ESKD=end stage kidney disease; FIM=Functional Independence Measurement; hs-CTnT=high-sensitivity cardiac troponin T; HTN=hypertension; ICD=International Classification of Disease; IQR=interquartile range; ICU=intensive care unit; KRT=kidney replacement therapy; MRI=magnetic resonance imaging; PE=pulmonary embolism; POCUS=point-of-care ultrasound; RT-PCR: reverse transcriptase polymerase chain reaction; SpO2=Peripheral Capillary Oxygen Saturation; VTE=venous thromboembolism; WHO=World Health Organization



**TABLE 2. STUDY QUALITY APPRAISAL**

Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Akhtar, 2021 <sup>39</sup> Qatar Retrospective	Evaluated all acute stroke admissions	Single site; n=833 (32 COVID-19 positive)	Unclear how modified Rankin Scale was administered	Yes	Little information on COVID-19 severity
Al-Aly, 2021 <sup>104</sup> USA (Veterans) Retrospective	Hospitalized cohort within a specific time range	Multisite; n=13,654 (COVID-19 group)		ICD-10 codes	Little information on COVID-19 severity; no information on comorbid conditions but focus was on new onset diagnoses
Al Kasab, 2020 <sup>23</sup> USA, South America, Europe Prospective	Consecutive patients meeting eligibility criteria	Multisite; n=13	Unclear how modified Rankin Scale was administered	Yes	No information about comorbidities; little information about hospitalization
Alemanno, 2021 <sup>105</sup> Italy Prospective	Clear inclusion; consecutive patients for baseline but not follow-up	Single site; n=87 (56 at 1mo follow-up)	Unclear how outcome measures were administered	Established measures (MoCA, MMSE), but criteria for “deficit severity” unclear	Little information on COVID-19 severity Little patient demographic data No information to explain poor follow-up retention
Alharthy, 2021 <sup>24</sup> Saudi Arabia Prospective	Unclear if consecutive patients	Single site; n=89	Unclear how ultrasound was analyzed	Unclear – used point-of-care ultrasound (noted limited evidence supporting its diagnostic utility in COVID-19)	Unclear how subjects were identified for enrollment; limited patient information
Alharthy, 2020 <sup>75</sup> Saudi Arabia Prospective	Consecutive patients reviewed for eligibility	Single site: n=171	Unclear how ultrasound was analyzed - authors note that adequate training is required for reliable use	Unclear – authors note ‘scarce data’ on use of ultrasound in COVID-19	Little information on COVID-19 severity

Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Anand, 2020 <sup>40</sup> USA Retrospective	All patients during study period reviewed for eligibility	Single site; n=74	Unclear who reviewed medical records and whether data were verified	Medical records	Little information on COVID-19 severity
Arab-Zozani, 2020 <sup>76</sup> Iran Cross-sectional	'Systematic sampling' to select patients	Single site; n=409	Telephone interviews at 22 +/- 15 days post- discharge	Medical records and telephone interview for EQ-5D-5L	Little information on comorbidities or COVID-19 severity
Atalla, 2020 <sup>136</sup> USA Retrospective	Unclear if all patients were reviewed for eligibility	2 sites; n=339 (19 readmitted)	Only captured patients who presented to hospital they were discharged from	Medical records and post- discharge call to patients	Little information on COVID-19 severity
Ayoubkhani, 2021 <sup>106</sup> United Kingdom Retrospective	Included all patients with exclusions clearly noted in flow diagram	Multisite (NHS hospitals in England); n=47,780 (COVID-19 group)		ICD-10 codes	Little information on COVID-19 severity
Barbaro, 2020 <sup>41</sup> Multi Retrospective cohort	ECMO registry cases – unclear if all patients included in registry	International registry; n=1035	Outcome unclear for 10% (discharged to long-term acute care or unspecified location)	Registry data	Little information on COVID-19 severity; outcomes reported for ARDS cohort
Bellan, 2021 <sup>107</sup> Italy Prospective	Contacted all participants discharged	Single site; n=238	Used validated scales and measures	Yes	Little information on COVID-19 severity
Benussi, 2020 <sup>25</sup> Italy Retrospective cohort	Included all patients meeting eligibility criteria	Single site ("hub" for acute cerebrovascular diseases); n=56 with COVID-19	Unclear how Stroke Scale was administered	Unclear how stroke was diagnosed; NIH Stroke Scale for assessment	No information on COVID-19 severity
Bhatt, 2021 <sup>42</sup> USA Retrospective	Unclear if all eligible patients (authors note limitations of coding)	Multicenter database (over 1000 healthcare entities/health systems); n=8,393 COVID-19	Unclear whether coding was consistent across sites	Database records (ICD-10 codes)	Little information on COVID-19 severity



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Boari, 2021 <sup>108</sup> Italy Prospective	Contacted all surviving participants	Single site; n=94	Used validated scales and measures	Yes	Little information on COVID-19 severity and follow-up group
Bowles, 2020 <sup>77</sup> USA Retrospective cohort	All patients admitted to home health care following hospitalization	Single home health agency serving 64 hospitals in 1 city; n=1,409	Outcomes assessed by nurse or physical therapist; mandatory assessment	Outcome and Assessment Information Set (OASIS)	No information on COVID-19 severity; results reported by age groups
Brendish, 2020 <sup>78</sup> United Kingdom Prospective cohort	Convenience sample – all patients from clinical trial of point-of-care testing for SARS-CoV-2; concurrent comparison group COVID- 19 negative	Single site; n=1,054	Unclear how data were extracted from medical records (qualifications/verification)	Medical records	COVID-19 severity (respiratory support) not linked to outcomes
Brosnahan, 2020 <sup>79</sup> USA Retrospective	All patients discharged and re-presenting reviewed for eligibility	Single site; n=9 (limited to patients re- presenting to initial hospitalization site)	Broad definition of thrombotic event	Hospital records filtered for presumed events	No information on COVID-19 severity
Casas-Rojo, 2020 <sup>56</sup> Spain Retrospective cohort	Consecutive patients	150 hospitals in registry; n=15,111	Unclear how patients were followed for readmission data; unclear if patients may have presented to other hospitals	Electronic data capture system with procedures for verification of data	Little information on COVID-19 severity
Chan, 2021 <sup>102</sup> USA Retrospective	Unclear if consecutive patients	5 sites; n=3993 (1835 with AKI while hospitalized; 212 with post-discharge follow- up)	Unclear – AKI defined based on change from baseline creatinine; 63% had missing data and creatinine was imputed; used creatinine from 7 to 365 days prior to admission for others	Yes – dataset	No patient characteristics for the n=212 patients with post-discharge follow- up data



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Chevinsky 2021 <sup>109</sup> USA Retrospective	Hospitalized cohort within a specific time range	Multisite (922 hospitals), data from large US health plan; n=27,284 (COVID-19 group)		ICD-10 codes	Little information on COVID-19 severity; no information on comorbid conditions but focus was on new onset diagnoses
Chopra, 2020 <sup>80</sup> USA Retrospective	Sites may use pseudo- randomized procedure to select patients for inclusion if unable to abstract all hospitalizations	38 sites; n=1,250	Trained abstractors and surveyors; structured templates; 42% completed for 60 day post-discharge survey	Medical records and telephone survey	Little information on COVID-19 severity
Collins, 2020 <sup>26</sup> USA Retrospective	All patients reviewed for eligibility	3 sites; n=20	Yes	Electronic medical records	Little information on COVID-19 severity
Curci, 2020 <sup>27</sup> Italy Cross-sectional	Consecutive patients reviewed for eligibility; all eligible included	Single site; n=32	Yes	No spirometry measures	Yes; all patients were admitted to ICU
Daher, 2020 <sup>81</sup> Germany Prospective	Consecutive	Single site; n=33	Trained study team administered questionnaires; echocardiography	Follow-up at pulmonary disease outpatient clinic	Labeled as 'severe' but no information on criteria
Daugherty 2021 <sup>110</sup> USA Retrospective	Hospitalized cohort within a specific time range	Multisite, data from large US health plan; n=18,118 (COVID-19)		ICD-10 codes	Little information on COVID-19 severity; no demographic data for hospitalized subgroup
Dawson, 2020 <sup>143</sup> United Kingdom Prospectively collected/ retrospectively analyzed	All patients meeting eligibility criteria	Single site; n=208	Length of follow-up not reported	Unclear how readmission or re-presentation was assessed; unclear if patients may have gone to other hospitals	Little patient information; time post- hospital not reported

Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
DeBolt, 2020 <sup>43</sup> USA Retrospective	Unclear if all pregnant women were reviewed for study eligibility	4 sites; n=38	Unclear how data were abstracted	Electronic medical records	Yes; results also reported separately for 'critical' group
de Graaf, 2021 <sup>111</sup> the Netherlands Prospective	All patients in geographic region were eligible	Single site; n=87	Cardiac evaluation by cardiologist; no information about pulmonary function testing	Yes (cardiac and pulmonary function tests)	Accounted for all patients admitted; little information on COVID-19 severity
de Havenon, 2021 <sup>146</sup> USA Retrospective	Unclear if all cases during study period included; COVID-19 via ICD code U07.1	312 US Hospitals; n=2,086	Reported "favorable discharge" combining home and acute rehabilitation	Healthcare analytics database	Little information on COVID-19 severity
De Lorenzo, 2020 <sup>82</sup> Italy Prospective	Consecutive	Single site; n=185	Multi-disciplinary team	Medical records and outpatient clinical follow-up; mMRC Dyspnea scale, MoCA (cognitive)	Yes; some severity measures included in prediction analyses
De Michieli, 2021 <sup>112</sup> USA Retrospective	Consecutive patients; unclear if all were admitted	Multi-site; n=367	Method for obtaining follow-up data not reported	Unclear	Little information on COVID-19 severity; unclear why 7% of survivors not assessed at follow-up
Dennis, 2021 <sup>57</sup> United Kingdom Prospective	Unclear if consecutive patients; described population as "low-risk" (eg, younger, largely without risk factors, pre-existing disease, or hospitalization) but inclusion criteria don't require that	2 sites; n=37 hospitalized (of n=201 included)	Unclear whether MRI was dual-reviewed; unclear how questionnaires were administered	Validated questionnaires (patient-reported), blood tests, MRI	Unclear how subjects were identified for enrollment; little information on COVID-19 severity; time post-hospital unclear
Doher, 2020 <sup>145</sup> Brazil Retrospective cohort	Unclear if all patients were assessed for eligibility	Single site; n=201	Unclear	Medical records or telephone contact	Yes (all ICU patients); unclear if some patients remain hospitalized at end of study period



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Egol, 2020 <sup>58</sup> USA Prospective	Consecutive patients	7 hospitals served by a university orthopedic department; n=17	Unclear how patients were followed after discharge	Unclear – source of data not reported	Unclear how many were COVID-19 positive at admission vs during admission
EI Moheb, 2020 <sup>59</sup> USA Retrospective	All patients meeting eligibility criteria identified; report focused on 141 of 242 (58%) with COVID-19 whose gastrointestinal complications while hospitalized were previously reported; after propensity matching, 92 COVID-19 patients were included	Single site with 13 ICUs; n=92	Unclear – length of follow- up for emergency department readmission not reported	Unclear – source of data not reported	No information about patient disposition at discharge, post- discharge monitoring, or length of follow-up
Engelen, 2021 <sup>113</sup> Belgium Prospective	Evaluated all confirmed COVID-19 admissions 75 and younger	Single site; n=146	Yes	Yes	Little information on COVID-19 severity beyond ICU admission
Eswaran, 2021 <sup>114</sup> USA Retrospective	Yes	11 sites; n=447	Yes	Yes	No comorbidity information; Little information on COVID-19 severity
Fisher, 2020 <sup>28</sup> USA Retrospective	Unclear whether age 18 was included or excluded; not specified if all patients were reviewed for eligibility	3 hospitals in a healthcare system; n=3,345	Yes	Medical records	Disposition of all patients is unclear
Frija-Masson, 2020 <sup>60</sup> France Retrospective	Unclear if all patients were reviewed for eligibility	Single site; n=50	Yes	Did not perform CT measures at 30 days	Unclear how 30 days post-symptom onset relates to time post- discharge
Fuglebjerg, 2020 <sup>29</sup> Denmark Case series	Consecutive patients reviewed for eligibility; all eligible included	Single site; n=26	Yes	Authors note that clinical implications of hypoxia are not well described in the literature	Little information about comorbidities or COVID-19 severity



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Garrigues, 2020 <sup>61</sup> France Prospective, survey	Contacted all eligible patients	Single site; n=120	Telephone questionnaire administered by trained physicians	Some elements of questionnaire were developed by the study authors	Limited information about comorbidities
Goicoechea, 2020 <sup>30</sup> Spain Retrospective	All admitted patients on maintenance hemodialysis meeting eligibility criteria	Single site; n=36	No information	Unclear (“worsening or appearance of X-ray pulmonary infiltrates”)	No information about patients who were discharged
Grewal, 2020 <sup>31</sup> USA Retrospective	All patients admitted meeting eligibility criteria	Single site; n=13	Yes	Yes	Little information about COVID-18 severity
Gupta, 2021 <sup>139</sup> USA Cohort	Consecutive	67 hospitals; n=3,099	Data form completed by study personnel; automated and manual data verification protocols	Chart review with standardized data form	Yes (all ICU admissions)
Hall, 2021 <sup>115</sup> United Kingdom Retrospective	Clear inclusion, exclusion seems convenience based, only evaluated first 200 patients; patients excluded for frailty may have post- discharge outcomes	Single site; n=200	yes	yes	Little information on COVID-19 severity
Hamilton, 2020 <sup>83</sup> United Kingdom Retrospective	All patients during study period meeting eligibility criteria	4 hospitals; n=1,032	Data manually checked for duplicate values	Electronic medical records	Time post-hospital unclear
Han, 2021 <sup>116</sup> China Prospective	Unclear how sample was identified	Single site; n=114	Used CT scans and imaging	Yes	Yes
Hegde, 2020 <sup>44</sup> USA Retrospective case series	All patients during study period meeting eligibility criteria	4 hospitals; n=7	Unclear who did abstraction and if data were verified	Electronic medical records	Yes



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Hill, 2020 <sup>84</sup> USA Retrospective	All patients assessed for eligibility	14 hospitals; n=86	Multiple strategies to query electronic medical records; criteria for confirmation of thrombosis	Electronic medical records; authors note limitations of medical record search	No information on comorbidities; little information on COVID-19 severity
Hittesdorf, 2020 <sup>140</sup> USA Retrospective	Unclear if all patients assessed for eligibility	Single site; n=116	Unclear how patients were followed	Yes	Limited information on comorbidities; 100% ICU admission
Hu, 2020 <sup>45</sup> China Cross-sectional	Unclear if all patients assessed for eligibility	Single site; n=76	Unclear if artificial intelligence findings were reviewed by clinicians	Unclear – artificial intelligence program for detection of fibrosis	Yes
Huang C, 2021 <sup>85</sup> China Retro and Prospective	All patients evaluated for eligibility; many exclusion criteria; stratified disproportional random sampling (according to severity) used to select patients for pulmonary function and imaging	Single site; n=1,733	Follow-up at outpatient clinic with trained physicians; CT mages evaluated by clinicians; interpretation of ultrasound unclear	mMRC dyspnea, EQ-5D-5L, pulmonary function, ultrasound, CT	Yes; outcomes reported by level of severity (oxygen requirement)
Huang L, 2020 <sup>62</sup> China Retrospective	Consecutive patients meeting eligibility criteria	Single site; n=26	Yes	Yes	Little information on COVID-19 severity; no information on time post-discharge
Huang Y, 2020 <sup>63</sup> China Retrospective	Unclear if all patients were reviewed for eligibility	Single site; n=57	19% (13/70) eligible could not be contacted or declined participation	Yes	Little information on COVID-19 severity
Jacobs, 2020 <sup>117</sup> USA Prospective	Contacted all surviving participants, 52% response rate	Single site; n=183	Used validated scales and measures	Yes	Yes



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Karaarslan, 2021 <sup>118</sup> Turkey Prospective	Included 300 participants, threshold for analysis	Single site; n=300	Used validated scales and measures, phone interviews	Yes	No information on respiratory outcomes at baseline, little information on COVID-19 severity
Katz, 2020 <sup>32</sup> USA Retrospective	All patients meeting eligibility criteria	11 hospitals in a health system; n=86	Yes	Yes – chart review and other databases	Little information on comorbidities
Khalili, 2020 <sup>86</sup> Iran Prospective cohort	Unclear if all patients were reviewed for eligibility	Single site; n=254	Unclear how patients were followed	Number of patients discharged alive not reported so readmission only available for total number enrolled	Little information on COVID-19 severity; time post-hospital unclear
Knights, 2020 <sup>137</sup> United Kingdom Retrospective	All admitted patients	Single site; n=69 (survivors)	Unclear how post-discharge care needs were captured	Data from electronic and paper medical records; additional information from patients	No information about patients discharged to care home or other; time post-hospital unclear
Li, 2021 <sup>119</sup> China Prospective	Reached 44% of eligible participants with 18% inclusion rate	Single site; n=107	Used clinical and laboratory tests	Yes	Little information on baseline characteristics
Liotta, 2020 <sup>33</sup> USA Retrospective	Consecutive patients	10 hospitals in a health system; n=509	Modified Rankin Scale scores determined independently by 2 reviewers	Data from electronic medical records (including templates specific to COVID-19), clinical notes, diagnostic studies, and physician-documented diagnoses; modified Rankin Scale	Yes
Liu, 2020 <sup>141</sup> China Retrospective	Unclear if all patients were reviewed for eligibility	2 sites; n=51	CT interpreted by 2 experienced radiologists (consensus)	CT	Yes (note: patients had similar level of severity)



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Loerinc, 2021 <sup>87</sup> USA Retrospective	All patients assessed for eligibility	Single site; n=310	Data manually collected from notes; training or experience not reported	Unclear – data from physician discharge notes, discharge medication reconciliation, case management and nursing notes at discharge, and post- hospital transition care management notes	Little information on COVID-19 severity
Lovinsky-Desir, 2020 <sup>64</sup> USA Retrospective	Sequential patients	2 hospitals in network; n=1243 (n=95 in <21 age group not included - median age 14-15)	Unclear how patients were followed for readmission data	Medical records	Length of follow-up for readmission unclear
Lv, 2020 <sup>88</sup> China Retrospective	Consecutive	Single site; n=137	Automated pulmonary function testing system	Electronic medical records	Little information on comorbidities
Mathew, 2020 <sup>46</sup> India Retrospective	Consecutive cases	13 hospitals in 1 city; n=62	Unclear how modified Rankin Scale was administered	Unclear	No information on COVID-19 severity
Matsunaga, 2020 <sup>47</sup> Japan Registry	Patients entered into registry at discretion of principal investigator	227 sites; n=2638	'Research collaborators' input data to registry; funding for each patient enrolled	Unclear how self-care ability was defined; source of oxygen therapy required and RRT data not reported	Yes
Mo, 2020 <sup>34</sup> China Retrospective	Unclear if all patients were reviewed for eligibility	Unclear if single site; n=110	Yes	Lung function tests but no imaging	Yes
Monday, 2020 <sup>89</sup> USA (Veterans) Retrospective	All admitted patients	Single site; n=79	Unclear who abstracted data	Electronic medical records	Time post-hospital unclear

Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Morin, 2021 <sup>120</sup> (COMEBAC Study Group) France Prospective	Unclear if reviewed records of all patients hospitalized during study period for eligibility	Single site; n=478 for telephone assessment; n=177 for clinic assessment (invited based on prior ICU admission or current symptoms during telephone assessment)	Telephone assessment by “medical officer”; independent review of lung CT scans; neuropsychologist for cognitive assessment	Predominantly standard instruments	57% of eligible patients consented to participate in follow-up; reported demographics for those who did not consent were similar
Mowla, 2020 <sup>48</sup> Multinational Retrospective	Unclear if all patients were reviewed for eligibility	9 sites in 3 countries; n=13 (other sites invited but had no cases or could not meet timeline)	Unclear how modified Rankin Scale was administered; missing data for 33% (3/9 discharged)	Dual confirmation of cerebral venous sinus thrombosis; medical records	No information on comorbidities; limited information on COVID- 19 severity
Naar, 2020 <sup>49</sup> USA Prospective	Consecutive patients	Single site; n=206	Unclear	Unclear how RRT status was identified	Little information on COVID-19 severity
Nachegea, 2020 <sup>50</sup> Democratic Republic of the Congo Retrospective cohort	All admitted patients meeting eligibility criteria	7 sites; n=766	Unclear who abstracted/verified data	Database records	Yes
Nemer, 2021 <sup>51</sup> USA Retrospective	All admitted patients meeting eligibility criteria	2 sites; n=350	Unclear who abstracted/verified data	Electronic medical records	Yes
Nersesjan, 2021 <sup>142</sup> Denmark Prospective	Consecutive cases	Single site; n=61	Unclear who abstracted/verified data	MoCA and clock drawing tests; electronic medical records	Yes
Ng, 2020 <sup>35</sup> USA Retrospective	All admitted patients meeting eligibility criteria	13 hospitals in a health system; n=3,854 (2,771 survivors)	Yes	Data from chart reviews (hospital progress, discharge, and social worker notes)	Yes

Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Ntaios, 2020 <sup>36</sup> Multi-national Retrospective	Consecutive patients meeting eligibility criteria	28 sites in 16 countries; n=174 (96 survivors)	Unclear how modified Rankin Scale was administered	Global COVID-19 Stroke registry	Yes
Nugent, 2021 <sup>121</sup> USA Retrospective	Clear inclusion; consecutive patients	5 hospitals; n=1612 (182 COVID-19 positive)	eGFR calculated using the Chronic Kidney Disease Epidemiology equation	AKI classified into 3 stages according to creatine criteria; cut-offs unclear	No information on COVID-19 severity
Osikomaiya, 2021 <sup>122</sup> Nigeria Retrospective	Clear but broad inclusion, unclear exclusion; patients referred to discharge center	Patients referred from 6 isolation facilities; n=274	Limitation of self-reported symptoms	Limitation of self-reported symptoms	Yes; inclusion of COVID-19 severity and comorbidities
Overstad, 2020 <sup>52</sup> Norway Retrospective	All admitted patients meeting eligibility criteria	Single site; n=70	Unclear who abstracted/verified data	COVID-19 registry, medical records, electronic charts	Yes
Ozer, 2021 <sup>123</sup> Turkey Prospective	Consecutive patients reviewed for eligibility	Single site; n=74	Images reviewed by single cardiologist blinded to study data	Echocardiography	No information on COVID-19 severity
Parra, 2020 <sup>90</sup> Spain Case-control	Unclear if all patients assessed for eligibility	Single site; n=61	Unclear – authors note that some patients may have been readmitted at other hospitals	Unclear	Limited information on COVID-19 severity
Patell, 2020 <sup>65</sup> USA Retrospective	Consecutive patients meeting eligibility criteria	Single site; n=163	Unclear if all patients were followed	Medical records	Yes
Paterson, 2020 <sup>37</sup> United Kingdom Retrospective	Cases referred to COVID team meetings (“selective”)	Single site; n=43 (29 with definite COVID- 19)	Unclear	Unclear – little information about source of data	Yes
Perry, 2020 <sup>53</sup> United Kingdom Retrospective case-control	Unclear if all patients were reviewed for eligibility	13 sites; n=86	Unclear how modified Rankin Scale was administered	Case report forms from multiple study sites	Limited information on COVID-19 severity

Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Puntmann, 2020 <sup>66</sup> Germany Prospective	Unselected cohort	Single site; n=100	Yes	Yes	Little information about subgroups of hospitalized patients vs home recovery patients; time post-hospital unclear
Qin 2021 <sup>103</sup> China Prospective	Clear, minimal	Single site; n=81	Yes	Yes	Yes
Raman, 2021 <sup>124</sup> United Kingdom Prospective	“Unselected cohort”; patients approached by medical team for possible inclusion in study; also noted that all patients assessed for eligibility	Unclear if single site; n=58 COVID-19 and n=30 group-matched controls	Images assessed by multiple specialists independently; questionnaires completed with electronic data capture; training/experience of assessors unclear	Software for MRI image analysis; standard questionnaires for cognitive function and dyspnea; electronic medical records for clinical data	All moderate to severe COVID-19; controls were not hospitalized; no demographic information about the 36 patients excluded
Ramani, 2021 <sup>91</sup> USA Retrospective	Appears to include all patients admitted during the study period	Single site; n=28	Patients attended Post-COVID-19 ICU Clinic; little information on how specific outcomes were assessed	MoCA, Quality of Life in Neurological Disorders, pulmonary function	No information on comorbidities; little information on COVID-19 severity
Rashidi, 2020 <sup>92</sup> Iran Retrospective	Consecutive cases	3 sites; n=1,529	Interview by trained physician; clinic follow-up if symptomatic	Imaging or ventilation-perfusion scans	Little information on COVID-19 severity
Rass, 2021 <sup>125</sup> Austria Prospective	Clear inclusion; consecutive patients	Multisite; n=135 (103 hospitalized)	Neurological evaluation performed by neurological consultants or junior neurologists under supervision of consultants; limitation of self-reported cognitive symptoms	Established measures (MoCA, SS-16) with clear criteria, but no baseline comparator; limitation of self-reported cognitive symptoms	Yes; inclusion of COVID-19 severity and comorbidities



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Remy-Jardin, 2021 <sup>126</sup> France Retrospective	Yes	Unclear if single site; n=55	Yes	Yes	Yes
Richardson, 2020 <sup>67</sup> USA Case series	All admitted patients meeting eligibility criteria	12 sites; n=5,700	Authors note median follow-up of 4.4 days post- discharge	Yes – electronic medical records	Little information on patients who were discharged
Roberts, 2020 <sup>68</sup> United Kingdom Prospective	All events at designated sites (patients may have presented elsewhere during follow-up)	2 sites of 1 hospital; n=1877	No routine contact during follow-up (only captured patients who presented to hospital they were discharged from)	Yes – imaging required	No patient demographic data; little information on COVID-19 severity
Rodriguez, 2020 <sup>54</sup> USA Registry	Consecutive patients in American Heart Association COVID-19 Registry meeting eligibility	88 sites; n=7,868	Abstraction/verification unclear	Medical records	Yes
Sachdeva, 2020 <sup>69</sup> USA Retrospective	All patients meeting eligibility criteria	13 hospitals of a health system; n=11	Unclear – methods for obtaining follow-up information not specified; unclear if patients may have presented to other hospitals	Yes – electronic and manual chart review	Time post-hospital not reported
Salisbury, 2020 <sup>93</sup> United Kingdom Retrospective	Consecutive cases	Single site; n=303	Authors note links with other local hospitals if patients present there with VTE	Electronic medical records; radiographic confirmation	No information on comorbidities; little information on COVID- 19 severity
Sami, 2020 <sup>94</sup> Iran Prospective	Unclear – “first COVID-19 cases with complete information”	Single site; n=490	Data cross-checked by research team (professionals and clinical faculty)	Telephone follow-up	Yes
Shah, 2020 <sup>95</sup> Canada Prospective	Unclear if all assessed for eligibility – “referrals”	Single site; n=60	Unclear – methods not reported	Pulmonary function tests; dyspnea symptoms present/absent	Limited information on COVID-19 severity; time post-hospital unclear



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Sibilia, 2021 <sup>144</sup> Spain Retrospective	Unclear	Single site; n=172	Yes	Yes	Yes
Somani, 2020 <sup>70</sup> USA Retrospective	Unclear if all patients were reviewed for eligibility	5 hospitals of a health system; n=103	No systematic follow-up; unclear if patients may have presented to other hospitals	Yes – electronic health records	Yes
Sonnweber 2020 <sup>96</sup> Austria Prospective	Unclear if all patients were reviewed for eligibility	3 sites; n=145 (74% were hospitalized)	In-person follow-up visits; training of individuals interpreting the test findings is unclear	mMRC dyspnea score, pulmonary function, CT, echocardiography	No information about subset of patients hospitalized; time post-hospital unclear
Spinicci, 2021 <sup>127</sup> Italy Retrospective	Clear but broad inclusion; consecutive patients, all patients invited for follow-up	Single site; n=100	Limitation of self-reported symptoms	Limitation of self-reported symptoms	Yes; inclusion of COVID-19 severity and comorbidities
Stevens, 2020 <sup>97</sup> USA Retrospective	All patients reviewed for eligibility	Single site; n=115	Data abstraction/verification unclear	Electronic medical record	Yes (100% ICU admissions)
Suarez-Robles, 2021 <sup>128</sup> France Retrospective	Yes	Single site; n=134	Patient self-report	Moderate, self-report	Sparse baseline data (no co-morbidity data); Little information on COVID-19 severity
Suleyman, 2020 <sup>138</sup> USA Retrospective case series	Consecutive patients	5 sites; n=355 hospitalized patients	Follow-up to 30 days post-discharge	Yes – electronic medical records	Yes
Taquet, 2021 <sup>129</sup> USA Retrospective	Evaluated COVID-19 patients using network of electronic records	Multisite; n=236,379	Evaluated outcomes using ICD codes and medical records	Yes	Limited information on COVID-19 severity
Tomasoni, 2021 <sup>130</sup> Italy Cross-sectional	Unclear how participants were identified	Single site; n=105	Used validated scales and measures	Yes	Little information on baseline characteristics



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Tudoran, 2021 <sup>131</sup> Romania Prospective	All patients hospitalized during study period reviewed for eligibility	Single site; n=125; 182 of 254 eligible patients agreed to participate (57 subsequently ineligible)	“Experienced operator” performed all echocardiographic measurements	Echocardiography	Baseline TTE did not include careful measures of ventricular function so may not have ruled out pre-existing cardiovascular conditions; little demographic data
Venturelli, 2021 <sup>132</sup> Italy Prospective	Contacted all participants meeting eligibility	Single site; n=767	Used validated scales and measures	Yes	Yes
Vizcaychipi, 2020 <sup>38</sup> United Kingdom Prospective	Consecutive admissions evaluated for eligibility	2 hospitals of 1 institution; n=939	Source of disposition data not reported	Unclear	Limited information on comorbidities and COVID-19 severity
Vlachou, 2021 <sup>98</sup> United Kingdom Retrospective	Only patients who underwent CT pulmonary angiography during hospitalization for increasing O2 requirements, refractory hypoxia, elevated D-dimer or tachycardia	Single site; n=39	No information about interpretation or verification of findings	Yes	No information on comorbidities and COVID-19 severity
Wang, 2020 <sup>71</sup> China Prospective cohort	Unclear if all patients were reviewed for eligibility	Single site; n=131	Followed every 7 days up to 4 weeks; methods for follow-up data collection unclear	Data obtained by questionnaire	Limited information on comorbidities
Wu, 2021 <sup>133</sup> China Prospective	Screened all patients admitted during study period for eligibility	Single site; n=83	Images independently assessed by experienced radiologists; little information about administration of other measures	mMRC for dyspnea; imaging per guidelines	Little information about patients who could not be contacted, declined participation, or withdrew; “severe” COVID but excluded patients receiving mechanical ventilation



Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Xia, 2020 <sup>55</sup> China Retrospective	Unclear if all patients were reviewed for eligibility	Single site; n=282	CT scans evaluated by radiologists with experience in thoracic imaging evaluation	CT	Yes
Xiong, 2021 <sup>99</sup> China Prospective	Patients who were discharged and completed telephone contact (many exclusions); comparison cohort was not hospitalized	Single site; n=538	Post-discharge contact was by experienced clinicians	Self-report of newly diagnosed hypertension	Yes
Xu, 2020 <sup>72</sup> China Retrospective case series	Unclear if all patients were reviewed for eligibility	3 sites; n=239 ICU patients	Unclear if follow-up was complete for post-discharge patients	Self-report	Time post-hospital unclear
Yasin, 2021 <sup>134</sup> Egypt Retrospective	Clear but minimal	Unclear if single site; n=210	Yes	Yes	Sparse baseline data (no co-morbidity data); little information on COVID-19 severity
You, 2020 <sup>73</sup> China Case series	Unclear if study included all or consecutive patients	Single site; n=18	Yes	CT scans not completed at the same time as pulmonary function tests; CT scans reviewed independently by 2 cardiothoracic radiologists blinded to clinical information	Limited information on comorbidities and COVID-19 severity
Yu, 2020 <sup>74</sup> China Retrospective case series	Unclear if study included all patients	Single site; n=32	Specific criteria provided for many of the outcomes assessed from CT scan; scans reviewed independently by 3 experienced radiologists	CT	Yes
Zhang, 2020 <sup>100</sup> China Retrospective	All patient recovered from COVID-19	Single site; n=134	Imaging findings evaluated by experienced radiologists	CT	No information on comorbidities; limited information on COVID-19 severity

Author, Year Country Study Design	Clear Inclusion Criteria/Consecutive/ Complete?	Adequate Sample Size?	Condition/Outcome Measured in a Standard Reliable Way?	Valid Methods for Identification of the Condition/Outcome?	Adequate Information about Subjects & Setting?
Zhao, 2020 <sup>101</sup> China Retrospective	Patients identified through national influenza surveillance system	3 sites; n=55	Images reviewed by 2 radiologists blinded to clinical data; followed ATS-ERS guidelines for pulmonary testing	CT, pulmonary function	Limited information on comorbidities and COVID-19 severity
Zhou, 2021 <sup>135</sup> China Prospective	Referrals to Infectious Diseases clinic were invited to participate (also described as consecutive)	Single site; n=97 (only those who attended clinic and completed ECG or echocardiography)	Unclear – no information on interpretation of test results	MRI, ECG and echocardiography, CMRI; not all patients completed all follow-up tests (depended on screening test results)	Limited demographic data

Abbreviations: AKI=Acute kidney injury; CT=computed tomography; ECG=electrocardiogram; eGFR=estimated globular filtration rate; mMRC=Modified Medical Research Council Dyspnea Scale; MMSE=Mini Mental State Examination; MoCa=Montreal Cognitive Assessment; MRI=magnetic resonance imaging; TTE=transthoracic echocardiogram

Reference: JBI Critical Appraisal Checklist for Case Series

**TABLE 3. PULMONARY OUTCOMES**

Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
Al-Aly, 2021 <sup>104</sup> USA (Veterans) Retrospective  Outcomes vs individuals hospitalized with seasonal influenza	NR	NR	NR	<b>Shortness of breath<sup>a</sup></b> HR (adjusted) 1.14 (95%CI 1.04, 1.26) Excess burden per 1000 hospitalized COVID-19 patients at 6 months 13.22 (95%CI 3.68, 21.94)  COVID-19 group: N=13,654 Control group (influenza): N=13,997
Alharthy, 2021 <sup>24</sup> Saudi Arabia Prospective	NR	NR	NR	<b>Pleural Effusions at Discharge (Ultrasound)</b> 1.5% (1/64) (n=64 survivors)
Alharthy, 2020 <sup>75</sup> Saudi Arabia Prospective	NR	NR	NR	<b>Pleural Effusions (Ultrasound)</b> <b>2 months</b> 18.9% (24/127) <b>4 months</b> 11.8% (15/127) P=.0001
Ayoubkhani, 2021 <sup>106</sup> United Kingdom Retrospective  Controls from general population not meeting inclusion criteria for COVID-19; unclear if any were hospitalized at time of study	NR	NR	NR	<b>Respiratory Disease, new onset events</b> COVID-19 Group: 21.5% (6085/28,335) Rate per 1000 person-years 538.9 (95%CI 525.5, 552.6) General Population Control Group: 0.8% (240/28,335) Rate per 1000 person-years 19.7 (95%CI 17.3, 22.4) P<.001 between groups  <b>Respiratory Disease, all events</b> COVID-19 Group: 29.6% (14,140/47,780) Rate per 1000 person-years

Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
				770.5 (95%CI 757.8, 783.3) General Population Control Group: 5.4% (2585/47,780) Rate per 1000 person-years 129.1 (95%CI 124.2, 134.2) P<.001 between groups  NOTE: study reported increased respiratory disease among those admitted to ICU (data NR)
Bellan, 2021 <sup>107</sup> Italy Prospective	NR	NR	<b>D<sub>LCO</sub> &lt;80%</b> 52% (113/219)	<b>Dyspnea</b> 6% (13/238)
Boari, 2021 <sup>108</sup> Italy Prospective	25% (24/94)	NR	<b>D<sub>LCO</sub> &lt;80%</b> 32% (30/94)	<b>Dyspnea</b> 36% (33/91)
Chevinsky 2021 <sup>109</sup> USA Retrospective  Controls were hospitalized individuals who did not meet inclusion criteria for COVID-19 and were not diagnosed with COVID-19 during the 4 months after index encounter	NR	NR	NR	<b>Respiratory failure; insufficiency; arrest</b> vs hospitalized non-COVID-19 control group 1-30 days after discharge OR (adjusted) 3.3 (95%CI 2.6, 4.1) 31-60 days after discharge OR (adjusted) 1.0 (95%CI 0.70, 1.4) 60-90 days after discharge OR (adjusted) 0.93 (95%CI 0.65, 1.3) 90-120 days after discharge OR (adjusted) 0.73 (95%CI 0.47, 1.1)  <b>Pneumonia (except that caused by tuberculosis)</b> vs hospitalized non-COVID-19 control group 1-30 days after discharge OR (adjusted) 5.5 (95%CI 4.1, 7.5) 31-60 days after discharge



Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
				OR (adjusted) 1.3 (95%CI 0.89, 2.0) 60-90 days after discharge OR (adjusted) 0.88 (95%CI 0.53, 1.5) 90-120 days after discharge OR (adjusted) 1.0 (95%CI 0.58, 1.9)  N=27,284 for COVID and control groups
Curci, 2020 <sup>27</sup> Italy Cross-sectional	NR	NR	NR	<p><b>PaO<sub>2</sub>/FiO<sub>2</sub> (mmHg)</b>                      Mild alteration (300-399): 22% (7/32)                      Moderate alteration (200-299): 38% (12/32)                      Severe alteration (&lt;200): 41% (13/32)</p> <p><b>Respiratory Supports Needed</b>                      None: 13% (4/32)                      Nasal cannula: 41% (13/32)                      Oxygen mask: 13% (4/32)                      Venturi mask: 25% (8/32)                      Non-rebreather mask: 9% (3/32)</p> <p><b>mMRC Dyspnea Scale</b>                      Grade 4: 13% (4/32)                      Grade 5: 88% (28/32)</p>
Daher, 2020 <sup>81</sup> Germany Prospective	NR	NR	NR	<p><b>Dyspnea (symptom questionnaire)</b>                      Admission: 48% (16/33)                      6 week follow-up: 33% (11/33)</p>
Daugherty 2021 <sup>110</sup> USA Retrospective  Controls did not have a COVID-19 diagnosis and were not admitted to a	NR	NR	NR	<p><b>New Clinical Diagnoses at 4 months</b></p> <p><b>Respiratory failure (acute respiratory failure, chronic respiratory failure, interstitial lung disease)</b>                      COVID group: 2.60%                      Non-COVID Control group: 0.19%                      Risk difference 2.41% (95%CI 1.35, 3.2)</p>



Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
hospital for COVID-19				<p><b>Acute respiratory failure</b>                      COVID group: 2.58%                      Non-COVID Control group: 0.18%                      Risk difference 2.4%                      (95%CI 1.67, 3.43)</p> <p><b>Chronic respiratory failure</b>                      COVID group: 1.53%                      Non-COVID Control group: 0.05%                      Risk difference 1.48%                      (95%CI 0.97, 1.75)</p> <p><b>Interstitial lung disease</b>                      COVID group: 1.60%                      Non-COVID Control group: 0.13%                      Risk difference 1.47%                      (95%CI 1.14, 1.98)                      P&lt;.001 for all outcomes</p> <p>N=18,118 for both groups</p>
Dawson, 2020 <sup>143</sup> United Kingdom Prospectively collected/ retrospectively analyzed	NR	NR	NR	<p><b>New Aspiration Pneumonia</b>                      0% (0/208)</p>
De Lorenzo, 2020 <sup>82</sup> Italy Prospective	NR	NR	NR	<p><b>mMRC Dyspnea Scale</b>                      Mild: 25% (31/126)                      Moderate: 3% (4/126)                      Severe: 2% (3/126)                      Very Severe: 2% (2/126)</p>
Dennis, 2021 <sup>57</sup> United Kingdom Prospective	NR	NR	NR	<p><b>Deep Breathing Fractional Area Change &lt;31%</b>                      12% (4/34)                      (n=3 missing data)</p>

Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
Frija-Masson, 2020 <sup>60</sup> France Retrospective	NR	<b>Pulmonary Function Test Interpretation</b> Normal: 48% (24/50)	NR	Restrictive pattern: 8% (4/50) Restriction with altered diffusion capacity: 18% (9/50) Altered diffusion capacity only: 26% (13/50)
Fuglebjerg, 2020 <sup>29</sup> Denmark Case series  Hypoxia and dyspnea elicited by 6-minute walking test  Exercise-induced hypoxia: SpO <sub>2</sub> <90% (test terminated)	NR	NR	NR	<b>Exercise-Induced Hypoxia, % (n/N)</b> 50% (13/26) NOTE: PE confirmed in 67% (4/6) who underwent further testing  SpO <sub>2</sub> <90% was not associated with an increase in subjective dyspnea (Borg scale)
Garrigues, 2020 <sup>61</sup> France Prospective, survey	NR	NR	NR	<b>mMRC Dyspnea Scale Grade 2 or More</b> 29% (35/120)  Ward patients: 28% (27/96) ICU patients: 33% (8/24)
Goicoechea, 2020 <sup>30</sup> Spain Retrospective	NR	NR	NR	<b>“Lung Abnormalities”</b> 86% (6/7) (Worsening or appearance of X-ray pulmonary infiltrates)
Hall, 2021 <sup>115</sup> United Kingdom Retrospective	NR	FVC ≤80% 27% of (16/59) of patients with complete lung function tests  TLC ≤70%	NR	<b>Reduction in mMRC Dyspnea score (range 1 to 5) ≥2 points</b> 18% (36/200)

Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
		44% of (26/59) of patients with complete lung function tests  Oxygen desaturation ≥4% 20% (34/170) of patients that underwent 6-minute walk test		<b>Persistent interstitial change (parenchymal abnormality) via CT</b> 32% (64/200)
Han, 2021 <sup>116</sup> China Prospective	“Fibrotic like changes at 6 months” 35% (40/114) De novo in 95% (38/40)	NR	<b>DLCO &lt;80%</b> 26% (27/104)	<b>Dyspnea</b> 14% (16/114)  <b>Pleural Effusions (CT scan)</b> 9% (10/114)
Hu, 2020 <sup>45</sup> China Cross-sectional	61% (46/76)	NR	NR	NR
Huang C, 2021 <sup>85</sup> China Retro and Prospective Scale 3 = No supplemental O <sub>2</sub> Scale 4 = Requiring supplemental O <sub>2</sub> Scale 5-6 = Requiring HFNC, NIV or IMV	NR	<b>FEV<sub>1</sub> &lt;80% Predicted</b> Scale 3: 8% (7/89) Scale 4: 2% (4/172); P<.05 vs Scale 3 Scale 5-6: 13% (11/88) <b>FVC&lt;80% Predicted</b> Scale 3: 3% (3/89) Scale 4: 1% (1/172) Scale 5-6: 11% (10/88) <b>FEV<sub>1</sub>/FVC &lt;70%</b> Scale 3: 8% (7/89) Scale 4: 8% (13/172) Scale 5-6: 2% (2/88) <b>TLC &lt;80% of Predicted</b> Scale 3: 11% (9/83) Scale 4: 10% (17/165) Scale 5-6: 35% (30/86)	<b>DLCO &lt;80% Predicted</b> Scale 3: 22% (18/83) Scale 4: 29% (48/165) Scale 5-6: 56% (48/86); P<.05 vs Scale 3	<b>mMRC scores ≥1</b> Overall: 26% (419/1615) Scale 3: 24% (102/425) Scale 4: 26% (277/1079) Scale 5-6: 36% (40/111), P<.05 vs Scale 3  <b>At Least 1 Abnormal CT Pattern</b> Scale 3: 52% (49/89) Scale 4: 54% (87/161) Scale 5-6: 54% (50/92)
Huang Y, 2020 <sup>63</sup> China Retrospective	7% (4/57)	<b>FEV<sub>1</sub> &lt;80% Predicted</b> 9% (5/57) (mild impairment) <b>FVC&lt;80% Predicted</b> 11% (6/57) (5 mild impairment, 1 moderate)	<b>DLCO &lt;80% Predicted</b> 53% (30/57) (26 mild impairment, 4 moderate)  Subgroups	<b>Respiratory Muscle Strength</b> Pimax <80% Predicted 49% (28/57) Pemax <80% Predicted 23% (13/57)



Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
		<p><b>FEV<sub>1</sub>/FVC &lt;80%</b> 44% (25/57) (mild impairment) <b>TLC &lt;80% of Predicted</b> 12% (7/57) (6 mild, 1 moderate)</p> <p>Outcomes did not differ by severity of COVID-19</p>	<p>Severe COVID-19: 77% (13/17) Non-severe: 43% (17/40) P=.02</p>	<p><b>CT Residual Abnormality</b> 54% (31/57) Subgroups Severe COVID-19: 94% (16/17) Non-severe: 38% (15/40) P Not Reported</p> <p><b>Obstructive Pulmonary Dysfunction</b> 11% (6/57)</p> <p><b>Restrictive Pulmonary Function</b> 12% (7/57)</p> <p><b>Combined Obstructive and Restrictive</b> 4% (2/57)</p>
Jacobs, 2020 <sup>117</sup> USA Prospective	NR	NR	NR	<p><b>Dyspnea (persisting from hospital discharge)</b> 45% (58/128) Or 32% (58/183 enrolled)</p>
Karaarslan, 2021 <sup>118</sup> Turkey Prospective	NR	NR	NR	<p><b>Dyspnea (2 weeks)</b> 38% (114/300)</p> <p><b>Dyspnea (4 weeks)</b> 26% (78/300)</p>
Li, 2021 <sup>119</sup> China Prospective	NR	NR	NR	<p><b>Lesions (incomplete resolution) (CT at 3-6 months)</b> 72% (44/61)</p>
Liu, 2020 <sup>141</sup> China Retrospective	NR	NR	NR	<p><b>Ground Glass Opacity (CT)</b> Discharge: 18% (9/51) First follow-up (~2 weeks): 10% (5/51) Second follow-up (~4 weeks): 10% (5/51)</p> <p><b>Consolidation</b> Discharge: 49% (25/51)</p>



Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
				First follow-up (~2 weeks): 8% (4/51) Second follow-up (~4 weeks): 2% (1/51)
Lv, 2020 <sup>88</sup> China Retrospective	NR	<p><b>At 2 Weeks Post-discharge</b></p> <p><b>IVC&lt;80% Predicted</b> 81% (111/137) Severe cases: 89% (24/27) Non-severe cases: 79% (87/110)</p> <p><b>FVC&lt;80% Predicted</b> 24% (33/137) Severe cases: 56% (15/27) Non-severe cases: 16% (18/110)</p>	NR	NR
Mo, 2020 <sup>34</sup> China Cross-sectional	NR	<p><b>FEV<sub>1</sub> &lt;80% Predicted</b> 14% (15/110)</p> <p><b>FVC &lt;80% Predicted</b> 9% (10/110)</p> <p><b>FEV<sub>1</sub>/FVC &lt;70%</b> 5% (5/110)</p> <p>Outcomes above did not differ by severity of COVID-19</p> <p><b>TLC &lt;80% Predicted</b> 25% (27/110)</p> <p>COVID-19 Severity Subgroups Mild: 17% (4/24) Pneumonia: 21% (14/67) Severe Pneumonia: 47% (9/19) P&lt;.05 overall and for Severe Pneumonia vs Pneumonia or vs Mild</p>	<p><b>DLCO &lt;80% Predicted 47%</b> (51/110)</p> <p>COVID-19 Severity Subgroups Mild: 30% (7/24) Pneumonia: 42% (28/67) Severe Pneumonia: 84% (16/19) P=.001 overall</p> <p>P&lt;.01 for Severe Pneumonia vs Pneumonia or vs Mild</p>	NR
Morin, 2021 <sup>120</sup> (COMEBAC Study Group) France Prospective	<p><b>Fibrotic Lesions</b> 19% (33/170) (clinic assessment; median 125 days) Intubated: 36.7% (18/49)</p>	NR	<p><b>DLCO &lt;70%</b> 22% (33/152) (clinic assessment; median 125 days)</p>	<p><b>Nijmegen Score (Dysfunctional Breathing=score &gt;22)</b> 21% (37/177) (clinic assessment; median 125 days) <b>Dyspnea (new onset during or after hospitalization for COVID-19 and persistent at time of assessment)</b></p>



Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
	Non-intubated: 12.4% (15/121)			16% (78/478) (telephone assessment; median 113 days) <b>Abnormal Lung CT scan</b> 53% (108/171) (clinic assessment; median 125 days) <b>Persistent GGO</b> 42% (72/170) (clinic assessment; median 125 days)
Osikomaiya, 2021 <sup>122</sup> Nigeria Retrospective	NR	NR	NR	<b>Dyspnea</b> 10% (26/274)
Qin 2021 <sup>103</sup> China Prospective	<b>Pulmonary Interstitial damage</b> (from subset of 45 patients who received chest CT): 71% (32/45)	<b>TLC &lt;80% Predicted</b> 10% (8/81)  <b>FVC &lt;80% Predicted</b> 21% (17/81)  <b>FEV<sub>1</sub>/FVC &lt;70%</b> 4% (3/81)  Pulmonary function test results were available for 81 patients (41 non severe and 40 severe patients)	<b>DLCO &lt;80% Predicted</b> 54% (44/81)	<b>Dyspnea</b> 9% (56/647)  <b>Dyspnea (non-severe)</b> 7% (26/399)  <b>Dyspnea (severe)</b> 12% (30/248)
Raman, 2021 <sup>124</sup> United Kingdom Prospective  Controls were negative for SARS-CoV-2 and asymptomatic; community	NR	<b>FVC &lt;80% Predicted</b> COVID-19 13% (7/56) Controls 0% (0/28) P=.09 <b>FEV<sub>1</sub>&lt;80% Predicted</b> COVID-19 11% (6/56) Controls 4% (1/28)	NR	<b>Dyspnea – mMRC ≥2 (significant breathlessness)</b> COVID-19 Group 64% (36/56) Control Group 10% (3/29) P<.0001 <b>Lung Parenchymal Abnormalities</b> COVID-19 Group 60% (32/53) Control Group



Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
dwelling (not hospitalized)		P=.42 Median 1.6 months post-discharge		11% (3/28) P<.0001 <b>VO<sub>2</sub> Peak &lt;80% of Predicted Maximum</b> COVID-19 Group 55% (28/51) Control Group 7% (2/27) P<.0001 Median 1.6 months post-discharge
Ramani, 2021 <sup>91</sup> USA Retrospective	NR	NR	<b>Reduced diffusion capacity:</b> 27% (7/26)	At 40 days (median) Normal lung function: 62% (16/26) Obstructions: 15% (4/26) Restriction: 19% (5/26) Mixed obstruction and restriction: 4% (1/26)
Remy-Jardin, 2021 <sup>126</sup> France Retrospective	12.7% (7/55)	NR	NR	<b>COVID-19 Lung Infiltration (“residual findings”) on CT (median of 144 days)</b> 73% (40/55) <b>Emphysema (median of 144 days)</b> 18% (10/55)
Sami, 2020 <sup>94</sup> Iran Prospective	NR	NR	NR	<b>Dyspnea (symptom questionnaire)</b> Week 1 Non-severe: 22% (86/400) Severe: 19% (10/53) Week 4 Non-severe: 15% (59/400) Severe: 19% (10/52)
Shah, 2020 <sup>95</sup> Canada Prospective	NR	<b>FEV<sub>1</sub>/FVC &lt;70%</b> 11% (7/60)	<b>Abnormal DLCO</b> 52% (31/60)	<b>Dyspnea (present/absent on symptom questionnaire)</b> At 12 weeks 21% (12/60)

Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
				<p><b>Imaging (CT)</b>  <b>Ground Glass Abnormality:</b> 83% (50/60)  <b>Reticulation:</b> 65% (39/60)  <b>Neither:</b> 12% (7/60)</p>
Sibilia, 2021 <sup>144</sup> Spain Retrospective	NR	<p><b>FEV &lt;80% Predicted</b> 25% (43/172)</p> <p><b>FVC &lt;80% Predicted</b> 24% (42/172)</p>	<p><b>DLCO &lt;80% Predicted</b> 57% (98/172)</p>	<p><b>Dyspnea</b> 40% (68/172)</p>
Sonnweber 2020 <sup>96</sup> Austria Prospective	NR	<p><b>FVC&lt;80% Predicted Normal</b> 60 days: 27% (34/125) 100 days: 22% (29/132)</p> <p><b>FEV<sub>1</sub> &lt;80% Predicted Normal</b> 60 days: 22% (28/127) 100 days: 22% (30/136)</p> <p><b>FEV<sub>1</sub>/FVC &lt;70%</b> 60 days: 4% (5/125) 100 days: 8% (11/138)</p> <p><b>TLC&lt;80% Predicted Normal</b> 60 days: 11% (14/127) 100 days: 11% (15/137)</p> <p><b>Lung Function Impaired</b> 60 days: 42% (53/126) 100 days: 36% (48/133)</p>	<p><b>DLCO&lt;80% Predicted Normal</b> 60 days: 31% (39/125) 100 days: 21% (28/133)</p>	<p><b>Dyspnea (assessed by questionnaire [or mMRC])</b> 60 days: 68% [2% (3/145; severe; mMRC 3-4) 100 days: 36% [4% (5/133) severe; mMRC 3-4]</p> <p><b>Pathological CT findings Overall</b> 60 days: 77% (112/145) 100 days: 63% (84/133)</p> <p><b>Ground Glass Opacities</b> 60 days: 77% 100 days: 63%</p> <p><b>Reticulation</b> 60 days: 58% 100 days: 51%</p> <p><b>Consolidations</b> 60 days: 54% 100 days: 7%</p>
Spinicci, 2021 <sup>127</sup> Italy Retrospective	NR	NR	NR	<p><b>Dyspnea</b> 30% (30/100)</p> <p><b>Respiratory Failure (pO<sub>2</sub>&lt;60 mmHg)</b> 0% (0/100)</p>

Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
Suarez-Robles, 2021 <sup>128</sup> France Retrospective	NR	NR	NR	<b>Dyspnea</b> 40% (54/134)
Tomasoni, 2021 <sup>130</sup> Italy Cross-sectional	NR	NR	NR	<b>Dyspnea (on-going)</b> 7% (7/105)
Venturelli, 2021 <sup>132</sup> Italy Prospective	NR	NR	<b>DL<sub>CO</sub> Reduced</b> 19% (136/716)	<b>Dyspnea (overall)</b> 30% (228/767)  <b>Dyspnea (mild)</b> 23% (176/767)  <b>Dyspnea (moderate)</b> 6% (42/767)  <b>Dyspnea (severe/very severe)</b> 1% (10/767)  <b>Pulmonary Obstruction</b> 4% (27/716)  <b>Pulmonary Restriction</b> 12% (85/716)
Wang, 2020 <sup>71</sup> China Prospective cohort	NR	NR	NR	<b>Chest CT Deteriorated</b> 1-2 weeks post-discharge 5.6% (2/36) (1 with enhanced inflammatory infiltrates, 1 with multiple bilateral GGO) 3-4 weeks post-discharge 0% (0/54)  Outcomes did not differ by severity of COVID-19

Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
Wu, 2021 <sup>133</sup> China Prospective	<b>Fibrosis</b> 0% (0/83) at all follow-up	<b>FVC&lt;80% Predicted</b> 3 months: 23% (19/83) 6 months: 16% (13/83) 12 months: 11% (9/83) <b>FEV<sub>1</sub>&lt;80% Predicted</b> 3 months: 30% (25/83) 6 months: 24% (20/83) 12 months: 16% (13/83) <b>TLC&lt;80% Predicted</b> 3 months: 27% (22/83) 6 months: 19% (16/83) 12 months: 15% (12/83) <b>NOTE: no pulmonary function tests at 9 months</b>	<b>DLCO&lt;80% Predicted</b> 3 months: 55% (46/83) 6 months: 54% (45/83) 12 months: 33% (27/83)	<b>Dyspnea – mMRC</b> Score of at least 1 (shortness of breath when hurrying on the level or walking up a slight hill) 3 months: 81% (67/83) 6 months: 30% (25/83) 9 months: 12% (10/83) 12 months: 5% (4/83) <b>Residual Changes on CT</b> 3 months: 78% (65/83) 6 months: 48% (40/83) 9 months: 27% (22/83) 12 months: 24% (20/83)
Xia, 2020 <sup>55</sup> China Retrospective	NR	NR	NR	<b>CT At Discharge</b> Residual infiltrates without fibrosis: 82% (233/282) Residual infiltrates and consolidation fibrosis: 14% (39/282)
Yasin, 2021 <sup>134</sup> Egypt Retrospective	48% (101/210)	NR	NR	NR
You, 2020 <sup>73</sup> China Case Series	22% (4/18) GGO plus pulmonary fibrosis: 61% (11/18) Normal: 6% (1/18) Not available: 11% (2/18)	<b>VC<sub>max</sub> &lt;80% Predicted</b> 17% (3/18) <b>FEV<sub>1</sub> &lt;80% Predicted</b> 17% (3/18) <b>FVC &lt;80% Predicted</b> 17% (3/18) <b>FEV<sub>1</sub>/FVC &lt;70%</b> 17% (3/18) Outcomes did not differ by severity of COVID-19	NR	<b>Ventilation Impairment</b> Normal: 67% (12/18) Obstructive Ventilatory Impairment: 17% (3/18) Restrictive Ventilatory Impairment: 17% (3/18)
Yu, 2020 <sup>74</sup> China	44% (14/32)	NR	NR	NR



Author, Year Country Study Design	Pulmonary Fibrosis % (n/N)	Lung Volume	Diffusion Capacity	Other
Retrospective case series	Pulmonary fibrosis: combination of parenchymal bands, irregular interfaces, coarse reticular pattern, and traction bronchiectasis			
Zhang, 2020 <sup>100</sup> China Retrospective	<b>At 2 weeks</b> 31% (35/112)	NR	NR	<b>Chest CT at 2 weeks</b> Normal: 40% (45/112) Ground glass opacity: 35% (39/112)
Zhao, 2020 <sup>101</sup> China Retrospective	NR	<b>Abnormalities:</b> 26% (14/55) TLC: 7% (4/55) FEV <sub>1</sub> : 11% (6/55) FVC: 11%% (6/55)	<b>DLCO Abnormality:</b> 16% (9/55)	<b>CT at 3 months</b> Ground glass opacity: 13% (7/55)

*Abbreviations:* DLCO=diffusing capacity of the lung for carbon monoxide; FEV<sub>1</sub>=forced expiratory volume in 1 sec; FVC=forced vital capacity; GGO=ground-glass opacity; HFNC=high flow nasal cannula; IMV=invasive mechanical ventilation; mMRC=modified Medical Research Council; NIV=non-invasive ventilation; NR=not reported; PE=pulmonary embolism; TLC=total lung capacity

<sup>a</sup>Includes participants without history of the outcome in the past one year



**TABLE 4. CARDIOVASCULAR OUTCOMES**

Author, Year Country Study Design	Imaging	Blood Tests	Other
Al-Aly, 2021 <sup>104</sup> USA (Veterans) Retrospective  All outcomes vs individuals hospitalized with seasonal influenza	NR	NR	All outcomes vs individuals hospitalized with seasonal influenza <b>Acute coronary disease<sup>a</sup></b> HR (adjusted) 1.29 (95%CI 1.11, 1.5) Excess burden per 1000 hospitalized COVID-19 patients at 6 months 9.36 (95%CI 4.16, 13.86)  <b>Heart Failure<sup>a</sup></b> HR (adjusted) 1.19 (95%CI 1.03, 1.39) Excess burden per 1000 hospitalized COVID-19 patients at 6 months 6.31 (95% CI 1.02, 10.88)  COVID-19 group: N=13,654 Control group (influenza): N=13,997
Alharthy, 2021 <sup>24</sup> Saudi Arabia Prospective	<b>Pericardial Effusion (Ultrasound) at Discharge</b> 1.5% (1/64) (n=64 survivors)	NR	NR
Alharthy, 2020 <sup>75</sup> Saudi Arabia Prospective	<b>Pericardial Effusion (Ultrasound)</b> <b>2 months</b> 15.7% (20/127) <b>4 months</b> 11.0% (14/127)	NR	NR
Ayoubkhani, 2021 <sup>106</sup> United Kingdom Retrospective  Controls from general population not meeting inclusion criteria for COVID-19; unclear if	NR	NR	<b>MACE, new onset events</b> COVID-19 Group: 2.6% (945/36,130) Rate per 1000 person-years 65.9 (95%CI 61.8, 70.3) General Population Control Group: 0.5% (190/36,130) Rate per 1000 person-years 12.3 (95%CI 10.6, 14.1) P<.001 between groups



Author, Year Country Study Design	Imaging	Blood Tests	Other
any were hospitalized at time of study			<p><b>MACE, all events</b>                      COVID-19 Group: 4.8% (2315/47,780)                      Rate per 1000 person-years                      126.1 (95%CI 121.0, 131.4)                      General Population Control Group: 1.8% (855/47,780)                      Rate per 1000 person-years                      42.6 (95%CI 39.8, 45.5)                      P&lt;.001 between groups</p> <p>NOTE: decreased rate of MACE in patients admitted to ICU</p>
Daher, 2020 <sup>81</sup> Germany Prospective	<p><b>Echocardiography</b>                      LVEF – globally normal                      Admission: 94% (17/18)                      6 week follow-up: 88% (29/33)                      RVEF – globally normal                      Admission: 94% (17/18)                      6 week follow-up: 94% (31/33)</p> <p><b>Pericardial Effusion</b>                      0% (0/33)</p>	NR	NR
Daugherty 2021 <sup>110</sup> USA Retrospective  Controls did not have a COVID-19 diagnosis and were not admitted to a hospital for COVID-19	NR	NR	<p><b>New Diagnoses</b>  <b>Coronary disease overall (MI, acute coronary syndrome, cardiogenic shock)</b>                      COVID-19 Group: 1.05%                      Control Group: 0.18%                      Risk difference 0.87% (95%CI 0.54, 1.27)                      P&lt;.001</p> <p><b>Congestive Heart Failure</b>                      COVID-19 Group: 1.54%                      Control Group: 0.20%                      Risk difference 1.34% (95%CI 0.66, 1.55)                      P&lt;.001</p> <p><b>Myocarditis</b>                      COVID-19 Group: 0.09%</p>



Author, Year Country Study Design	Imaging	Blood Tests	Other
			Control Group: 0.01% Risk difference 0.08% (95%CI -0.06, 0.19) P=1.0  N=18,118 for both groups
de Graaf, 2021 <sup>111</sup> the Netherlands Prospective	<b>Echocardiography</b> <b>Abnormal LV function (LVEF&lt;52%): 22%</b> (18/81) (NOTE: known pre-existing condition for 2 patients)	<b>Elevated Troponin T (Cardiac Injury)</b> 19% (15/81)	NR
De Lorenzo, 2020 <sup>82</sup> Italy Prospective	NR	NR	<b>Uncontrolled Blood Pressure Requiring Therapeutic Change</b> 21% (26/126)
De Michieli, 2021 <sup>112</sup> USA Retrospective	NR	NR	<b>Acute MI</b> 0% (0/312) (median 49 days follow-up)
Dennis, 2021 <sup>57</sup> United Kingdom Prospective	<b>CMR</b> <b>Left Ventricular Ejection Fraction (%)</b> Impaired ( $\leq 51\%$ ): 11% (4/37) Normal ( $>51\%$ ): 89% (33/37) <b>Evidence of Myocarditis</b> 22% (8/37)	NR	NR
Eswaran, 2021 <sup>114</sup> USA Retrospective	NR	NR	<b>Non-ST Segment Myocardial Infarction</b> 0.08% (4/447) Within 30 days
Hall, 2021 <sup>115</sup> United Kingdom Retrospective	NR	NR	<b>Cardiac Causes of Breathlessness</b> 4% (8/200) or 10% (8/81 with breathlessness) at 4-6 weeks (previously undiagnosed or deterioration of existing problem, included pericarditis, persistent sinus tachycardia, hypertrophic cardiomyopathy and inferior regional wall motion abnormality, atrial septal defect, pulmonary hypertension, left ventricular



Author, Year Country Study Design	Imaging	Blood Tests	Other
			hypertrophy and worsening of pre-existing heart failure)
<p>Huang L, 2020<sup>62</sup> China Retrospective</p> <p>NOTE: of 26 patients tested, 15 (58%) were considered positive based on presence of positive conventional CMR findings (increased myocardial edema ratio [<math>&gt;2.0</math>] (n=7) and/or LGE presence (n=8)) and 11 (42%) were negative</p>	<p><b>CMR</b> <b>Myocardial Edema</b> 54% (14/26) 50% (7/14) with positive LGE 50% (7/14) with small pericardial effusion</p> <p><b>LGE</b> 31% (8/26) with focal linear subepicardial and patchy mid-wall LGE (includes 7 patients noted above)</p> <p>Native T1, T2, and ECV values were significantly elevated in recovered COVID-19 patients with positive CMR findings compared with healthy controls</p> <p>Right ventricular ejection fraction, cardiac index, and stroke volume area were decreased in recovered COVID-19 patients with positive CMR findings compared with healthy controls</p>	NR	NR
<p>Morin, 2021<sup>120</sup> (COMEBAC Study Group) France Prospective</p>	<p><b>Echocardiography</b> <b>LVEF &lt;50% (no patient &lt;40%)</b> 12% (10/83) (clinic assessment; median 125 days)</p>	NR	NR
<p>Ozer, 2021<sup>123</sup> Turkey Prospective</p>	<p><b>LV-GLS above -18% (subclinical myocardial deformation)</b> 38% (28/74) 57% (16/28) in group with myocardial injury based on troponin level during hospitalization 26% (12/46) in group without myocardial injury</p>	NR	NR

Author, Year Country Study Design	Imaging	Blood Tests	Other
<p>Puntmann, 2020<sup>66</sup> Germany Prospective</p> <p>NOTE: Data for 100 patients; 33% hospitalized</p>	<p><b>CMR</b></p> <p><b>Abnormal Native T1</b> COVID-19: 73% (73/100) Healthy Controls: 12% (6/50) Risk Factor-matched Controls: 58% (33/57) P&lt;.05 for COVID-19 vs Controls</p> <p><b>Abnormal Native T2:</b> COVID-19: 60% (60/100) Healthy Controls: 12% (6/50) Risk Factor-matched Controls: 26% (15/57) P&lt;.05 for COVID-19 vs Controls</p> <p><b>LGE</b></p> <p><i>Myocardial</i> COVID-19: 32% (32/100) Healthy Controls: 0% Risk Factor-matched Controls: 17% (9/57) P&lt;.05 for COVID-19 vs Controls</p> <p><i>Pericardial</i> COVID-19: 22% (22/100) Healthy Controls: 0% Risk Factor-matched Controls: 14% (8/57)</p> <p><b>Pericardial Effusion &gt;10 mm</b> COVID-19: 20% (20/100) Healthy Controls: 0% Risk Factor-matched Controls: 7% (4/57) P&lt;.05 for COVID-19 vs Controls</p>	<p><b>Detectable hsTNT (≥3 pg/mL)</b> COVID-19 71% (71/100) Healthy Controls 22% (11/50) Risk Factor-matched Controls 54% (31/57) P&lt;.05 for COVID-19 vs Controls</p> <p><b>Significantly elevated hsTNT (≥13.9 pg/mL)</b> COVID-19 5% (5/100) Healthy Controls 0% Risk Factor-matched Controls: 0% P&lt;.05 for COVID-19 vs Controls</p>	<p>NR</p>
<p>Raman, 2021<sup>124</sup> United Kingdom Prospective</p> <p>Controls were negative for SARS-CoV-2 and asymptomatic;</p>	<p><b>Left Ventricular Function</b> Normal and comparable between groups (data NR)</p> <p><b>Native T1 (Basal Myocardium) &gt;1197 ms (&gt;2 SD from control mean)</b> COVID-19: 26% (13/50) Controls: 4% (1/28) P=.015</p>	<p><b>Abnormal Troponin T</b> COVID-19 0% Controls 0% Median 1.6 months post-discharge</p>	<p>NR</p>



Author, Year Country Study Design	Imaging	Blood Tests	Other
community dwelling (not hospitalized)	<p><b>Native T1 (Mid Myocardium) &gt;1215 ms (&gt;2 SD from control mean)</b>                      COVID-19: 8% (4/51)                      Controls: 0% (0/28)                      P=.29</p> <p><b>Native T1 (Apical Myocardium) &gt;1275 ms (&gt;2 SD from control mean)</b>                      COVID-19: 2% (1/50)                      Controls: 4% (1/28)                      P=1.0</p> <p><b>LGE – Myocarditis Pattern</b>                      COVID-19: 12% (6/52)                      Controls: 7% (2/28)                      P=.47</p> <p><b>LGE - Pericardial Effusion &gt;10 mm</b>                      COVID-19: 2% (1/52)                      Controls: 0% (0/28)                      P=1.0</p> <p>Median 1.6 months post-discharge</p>		
Sonnweber 2020 <sup>96</sup> Austria Prospective	<p><b>Echocardiography</b>  <b>LVEF (&lt;53%)</b>                      60 days: 3% (4/145)                      100 days: 3% (4/134)</p> <p><b>Pericardial Effusion</b>                      60 days: 6% (8/145)                      100 days: 1% (1/134)</p>	NR	NR
Spinicci, 2021 <sup>127</sup> Italy Retrospective	NR	NR	<p><b>Palpitation</b>                      15% (15/100)</p> <p><b>Chest pain</b>                      12% (12/100)</p>
Tudoran, 2021 <sup>131</sup> Romania Prospective	<p><b>Diastolic Dysfunction</b>                      17% (21/125)</p> <p><b>Diastolic Dysfunction and Impaired Left Ventricular Systolic Function</b>                      9% (11/125)</p> <p>Assessed at 6-10 weeks post-discharge</p>	NR	NR

Author, Year Country Study Design	Imaging	Blood Tests	Other
Xiong, 2021 <sup>99</sup> China Prospective	NR	NR	<b>Newly Diagnosed Hypertension</b> COVID-19: 1% (7/538) Controls: 0% (0/184)
Zhou, 2021 <sup>135</sup> China Prospective	<p><b>LVEF &lt;50%</b> 1% (1/97) Median 11 days post-discharge</p> <p><b>CMRI (If Elevated Troponin or ECG Abnormality)</b> No evidence of acute myocarditis</p>	<p><b>Troponin &gt;99<sup>th</sup> Percentile of Upper Limit of Normal</b> 6% (6/97)</p>	<p><b>Atrial Fibrillation (newly detected)</b> 1% (1/97)</p> <p><b>Sinus Bradycardia (&lt;60 bpm) (newly detected)</b> 28% (27/97)</p> <p><b>T Wave Inversion (newly detected)</b> 8% (8/97) Median 11 days post-discharge</p>

Abbreviations: hsTNT=high-sensitivity Troponin T; LGE=late gadolinium enhancement; LVEF=left ventricular ejection fraction; LV-GLS=left ventricular global longitudinal strain; NR=not reported

<sup>a</sup> Includes participants without history of the outcome in the past one year

**TABLE 5. NEUROMUSCULAR OUTCOMES**

Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
Akhtar, 2021 <sup>39</sup> Qatar Retrospective	NR	<p><b>Good Prognosis Score ≤2 at Discharge</b>                      COVID-19 Group: 28% (9/32)</p> <p>Concurrent Non-COVID-19 Group: 52%                      (112/216)                      Pre-COVID-19 Era Group: 60%                      (348/585)                      P=.001</p>	NR
Al-Aly, 2021 <sup>104</sup> USA (Veterans) Retrospective  All outcomes vs individuals hospitalized with seasonal influenza	Stroke <sup>a</sup> HR (adjusted) 1.30 (95%CI 1.05, 1.6) Excess burden per 1000 COVID-19 persons at 6 months 4.79 (95%CI 1, 7.87)	NR	<p><b>Neurocognitive Disorders<sup>a</sup></b>                      Excess burden per 1000 COVID-19                      persons at 6 months                      16.16 (95%CI 10.40, 21.19)</p> <p><b>Memory Problems<sup>a</sup></b>                      HR (adjusted) 1.42                      (95%CI 1.23, 1.63)                      Excess burden per 1000 COVID-19                      persons at 6 months                      16.59 (95%CI 10.59, 21.84)</p> <p>COVID-19 group: N=13,654                      Control group (influenza): N=13,997</p>
Alemanno, 2021 <sup>105</sup> Italy Prospective	NR	NR	<p><b>MMSE</b>  <b>Cognitive Deficits (range in effect                      severity)</b>                      Group 1 9% (2/22) mild                      Group 2 8% (1/12) mild                      Group 3 35% (7/20) mild to moderate                      Group 4 50% (1/2) moderate</p> <p><b>MoCA</b>  <b>Cognitive Deficits</b>                      Group 1 55% (12/22)                      Group 2 83% (10/12)                      Group 3 85% (17/20)</p>



Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
			Group 4 100% (2/2)
Al Kasab, 2020 <sup>23</sup> USA, South America, Europe Prospective	NR	<p><b>“Functional Independence on Discharge” Score 0-2</b> 17% (2/12) NOTE: missing data for n=1</p> <p>Non-COVID-19 Group 30% (94/316) P=.52 NOTE: missing data for n=129</p>	NR
Benussi, 2020 <sup>25</sup> Italy Retrospective cohort	<p>Median (IQR) COVID-19 Group (n=43) 9.0 (1.0-19.0)</p> <p>Non-COVID-19 Group (n=68) 2.0 (0.0-6.8) P=.005</p>	<p><b>“Good Outcome” Score ≤2 at Discharge</b> COVID-19 Group: 25.6% (11/43)</p> <p>Non-COVID-19 Group: 70.6% (48/68) P&lt;.001</p>	NR
Bowles, 2020 <sup>77</sup> USA Retrospective cohort  NOTE: 1302 of 1409 patients had admission and discharge assessments	NR	NR	<p><b>Cognitive Function at Home Health Care Admission</b> Requires prompting: 23% (297/1302) Requires assistance and direction: 6% (76/1302)</p> <p><b>Cognitive Function at Home Health Care Discharge</b> Requires prompting: 10% (130/1302) Requires assistance and direction: 3% (42/1302)</p> <p><b>Confusion at Home Health Care Admission</b> In new and complex situations only: 41% (536/1302) Anytime: 5% (70/1302)</p>



Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
			<p><b>Confusion at Home Health Care Discharge</b>                      In new and complex situations only: 19% (251/1302)                      Anytime: 3% (38/1302)</p>
<p>Chevinsky 2021<sup>109</sup>                      USA                      Retrospective</p> <p>Controls were hospitalized individuals who did not meet inclusion criteria for COVID-19 and were not diagnosed with COVID-19 during the 4 months after index encounter</p>	<p>NR</p>	<p>NR</p>	<p><b>Neurocognitive Disorders</b>                      vs hospitalized non-COVID-19 control group patients                      1-30 days after discharge                      OR (adjusted) 1.6 (95%CI 1.2, 2.1)                      31-60 days after discharge                      OR (adjusted) 1.2 (95%CI 0.87, 1.7)                      60-90 days after discharge                      OR (adjusted) 1.1 (95%CI 0.77, 1.6)                      90-120 days after discharge                      OR (adjusted) 1.1 (95%CI 0.72, 1.7)                      N=27,284 for COVID-19 and control groups</p> <p><b>Myopathies</b>                      1-30 days after discharge                      OR (adjusted) 5.9 (95%CI 2.8, 12.4)                      Not reported at other follow-up intervals                      N=27,284 for COVID and control groups</p>
<p>Daher, 2020<sup>81</sup>                      Germany                      Prospective</p>	<p>NR</p>	<p>NR</p>	<p><b>Cognitive Disorders (unclear how defined)</b>                      6 week follow-up: 18% (6/33)</p>
<p>Daugherty 2021<sup>110</sup>                      USA                      Retrospective</p> <p>Controls did not have a COVID-19 diagnosis and were not admitted to a hospital for COVID-19</p>	<p>Stroke (ischemic and hemorrhagic)                      COVID Group: 1.12%                      Control Group: 0.29%                      Risk difference 0.83% (95%CI 0.4, 1.2)                      P&lt;.001                      N=18,118 for both groups</p>	<p>NR</p>	<p><b>New Clinical Diagnoses</b>  <b>Amnesia/memory difficulty</b>                      COVID-19 Group: 2.90%                      Control Group: 0.43%                      Risk difference 2.47% (95%CI 1.76, 2.96)                      P&lt;.001</p> <p><b>Dementia</b>                      COVID-19 Group: 0.23%</p>



Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
NOTE: this study also reported peripheral neuropathy, encephalopathy, seizure, and Guillain-Barre Syndrome			Control Group: 0.03% Risk difference 0.2% (95%CI 0.07, 0.3) P<.001  <b>Alzheimer</b> COVID-19 Group: 0.04% Control Group: 0.0% Risk difference 0.04% (95%CI 0.0, 0.1) P<.001  N=18,118 for both groups
de Graaf, 2021 <sup>111</sup> The Netherlands Prospective	NR	NR	<b>Cognitive Failures Questionnaire (CFQ-25)</b> (cognitive impairment: score ≥31) 27% (13/48) <b>Informant Questionnaire on Cognitive Functioning in the Elderly</b> (cognitive impairment: score >3.31) 26% (10/38)
De Lorenzo, 2020 <sup>82</sup> Italy Prospective	NR	NR	<b>Cognitive Impairment (MoCA&lt;24)</b> 29% (36/126)
Eswaran, 2021 <sup>114</sup> USA Retrospective	Of 9 total vascular thromboembolic events there was 1 ischemic stroke	NR	NR
Garrigues, 2020 <sup>61</sup> France Prospective, survey	NR	NR	<b>Attention Disorder</b> 27% (32/120) Ward patients: 29% (28/96) ICU patients: 17% (4/24) P=.327  <b>Memory Loss</b> 34% (41/120) Ward patients: 38% (36/96)



Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
			ICU patients: 21% (5/24) P=.194
Grewal, 2020 <sup>31</sup> USA Retrospective	Median (IQR) COVID-19 Group (n=13) 11 (4-23) 2020 Comparison Cohort: 3 (2-13) 2019 Comparison Cohort: 4 (1-11)	<b>“Poor Outcome” Score &gt;2 at Discharge</b> COVID-19 Group: 77% (10/13)  2020 Comparison Cohort: 47% (25/53) 2019 Comparison Cohort: 41% (36/88)	NR
Jacobs, 2020 <sup>117</sup> USA Prospective	NR	NR	<b>Confusion (persisting from hospital discharge)</b> 43% (16/37) or 9% (16/183 enrolled)
Liotta, 2020 <sup>33</sup> USA Retrospective	NR	<b>Score ≤2 at Discharge</b> Overall: 71.1% (362/509)  No neurological manifestation (during hospitalization): 70.0% (63/90) Any neurological manifestation: 71.4 (299/419)  No encephalopathy (during hospitalization): 89.3% (310/347) Encephalopathy: 32.1% (52/162)	
Mathew, 2020 <sup>46</sup> India Retrospective	NR	<b>Score ≤2 at Discharge</b> 19% (12/62)	NR
Morin, 2021 <sup>120</sup> (COMEBAC Study Group) France Prospective	NR	NR	<b>Q3PC Questionnaire</b> <b>Memory Difficulties</b> 18% (73/416) <b>Mental Slowness</b> 10% (42/415) <b>Concentration Problems</b> 10% (41/412)

Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
			(telephone assessment; median 113 days) <b>Cognitive Complaint (impaired McNair score, reported cognitive symptoms, or both)</b> 50% (79/159) <b>Cognitive Impairment (impairment of either MoCA or d2-R score)</b> 38% (61/159)
Mowla, 2020 <sup>48</sup> Multinational Retrospective	NR	<b>Score ≤2 at Discharge</b> COVID-19 Group (data available for 10/13 patients) 60% (6/10) Control Group: 77% (44/57) P=.26	NR
Nersesjan, 2021 <sup>142</sup> Denmark Prospective	NR	NR	<b>Stroke after Discharge</b> 4% (2/45 with 3 month follow-up data) <b>Encephalopathy after Discharge</b> 2% (1/45 with 3 month follow-up data) <b>Peripheral Neuropathy after Discharge</b> 9% (4/45 with 3 month follow-up data) <b>Readmission to neurological department</b> 24% (4/17 readmitted through 3 month follow-up)
Ntaios, 2020 <sup>36</sup> Multi-national registry Retrospective	NR	<b>Severe Disability at Discharge</b> COVID-19 Group: 51% (49/96 survivors)  <b>Median Score (propensity matched population (n=330))</b> COVID-19 Group: 4 [IQR 2-6] Matched Group: 2 [IQR 1-4] P<.001	NR
Osikomaiya, 2021 <sup>122</sup> Nigeria	NR	NR	<b>Attention or memory deficit</b> 5% (14/274)



Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
Retrospective			
Perry, 2020 <sup>53</sup> United Kingdom Retrospective case-control	NR	<b>Score ≤2 at Discharge</b> (estimated from plot) COVID-19 Group: 29% Control Group: 46%	NR
Raman, 2021 <sup>124</sup> United Kingdom Prospective  Controls were negative for SARS-CoV-2 and asymptomatic; community dwelling (not hospitalized)	NR	NR	<p><b>MoCA &lt;26 (Abnormal)</b>                      COVID-19 Group                      28% (16/58)                      Control Group                      17% (5/30)                      P=.30 (calculated)                      Median 1.6 months post-discharge</p> <p><b>Clinical Assessment of Brain MRI</b>  <b>Brain Abnormalities</b>                      COVID-19 Group                      24% (13/54)                      Control Group                      21% (6/28)                      P=.79                      Median 1.6 months post-discharge</p>
Ramani, 2021 <sup>91</sup> USA Retrospective	NR	NR	<p><b>Mild Cognitive Impairment</b>  <b>MoCA (&lt;26)</b>                      57% (16/28)</p> <p><b>Quality of Life in Neurological Disorders</b>                      22% (6/27)</p>
Rass, 2021 <sup>125</sup> Austria Prospective	NR	NR	<p><b>Diagnoses at 3 Months (not diagnosed before COVID-19)</b>  <b>Any Neurological Disease</b>                      15% (20/135)  <b>Polyneuropathy/myopathy</b>                      13% (17/135)  <b>Parkinsonism</b>                      1% (1/135)</p>



Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
			<p><b>Stroke with Clinical Symptoms</b> 1% (1/135)</p> <p><b>Mild Encephalopathy</b> 2% (2/135)</p> <p><b>MoCA</b> <b>Cognitive Deficits (score &lt;26)</b> 23% (29/135) Severe COVID-19:29% (8/31) Moderate COVID-19 30% (20/72) Mild COVID-19 3% (1/32)</p> <p><b>Forgetfulness</b> 25% (30/135) Severe COVID-19 26% (7/31) Moderate COVID-19 24% (16/72) Mild COVID-19 24% (7/32)</p>
Spinicci, 2021 <sup>127</sup> Italy Retrospective	NR	NR	<p><b>Mental confusion</b> 10% (10/100)</p>
Tacquet, 2021 USA Retrospective	NR	NR	<p><b>Ischemic Stroke (first)</b> 1.60% (741/46302)</p> <p><b>Intracranial Hemorrhage (first)</b> 0.63% (292/46,302)</p> <p><b>Dementia</b> 1.46% (676/46,302)</p> <p><b>Parkinsonism</b> 0.20% (93/46,302)</p> <p><b>Myoneural junction or muscle disease</b> 1.24% (574/46,302)</p>
Tomasoni, 2021 <sup>130</sup> Italy Cross-sectional			<p><b>Cognitive deficits (memory disorder, on-going)</b> 17% (18/105)</p> <p><b>MMSE (mild deficits 18-25)</b> 36% (9/25)</p>

Author, Year Country Study Design	NIH Stroke Scale	Modified Rankin Scale	Other
			<b>MMSE (pathological &lt;18)</b> 4% (1/25)
Venturelli, 2021 <sup>132</sup> Italy Prospective	NR	NR	<b>Confusion</b> 5% (23/510)  <b>MoCa (normal)</b> >99% (302/304)  <b>MoCa (pathologic =0)</b> <1% (2/304)

Abbreviation: IQR=interquartile range; MMSE=Mini Mental State Examination; MoCA=Montreal Cognitive Assessment; NR=not reported; Q3PC=European AIDS Clinical Society cognitive screening questions

<sup>a</sup> Includes participants without history of the outcome in the past one year

**TABLE 6. RENAL OUTCOMES**

Author, Year Country Study Design	Acute Kidney Disease	Chronic Kidney Disease	Need for Renal Replacement Therapy	Imaging Findings
<p>Al-Aly, 2021<sup>104</sup> USA (Veterans) Retrospective</p> <p>All outcomes vs. individuals hospitalized with seasonal influenza and including individuals without history of the outcome in past 1 year</p>	<p><b>Acute Renal Failure<sup>a</sup></b> Excess burden per 1000 COVID-19 persons at 6 months 27.93 (95%CI 20.67, 34.50)</p> <p><b>Acute Kidney Injury<sup>a</sup></b> HR (adjusted) 1.24 (95%CI 1.1, 1.4) Excess burden per 1000 COVID-19 persons at 6 months 11.21 (95%CI 5.36, 16.43)</p> <p>COVID-19 Group: N=13,654 Control Group (influenza): N=13,997</p>	<p><b>CKD<sup>a</sup></b> HR (adjusted) 1.35 (95%CI 1.1, 1.65) Excess burden per 1000 COVID-19 persons at 6 months 6.03 (95%CI 2.17, 9.2)</p> <p>COVID-19 Group: N=13,654 Control Group (influenza): N=13,997</p>	<p>NR</p>	<p>NR</p>
<p>Ayoubkhani, 2021<sup>106</sup> United Kingdom Retrospective</p> <p>Controls from general population not meeting inclusion criteria for COVID-19; unclear if any were hospitalized at time of study</p>	<p>NR</p>	<p><b>CKD, new onset events at 140 (mean) days</b> COVID-19 Group: 0.6% (240/41,705) Rate per 1000 person-years 14.6 (95%CI 12.8, 16.6) General Population Control Group: 0.3% (125/41,705) Rate per 1000 person-years 7.2 (95%CI 6, 8.5)</p> <p><b>CKD, all events</b> COVID-19 Group: 1.5% (710/47,780) Rate per 1000 person-years 38.7 (95%CI 35.9, 41.6) General Population Control Group: 0.9% (410/47,780) Rate per 1000 person-years 20.4 (95%CI 18.5, 22.5)</p>	<p>NR</p>	<p>NR</p>



Author, Year Country Study Design	Acute Kidney Disease	Chronic Kidney Disease	Need for Renal Replacement Therapy	Imaging Findings
		NOTE: Similar rates whether admitted to ICU or not		
<p>Chan, 2021<sup>102</sup> USA Retrospective</p> <p>Compared last hospital creatinine with baseline; grouped as recovered or with AKD Stage 1, 2, or 3</p> <p>Recovered: difference in creatinine <math>\leq 0.3</math>; change in % <math>\leq 25\%</math></p> <p>Stage 1: difference <math>&gt;0.3</math> and change <math>&gt;25\%</math> and <math>\leq 100\%</math></p> <p>Stage 2: change in % <math>&gt;100\%</math> and <math>\leq 200\%</math></p> <p>Stage 3: change in % <math>&gt;200\%</math></p>	<p><b>At Discharge</b> 35% (291/832)<sup>b</sup> AKD Stage 1: 23% AKD Stage 2: 6% AKD Stage 3: 6%</p> <p><b>Follow-up (median 21 [IQR 8-38] days)</b> Data available for n=77 with AKD at discharge Recovered: 36% (28/77) AKD Stage 1: 33% (25/77) AKD Stage 2: 13% (10/77) AKD Stage 3: 18% (14/77)</p> <p>Data available for n=135 who had recovered at discharge Remain recovered: 86% (116/135) New AKD Stage 1: 10% (14/135) New AKD Stage 2: 2% (3/135) New AKD Stage 3: 2% (3/135)</p>	NR	NR	NR
<p>Chevinsky 2021<sup>109</sup> USA Retrospective</p> <p>Controls were hospitalized individuals who did not meet inclusion criteria for COVID-19 and were not diagnosed with COVID-19 during the 4</p>	<p>Acute and unspecified kidney failure vs hospitalized non-COVID-19 control group patients</p> <p>1-30 days after discharge OR (adjusted) 1.3 (95%CI 1.0, 1.6)</p> <p>31-60 days after discharge OR (adjusted) 0.74 (95%CI 0.56, 0.99)</p> <p>60-90 days after discharge</p>	NR	NR	NR



Author, Year Country Study Design	Acute Kidney Disease	Chronic Kidney Disease	Need for Renal Replacement Therapy	Imaging Findings
months after index encounter	OR (adjusted) 0.67 (95%CI 0.48, 0.92) 90-120 days after discharge OR (adjusted) 0.56 (95%CI 0.39, 0.80) N=27,284 for COVID and control groups			
Daugherty 2021 <sup>110</sup> USA Retrospective  Controls did not have a COVID-19 diagnosis and were not admitted to a hospital for COVID- 19	<b>Kidney injury (acute and chronic)</b> COVID-19 Group: 3.02% Control Group: 0.79% Risk difference 2.22% (95%CI 1.42, 2.79)  <b>Acute kidney injury</b> COVID-19 Group: 2.85% Control Group: 0.47% Risk difference 2.38% (95%CI 1.67, 3.11)	<b>CKD</b> COVID-19 Group: 2.06% Control Group: 0.70% Risk difference 1.36% (95%CI 0.72, 1.82)  P<.001 for both outcomes N=18,118 for both groups	NR	NR
Dennis, 2021 <sup>57</sup> United Kingdom Prospective	NR	NR	NR	<b>Kidney Cortex T1</b> Normal: 95% (35/37) Impairment: 5% (2/37)
Doher, 2020 <sup>145</sup> Brazil Retrospective cohort	NR	NR	<b>At Discharge (among RRT group, n=34)</b> 11% (1/9) 34 required RRT during hospitalization; NOTE: 12 died in the hospital, unclear if 9 represents patients discharged by end of study period	NR
Gupta, 2021 <sup>139</sup> USA Cohort	NR	NR	<b>At Discharge</b> 34% (73/216 discharged) <b>At 60 Days after ICU Admission</b> 57% (39/69 alive at day 60) 18% (39/216 discharged)	NR



Author, Year Country Study Design	Acute Kidney Disease	Chronic Kidney Disease	Need for Renal Replacement Therapy	Imaging Findings
Hamilton, 2020 <sup>83</sup> United Kingdom Retrospective	NR	NR	<b>At Discharge</b> 6% (2/32 who required RRT during hospitalization)	NR
Hittesdorf, 2020 <sup>140</sup> USA Retrospective	NR	NR	<b>At Discharge and 90 Days after Admission</b> 4% (2/45 who required RRT during hospitalization) 7% (2/27 surviving at 90 days)	NR
Huang C, 2021 <sup>85</sup> China Retro and Prospective	NR	NR	NR	<b>Abnormal morphology (ultrasound)</b> Scale 3: 0% (0/40) Scale 4: 0% (0/113) Scale 5-6: 0% (0/28)
Matsunaga, 2020 <sup>47</sup> Japan Registry	NR	NR	<b>At Discharge</b> 1% (16/2431)	NR
Morin, 2021 <sup>120</sup> (COMEBAC Study Group) France Prospective	NR	<b>Persistent Alteration of Kidney Function at 4 Months</b> 2% (2/95 who had AKI during hospitalization) or 0.4% (2/478 overall)	NR	NR
Naar, 2020 <sup>49</sup> USA Prospective	NR	NR	<b>At Discharge</b> 11% (5/46 who required RRT during hospitalization) 3% (5/148 who developed AKI) 2% (5/206 enrolled)  NOTE: 3% (7/206) were dialysis-dependent before hospital admission	NR
Ng, 2020 <sup>35</sup> USA Retrospective	<b>At Discharge KRT</b>	NR	<b>RRT</b> 92% (33/36) who had not recovered needed RRT at discharge (30.6% [33/108] of	NR



Author, Year Country Study Design	Acute Kidney Disease	Chronic Kidney Disease	Need for Renal Replacement Therapy	Imaging Findings
	17% (108/638) survived; 33% (36/108) had not recovered kidney function <b>Non-KRT</b> 52% (1663/3216) survived; 26% (430/1663) had not recovered kidney function		survivors who required hospital RRT) NOTE: 58% (19/33) had underlying CKD on admission	
Nugent, 2021 <sup>121</sup> USA Retrospective  COVID and non-COVID groups developed AKI during hospitalization	NR	<b>Kidney Dysfunction Post-Discharge (3-6 months)</b> 8% (15/182) <b>Kidney Recovery after Discharge</b> (rate per 100 patient-days) COVID-19 Group (n=32) 0.95 (0.62, 1.46) Non-COVID Group (n=287) 1.74 (1.51, 2.00) HR (adj): 0.57 (0.35, 0.92); P=.02	NR	NR
Raman, 2021 <sup>124</sup> United Kingdom Prospective  Controls were negative for SARS-CoV-2 and asymptomatic; community dwelling (not hospitalized)	NR	<b>Residual Renal Impairment at 2-3 Months (not present prior to COVID-19)</b> 3% (2/58) NOTES: Outcome NR for control group; 6 patients developed AKI during hospitalization; 2 required RRT	NR	NR
Stevens, 2020 <sup>97</sup> USA Retrospective	NR	NR	<b>At 30 days (median) from RRT Initiation</b> 8% (9/115) (NOTE: 2 of the 9 had been discharged with dialysis-dependent AKI; others remained hospitalized)	NR

Abbreviations: AKD=acute kidney disease; CKD=chronic kidney disease; IQR=interquartile range; KRT=kidney replacement therapy

<sup>a</sup>Includes participants without history of the outcome in the past one year

<sup>b</sup>Of 1835 with AKI while hospitalized, 832 were discharged; 291 with acute kidney disease



**TABLE 7. ENDOCRINE OUTCOMES**

Author, Year Country Study Design	Diabetes Mellitus
<p>Al-Aly, 2021<sup>104</sup> USA (Veterans) Retrospective</p> <p>All outcomes vs. individuals hospitalized with seasonal influenza and including individuals without history of the outcome in past 1 year</p>	<p><b>Diabetes<sup>a</sup></b> HR (adjusted) 1.6 (95%CI 1.36, 1.87) Excess burden per 1000 hospitalized COVID-19 patients at 6 months 21.39 (95%CI 15.1, 26.77)</p> <p>COVID-19 Group: N=13,654 Control Group (influenza): N=13,997</p>
<p>Ayoubkhani, 2021<sup>106</sup> United Kingdom Retrospective</p> <p>Controls from general population not meeting inclusion criteria for COVID-19; unclear if any were hospitalized at time of study</p>	<p><b>Diabetes, new onset events</b> COVID-19 Group: 1.1% (400/36,100) Rate per 1000 person-years 28.7 (95%CI 26, 31.7) General Population Control Group: 0.3% (125/36,100) Rate per 1000 person-years 8.2 (95%CI 6.9, 9.8) P&lt;.001 between groups</p> <p><b>Diabetes, all events</b> COVID-19 Group: 4.9% (2330/47,780) Rate per 1000 person-years 126.9 (95%CI 121.8, 132.2) General Population Control Group: 3.6% (1725/47,780) Rate per 1000 person-years 86.0 (95%CI 82.0, 90.2) P&lt;.001 between groups</p> <p>NOTE: Increased rates of diabetes for patients admitted to ICU</p>
<p>Daugherty 2021<sup>110</sup> USA Retrospective</p> <p>Controls did not have a COVID-19 diagnosis and were not admitted to a hospital for COVID-19</p>	<p><b>New Clinical Diagnoses Diabetes (Type 2)</b> COVID Group: 3.04% Control Group: 0.83% Risk difference 2.21% (95%CI 1.4, 3.16) P&lt;.001 N=18,118 for both groups</p>

<sup>a</sup> Includes participants without history of the outcome in the past one year  
CI=confidence interval; HR=hazard ratio;

**TABLE 8. GASTROINTESTINAL OUTCOMES**

Author, Year Country Study Design	Imaging Findings	Other Findings
Al-Aly, 2021 <sup>104</sup> USA (Veterans) Retrospective  Outcomes vs individuals hospitalized with seasonal influenza and including individuals without history of the outcome in past 1 year	<p><b>“Gastrointestinal Disorders” (includes dysphagia)<sup>a</sup></b>                      Excess burden per 1000 COVID-19 persons at 6 months                      19.28 (95%CI 12.75, 25.13)</p> <p>COVID-19 Group: N=13,654                      Control Group (influenza): N=13,997</p>	
Ayoubkhani, 2021 <sup>106</sup> United Kingdom Retrospective  Controls from general population not meeting inclusion criteria for COVID-19; unclear if any were hospitalized at time of study	<p><b>Chronic Liver Disease, new onset</b>                      (mean 140 days follow-up)                      COVID-19 Group: 0.2% (70/46,395)                      Rate per 1000 person-years                      4.0 (95%CI 3.2, 5.1)                      General Population Control Group: 0.04%                      (15/46,395)                      Rate per 1000 person-years                      0.9 (95%CI 0.5, 1.4)                      P&lt;.001 between groups*</p> <p><b>Chronic Liver Disease, all events</b>                      COVID-19 Group: 0.3% (135/47,780)                      Rate per 1000 person-years                      7.2 (95%CI 6.1, 8.6)                      General Population Control Group: 0.1%                      (50/47,780)                      Rate per 1000 person-years                      2.5 (95%CI 1.9, 3.3)                      *calculated by review authors</p>	
Daugherty 2021 <sup>110</sup> USA Retrospective  Controls did not have a COVID-19 diagnosis and were not admitted to a hospital for COVID-19	<p style="text-align: center;"><b>NR</b></p>	<p><b>Liver Test Abnormality (at 4 months)</b>                      COVID-19 Group: 3.30%                      Control Group: 1.36%                      Risk difference 1.95% (95%CI 1.06, 2.58)                      P&lt;.001                      N=18,118 for both groups</p>
Dennis, 2021 <sup>57</sup> United Kingdom Prospective	<p><b>Liver Inflammation (cT1 in ms)</b>                      Normal (&lt;784 ms): 76% (28/36)                      Impaired (≥784 ms): 24% (9/36)                      Missing data for n=1</p>	
de Graaf, 2021 <sup>111</sup> The Netherlands Prospective	<p style="text-align: center;"><b>NR</b></p>	<p><b>Elevated Liver Enzyme at 1.5 Months</b>                      2% (2/81)</p>
Huang C, 2021 <sup>85</sup> China Retro and Prospective	<p><b>Abnormal Liver Morphology (ultrasound)</b>                      Scale 3: 0% (0/100)                      Scale 4: 0% (0/185)                      Scale 5-6: 0% (0/105)</p>	

<sup>a</sup> Includes participants without history of the outcome in the past one year



**TABLE 9. HEMATOLOGIC OUTCOMES**

Author, Year Country Study Design	Thromboembolism	Hemorrhage	Coagulation Disorder
Al-Aly, 2021 <sup>104</sup> USA (Veterans) Retrospective  All outcomes vs individuals hospitalized with seasonal influenza and including individuals without history of the outcome in past 1 year	<b>Thromboembolism<sup>a</sup></b> HR (adjusted) 2.26 (95% CI 1.94, 2.64) Excess burden per 1000 hospitalized COVID-19 patients at 6 months 29.77 (95% CI 25.74, 33.24)  <b>Pulmonary Embolism<sup>a</sup></b> Excess burden per 1000 COVID-19 persons at 6 months 18.31 (95%CI 15.83, 20.25)  COVID-19 Group: N=13,654 Control Group (influenza): N=13,997	NR	<b>Coagulation Disorder<sup>a</sup></b> Excess burden per 1000 COVID-19 persons at 6 months 14.31 (95%CI 10.08, 17.89)  COVID-19 Group: N=13,654 Control Group (influenza): N=13,997
Alharthy, 2021 <sup>24</sup> Saudi Arabia Prospective	<b>DVT at Discharge</b> 12.5% (8/64) (n=64 survivors)	NR	NR
Alharthy, 2020 <sup>75</sup> Saudi Arabia Prospective	<b>DVT</b> <b>2 months</b> 14.2% (18/127) <b>4 months</b> 7.1% (9/127)	NR	NR
Brosnahan, 2020 <sup>79</sup> USA Retrospective	<b>Re-presented with Concern for Thrombotic Event</b> 0.46% (9/1,975) (included DVT, PE, limb ischemia due to coronary thrombosis, acute stroke, rapidly evolving hemodynamic instability with elevated D-dimer at time of presentation)	NR	NR
Chevinsky 2021 <sup>109</sup> USA Retrospective  Controls were hospitalized individuals who did not meet inclusion criteria for COVID-19 and were not diagnosed with COVID-	<b>Acute Pulmonary Embolism vs hospitalized non-COVID-19 control group patients</b> 1-30 days after discharge OR (adjusted) 1.5 (95%CI 1.0, 2.1) 31-60 days after discharge OR (adjusted) 1.4 (95%CI 0.93, 2.1) 60-90 days after discharge OR (adjusted) 1.2 (95%CI 0.72, 1.9) 90-120 days after discharge OR (adjusted) 1.2 (95%CI 0.70, 2.1)	NR	<b>Coagulation and Hemorrhagic Disorders</b> vs hospitalized non-COVID-19 control group patients 1-30 days after discharge OR (adjusted) 1.3 (95%CI 1.0, 1.6) 31-60 days after discharge OR (adjusted) 1.3 (95%CI 0.95, 1.7) 60-90 days after discharge OR (adjusted) 0.65 (95%CI 0.46, 0.90) 90-120 days after discharge

Author, Year Country Study Design	Thromboembolism	Hemorrhage	Coagulation Disorder
19 during the 4 months after index encounter	N=27,284 for COVID and control groups		OR (adjusted) 0.66 (95%CI 0.45, 0.97) N=27,284 for COVID and control groups
Daher, 2020 <sup>81</sup> Germany Prospective	<b>Thromboembolic events at 6 weeks</b> 0% (0/33)	NR	NR
Daugherty 2021 <sup>110</sup> USA Retrospective  Controls did not have a COVID-19 diagnosis and were not admitted to a hospital for COVID-19	<b>DVT</b> (at 4 months) COVID-19 Group: 2.28% Control Group: 0.30% Risk difference 1.99% (95%CI 1.52, 2.46)  <b>PE</b> (at 4 months) COVID-19 Group: 1.25% Control Group: 0.14% Risk difference 1.11% (95%CI 0.69, 1.39)  P<.001 for all outcomes N=18,118 for both groups	NR	<b>Hypercoagulability</b> (at 4 months) COVID-19 Group: 3.15% Control Group: 0.37% Risk difference 2.78% (95%CI 2.29, 3.62) P<.001 N=18,118 for both groups
Engelen, 2021 <sup>113</sup> Belgium Prospective	<b>DVT</b> (asymptomatic, no prophylaxis, 6 weeks post-discharge) 0% (0/146) <b>PE</b> (symptomatic, receiving prophylaxis, 6 weeks post- discharge) 1% (1/146)	<b>Major Bleeding</b> (6 weeks post-discharge) 0% (0/146) including in 41 patients receiving post- discharge prophylaxis	NR
Eswaran, 2021 <sup>114</sup> USA Retrospective	<b>Total Vascular Thromboembolic Events</b> (within 30 days) 2% (9/447), 6 were arterial and 3 were venous events. (see cardiovascular and neurologic outcomes)  <b>Pulmonary Embolism</b> (within 30 days) 1% (3/447)	NR	NR
Hall, 2021 <sup>115</sup> United Kingdom Retrospective	<b>Pulmonary Embolism (at 4-6 weeks)</b> 2% (4/200)  <b>Lung Infarcts</b> 1% (2/200)	NR	NR
Hill, 2020 <sup>84</sup> USA Retrospective	<b>VTE</b> (median follow-up 21 days) 0.14% (3/2075 survivors)	NR	NR

Author, Year Country Study Design	Thromboembolism	Hemorrhage	Coagulation Disorder
	NOTE: authors report 1 additional VTE in patient who was evaluated and discharged		
Huang C, 2021 <sup>85</sup> China Retro and Prospective	<b>DVT of Lower Limbs (ultrasound)</b> Scale 3: 0% (0/100) Scale 4: 0% (0/185) Scale 5-6: 0% (0/105)	NR	NR
Patell, 2020 <sup>65</sup> USA Retrospective	2.5% (4/163) at median of 23 days [IQR 12-33] 1 each: PE, intracardiac thrombus, thrombosed arteriovenous fistula, ischemic stroke  Among 13 patients discharged on thromboprophylaxis: no observed thrombotic or hemorrhagic complications	3.7% (6/163) at median of 27 days [IQR 16-31] 2 "major bleeds" (both following falls), 4 "clinically relevant non-major bleeding"	NR
Rashidi, 2020 <sup>92</sup> Iran Retrospective	<b>Acute PE at 45 Days</b> 0.2% (3/1529) including 1 alive at follow-up who reported potential symptoms and 2 deaths due to pulmonary embolism	NR	NR
Remy-Jardin, 2021 <sup>126</sup> France Retrospective	<b>PE-type Perfusion Defects (median 144 days)</b> 11% (6/55) or 16.6% (6/36 with perfusion abnormalities) <b>Detectable Clot</b> 2% (1/55) (Note: 1 of the 6 patients with perfusion defects had detectable clot)	NR	NR
Roberts, 2020 <sup>68</sup> United Kingdom Prospective	<b>VTE</b> COVID-19 Group 0.48% (9/1877) at median of 8 days [range 3-33] 2 DVT, 7 PE  Control Group (Medical Admissions in 2019) 0.31% (56/18,159) 8 proximal, 10 distal, 5 line-associated upper-limb DVT, 33 PE  OR 1.6 (95%CI 0.77, 3.1); P=.2	NR	NR

Author, Year Country Study Design	Thromboembolism	Hemorrhage	Coagulation Disorder
Salisbury, 2020 <sup>93</sup> United Kingdom Retrospective	<b>VTE</b> (secondary analysis- subgroup [n=152] discharged without an indication for therapeutic anticoagulation and followed for 42 days) 3% (4/152) (all PE, median of 14 days after discharge [range 4-26 days])	<b>Bleeding</b> (secondary analysis- subgroup [n=152] discharged without an indication for therapeutic anticoagulation and followed for 42 days) 0% (0/152)	<b>NR</b>
Vlachou, 2021 <sup>98</sup> United Kingdom Retrospective	<b>Admitted with PE (post- recovery)</b> At up to 4 weeks 1% (4/370 enrolled) (NOTE: number discharged alive NR)	NR	NR

*Abbreviations:* DVT=deep venous thrombosis; IQR=interquartile range; PE=pulmonary embolism

<sup>a</sup> Includes participants without history of the outcome in the past one year

**TABLE 10. HEALTHCARE/RESOURCE UTILIZATION OUTCOMES**

Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
Anand, 2020 <sup>40</sup> USA Retrospective	NR	Home Without Services: 34% (22/64 discharged) Skilled Nursing Facility: 32% (20/64 discharged) Acute Rehabilitation: 14% (9/64 discharged) Home With Services: 8% (5/64 discharged) Inpatient Hospice: 6% (4/64 discharged) Long-term Acute Care Hospital: 5% (3/64 discharged) Home with Hospice: 2% (1/64 discharged)	NR	NR
Arab-Zozani, 2020 <sup>76</sup> Iran Cross-sectional	NR	NR	NR	<b>Self-care</b> 88% reported no problems with self-care post-discharge
Atalla, 2020 <sup>136</sup> USA Retrospective	5.6% (19/339)  Median of 5 days [IQR 3-13] post discharge  <b>Clinical Course During 2<sup>nd</sup> Admission</b> Length of Stay: 7 days Intensive Care: 31%  NOTE: 3 patients required a third admission	<b>For 19 Patients Readmitted</b> Skilled Nursing Facility: 26% (5/19) Home (n=11) or Hotel for COVID+Homeless (n=3): 74% (14/19)	NR	<b>Reasons for Readmission</b> Bacterial pneumonia secondary to COVID-19 infection: 21% (4/19) Prolonged COVID-19 Course: 21% (4/19) Psychiatric episodes: 16% (3/19) Metabolic encephalopathy: 11% (2/19) Thrombotic episodes: 11% (2/19) Alcohol intoxication, orthostatic hypotension, gastroenteritis, fall/trauma (1 each): 21% (4/19)
Barbaro, 2020 <sup>41</sup> Multi (International Registry)	NR	Home or acute rehabilitation center: 53% (311/588 discharged alive)	NR	NR

Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
Retrospective cohort		Long-term acute care center or unspecified: 17% (101/588 discharged alive) Another hospital: 30% (176/588 discharged alive)  NOTE: a total of 588/1035 were discharged alive		
Bhatt, 2021 <sup>42</sup> USA Retrospective	NR	<p><b>COVID-19 Patients without History of Heart Failure</b>                      Hospice: 4% (4,320/121,813 discharged)                      Skilled nursing or rehabilitative care: 19% (22,601/121,813 discharged)</p> <p><b>COVID Patients with a History of Heart Failure</b>                      Hospice: 7% (428/6,357 discharged)                      Skilled nursing or rehabilitative care: 41% (2,605/6,357 discharged)</p> <p><b>Non-COVID Patients with History of Heart Failure (Non-Heart Failure Hospitalization)</b>                      Hospice: 4% (4,068/95,556 discharged)                      Skilled nursing or rehabilitative care: 21% (20,352/95,556 discharged)</p>	NR	NR
Bowles, 2020 <sup>77</sup> USA Retrospective cohort	<b>During Home Health Care (post-discharge)</b> 10% (137/1409)	NR	NR	NR



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
Brendish, 2020 <sup>78</sup> United Kingdom Prospective cohort	<b>Within 30 days</b> COVID-19 Positive: 10.6% (30/352) COVID-19 Negative: 17.7% (105/702) MD -7.2% [95%CI -11.7, 2.2; P=.0033]	NR	NR	NR
Casas-Rojo, 2020 <sup>56</sup> Spain Retrospective cohort	3.9% (573/14,709) Of patients discharged: 4.8% (573/11,928)  Not discharged at end of follow-up (after readmission) 0.2% (31/15,150) Of patients readmitted: 5.4% (31/573)	NR	NR	NR
Chopra, 2020 <sup>80</sup> USA Retrospective	<b>Within 60 days</b> 15% (189/1250)	Home: 78% (975/1250) Skilled nursing or rehabilitation: 13% (158/1250) Doesn't total 100% or 1250	<b>Oxygen use:</b> 7% (32/488) <b>New use of CPAP or other when asleep:</b> 7% (34/488)	<b>Primary care follow-up within 60 days</b> 78% (382/488 who completed follow- up telephone survey) <b>Home Health Services</b> 20% (98/488) <b>Unable to return to normal activity</b> 39% (188/488)
Collins, 2020 <sup>26</sup> USA Retrospective	NR	Home: 65% (13/20) or 81% (13/16 discharged) Nursing facility (permanent residence): 5% (1/20) or 6% (1/16 discharged) Hotel for those with confirmed COVID-19: 10% (2/20) or 13% (2/16 discharged)	NR	NR
Daher, 2020 <sup>81</sup> Germany Prospective	NR	NR	<b>Needing Oxygen Therapy</b> Admission: 82% (27/33) 6 week follow-up: 3% (1/33)	

Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
Dawson, 2020 <sup>143</sup> United Kingdom Prospectively collected/ retrospectively analyzed	0% (0/208) Length of follow-up NR	NR	NR	NR
DeBolt, 2020 <sup>43</sup> USA Retrospective	NR	<p><b>Home without Oxygen Requirement</b> Pregnant: 92% (35/38) Non-pregnant COVID-19 Controls: 85% (77/91 discharged alive)</p> <p><b>Skilled Nursing Facility, Long-term Acute Care, or Home with Oxygen Requirement</b> Pregnant: 8% (3/38) Non-pregnant COVID-19 Controls: 15% (14/91 discharged alive)</p>	NR	NR
de Havenon, 2021 <sup>146</sup> USA Retrospective	NR	<p><b>Favorable Discharge (Home or Acute Rehabilitation)</b> COVID-19 group: 34% (707/2,086) Historical controls: 66% (110,546/166,586) P&lt;.001</p>	NR	NR
De Lorenzo, 2020 <sup>82</sup> Italy Prospective	NR	NR	NR	<p><b>Need for Follow-up</b> 60% (75/126) (defined as presence at follow-up evaluation of at least 1 of: respiratory rate &gt;20 breaths/min, uncontrolled blood pressure requiring therapeutic change, moderate to very severe dyspnea, malnutrition, or new-onset cognitive impairment)</p>



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
De Michieli, 2021 <sup>112</sup> USA Retrospective	<b>COVID-19 Related</b> 9.6% (30/312) (median 49 days follow-up)	NR	NR	NR
Egol, 2020 <sup>58</sup> USA Prospective	<b>Within 30 Days</b> COVID-19 Positive: 11.8% (2/17)  COVID-19 Suspected: 7.1% (1/14)  COVID-19 Negative: 2.8% (3/107) P=.21	NR	<b>Post-Acute Rehabilitation</b> COVID-19 Positive: 90.0% (9/17)  COVID-19 Suspected: 84.6% (11/14)  COVID-19 Negative: 78.3% (83/107) P=.61	NR
El Moheb, 2020 <sup>59</sup> USA Retrospective	<b>Emergency Department Readmission</b> 11% (10/92) Matched comparison group of non-COVID-19 ARDS patients: 11% (10/92) Length of follow-up not reported	NR	NR	NR
Fisher, 2020 <sup>28</sup> USA Retrospective	NR	<b>Nursing Home</b> COVID-19 positive: 14.7% (492/3345) or 23% (492/2142 discharged) COVID-19 negative: 12.8% (152/1265) or 17% (162/950 discharged) RR (total study population): 1.2 (95%CI 1.0, 1.4) Historical control: 14.6% (1436/9859) or 15% (1436/9544 discharged) RR (COVID positive vs control, total study population): 1.0 (95%CI 0.9, 1.1)	NR	NR



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
		<p><b>Home</b>                      COVID-19 positive: 49.3% (1650/3345) or 77% (1650/2142 discharged)                      COVID-19 negative: 62.3% (788/2365) or 83% (788/950 discharged)                      RR (total study population): 0.08 (95%CI 0.7, 0.8)                      Historical control: 82.2% (8108/9859) or 85% (8108/9544 discharged)                      RR (COVID positive vs control, total study population): 0.6 (95%CI 0.57, 0.62)</p>		
Grewal, 2020 <sup>31</sup> USA Retrospective	NR	Disposition reported for 10/13 survivors (remaining patients: 2 deaths, 1 unknown disposition) Home: 30% (3/10) (2/6 in 'COVID' group, 1/4 in 'Neuro' group) Acute rehabilitation: 50% (5/10) (3/6 in 'COVID' group, 2/4 in 'Neuro' group) Long-term acute care: 20% (2/10) (1/6 in 'COVID' group, 1/4 in 'Neuro' group)	NR	NR
Hamilton, 2020 <sup>83</sup> United Kingdom Retrospective	<b>Within 30 days</b> 8% (86/1032)	NR	NR	NR
Huang C, 2021 <sup>85</sup> China Retro and Prospective	NR	NR	NR	<b>Personal care problems</b> 1% (11/1622)

Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
Hegde, 2020 <sup>44</sup> USA Retrospective case series	NR	Skilled Nursing Facility: 14% (1/7) or 33% (1/3 discharged) Long-term Acute Care: 29% (2/7) or 67% (2/3) discharged	NR	NR
Katz, 2020 <sup>32</sup> USA Retrospective	NR	<p>COVID-19 group Home: 29% (25/86) or 45% (25/56 discharged) Rehabilitation: 36% (31/86) or 55% (31/56 discharged) (Additional 30 patients died or in hospice care)</p> <p>Non-COVID-19 group Home: 46% (228/499) or 52% (228/438 discharged) Rehabilitation: 42% (210/499) or 48% (210/438 discharged) (Additional 61 patients died or in hospice care)</p> <p>Overall P&lt;.001</p>	NR	NR
Khalili, 2020 <sup>86</sup> Iran Prospective cohort	<b>Within 90 days of admission</b> 4% (10/254)	NR	NR	NR
Knights, 2020 <sup>137</sup> United Kingdom Retrospective	5.4% (3/56 patients discharged home)	Home: 81% (56/69 discharged) Care Home: 14% (10/69 discharged) Other (not specified): 4% (3/69 discharged)	New “packages of care”: 2.9% (2/69 discharged) New care home placement: 7.2% (5/69 discharged) Increase in mobility aids: 11.6% (8/69 discharged)	NR
Loerinc, 2021 <sup>87</sup> USA Retrospective	<b>Within 30 days</b> <b>Emergency department visits:</b> 7.7% (24/310) with 54% (13/24) attributed to COVID-19	Home: 91% (281/310) Skilled nursing facility: 8% (25/310) State Public Health quarantine facility: 1% (4/310)	<b>Any home health or oxygen:</b> 24% (75/310) including physical or occupational therapy: 14% (42/310);	<b>Recommended follow-up appointments</b> Primary care: 83% (258/310)



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
	<p><b>Readmission</b> 5.2% (16/310) with 69% (11/16) attributed to COVID-19</p>		<p>home nursing service: 5% (16/310); new home oxygen therapy: 13% (41/310)</p> <p><b>New short-term medications:</b> 67% (207/310); average of 2.2 new prescriptions per patient</p> <p><b>New long-term medications:</b> 23% (72/310); average of 1.6 per patient</p>	<p>Specialist: 29% (90/310) including nephrology [7% (23/310)] and cardiology [5% (14/310)]</p> <p><b>Follow-up bloodwork ordered:</b> 10% (31/310)</p> <p><b>Follow-up radiology ordered:</b> 7% (21/310)</p>
<p>Lovinsky-Desir, 2020<sup>64</sup> USA Retrospective</p>	<p>Age 21-39 No Asthma: 5% (12/261) Asthma: 10% (4/39) P=.14</p> <p>Age 40-65 No Asthma: 5% (40/832) Asthma: 5% (5/111) P=1.0</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>
<p>Matsunaga, 2020<sup>47</sup> Japan Registry</p>	<p>NR</p>	<p>Home: 72% (1762/2437 discharged) Long-term care facility: 2% (44/2437 discharged) Transfer to another hospital: 18% (437/2437 discharged) Transfer to non-medical (isolation) facility: 8% (194/2437 discharged)</p>	<p><b>Oxygen therapy required:</b> 8% (182/2430)</p>	<p><b>Self-care Ability</b> Same as before onset of COVID-19: 84% (2045/2425) Worsened: 10% (237/2425) Improved: 4% (106/2425) Unknown: 2% (27/2425)</p>
<p>Monday, 2020<sup>89</sup> USA (Veterans) Retrospective</p>	<p><b>Within 30 days of admission</b> 14% (8/57 discharged alive)</p>	<p>NR</p>	<p><b>Home oxygen at discharge</b> 39% (22/57 discharged alive)</p>	<p>NR</p>



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
Nachega, 2020 <sup>50</sup> Democratic Republic of the Congo Retrospective cohort	NR	Transferred to Home Care 2.6% (20/766 enrolled) 3.0 % (20/665 recovered/alive)	NR	NR
Nemer, 2021 <sup>51</sup> USA Retrospective	NR	<b>Home:</b> 79% (278/350 enrolled) or 85% (278/328 discharged alive) <b>Subacute facility:</b> 11% (40/350 enrolled or 12% (40/328 discharged alive) <b>Hospice:</b> 2% (8/350 enrolled) or 2% (8/328 discharged alive)	NR	NR
Nersesjan, 2021 <sup>142</sup> Denmark Prospective	<b>At 3 months</b> 38% (17/45 with 3 month data)	NR	NR	NR
Overstad, 2020 <sup>52</sup> Norway Retrospective	4 were readmitted during study period (denominator unclear)	<b>Home</b> 74% (52/70) or 83% (52/63 discharged alive) ICU patients: 38% (5/13) or 63% (5/8 discharged alive) Ward patients: 86% (49/57) or 89% (49/55 discharged alive) <b>Discharged to 24-hour care</b> 16% (11/70) or 17% (11/63 discharged alive) ICU patients: 23% (3/13) or 38% (3/8 discharged alive) Ward patients: 12% (7/57) or 13% (7/55 discharged alive)	NR	NR
Parra, 2020 <sup>90</sup> Spain Case-control	4.4% (61/1368) Median time to readmission: 6 days	NR	NR	<b>Reasons for Readmission</b> Pneumonia: 56% (34/61)



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
				PE, DVT, or lower limb arterial thrombosis: 16% (10/61) Heart failure: 10% (6/61) Bacterial infection: 7% (4/61) Myocardial acute infarction, acute kidney failure, severe bleeding, diabetes, generalized edema, threatened miscarriage: each 2% (1/61)
Patell, 2020 <sup>65</sup> USA Retrospective	7% (12/163)	NR	NR	NR
Paterson, 2020 <sup>37</sup> United Kingdom Retrospective	NR	<b>Patients with Encephalopathy</b> (n=7 discharged) Home: 86% (6/7) Rehabilitation Unit: 14% (1/7) <b>Patients with Inflammatory Central Nervous System Syndromes</b> (n=2 discharged) Home: 100% (2/2) <b>Patients with Ischemic Stroke</b> (n=4 discharged) Rehabilitation Unit: 75% (3/4) Stroke Unit: 25% (1/4) <b>Patients with Peripheral Neurological Syndromes</b> (n=2 discharged; location NR) <b>Uncharacterized Condition</b> (n=1 discharged to home)	NR	NR
Richardson, 2020 <sup>67</sup> USA Case series	Overall: 2.2% (45/2081) <18 years: 3.1% (1/32) 18-65 years: 1.6% (22/1,373) >65 years: 3.3% (22/676)  <b>Time to Readmission</b> (median [IQR]) 3 [1.0, 4.5] days	<b>Facility (eg, nursing home, rehabilitation)</b> Overall: 5.9% (122/2081) <18 years: 0% (0/32) 18-65 years: 2.0% (28/1373) >65 years: 13.9% (94/676)  <b>Home</b>	NR	NR



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
		Overall: 94.1% (1959/2081) <18 years: 100% (32/32) 18-65 years: 98% (1345/1373) >65 years: 86% (582/676)		
Rodriguez, 2020 <sup>54</sup> USA Registry	NR	<b>Home:</b> 74% (4746/6421 discharged) <b>Hospice:</b> 3% (192/6421 discharged) <b>Nursing facility:</b> 17% (1097/6421 discharged) <b>Transfer to another hospital:</b> 5% (317/6421 discharged)	NR	NR
Sachdeva, 2020 <sup>69</sup> USA Retrospective	9% (1/9 discharged home) Length of follow-up NR	NR	NR	NR
Somani, 2020 <sup>70</sup> USA Retrospective	<b>Returned for Emergency Care</b> 3.6% (103/2864) Median 4.5 days  <b>Inpatient Admission</b> 54.4% (56/103) or 2% (56/3864) overall	NR	NR	<b>Reasons for Return</b> Respiratory distress: 50% Chest pain: 6% Other pain: 6% Altered mental status: 5% Falls: 5% Soft tissue infection: 5%  <b>Need for ICU level care</b> 10.7% (6/56)
Spinicci, 2021 <sup>127</sup> Italy Retrospective	10% (10/100)	NR	<b>Long-term Oxygen Therapy (discharge)</b> 15% (15/100)  <b>Long-term Oxygen Therapy (2 months)</b> 5% (5/100)	<b>Reasons for Readmission</b> <b>Cardiac disease (ie, heart failure and myocardial infarction)</b> 5% (5/100)  <b>Infectious diseases</b> 2% (2/100)  <b>Respiratory symptoms</b> 1% (1/100)



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
				<b>Neurologic disorders</b> 2% (2/100)
Suarez-Robles, 2021 <sup>128</sup> France Retrospective	<b>Bacterial respiratory infection, pulmonary thromboembolism, exacerbated COPD</b> 5% (7/134)	NR	Oxygen therapy 3% (4/134)	NR
Suleyman, 2020 <sup>138</sup> USA Retrospective case series	<b>30-day Hospital</b> Overall: 11.2% (29 cases) General practice unit: 27 cases ICU: 2 cases P<.001  NOTE: among patients initially discharged home from ED, 4% (4/108) were readmitted within 30 days	<b>Home</b> General practice unit: 96% (183/191 with known discharge disposition) ICU: 79% (49/62 with known discharge disposition)  <b>Rehabilitation center</b> General practice unit: 4% (8/191) ICU: 21% (13/62)	NR	<b>30-day Mortality (includes hospital mortality)</b> General practice unit: 7% (15/214) ICU: 40% (57/141) P<.001
Vizcaychipi, 2020 <sup>38</sup> United Kingdom Prospective	NR	<b>Home (usual residence)</b> 92.5% (614/664 discharged alive)  <b>Temporary Home</b> 2.4% (16/664)  <b>Residential Care Home</b> 5.1% (34/664)	NR	NR
Wang, 2020 <sup>71</sup> China Prospective cohort	1-2 weeks post-discharge 4% (5/131) 3-4 weeks post-discharge 2% (3/131)	<b>Home Quarantine</b> 1-2 weeks post-discharge 87% (114/131) 3-4 weeks post-discharge 92% (121/131) <b>Community Quarantine</b> 1-2 weeks post-discharge 9% (12/131) 3-4 weeks post-discharge 3% (4/131) <b>Designated Hospital</b>	<b>Oxygen therapy</b> 1-2 weeks post-discharge 7% (9/131) 3-4 weeks post-discharge 1% (1/131) <b>Corticosteroids</b> 1-2 weeks post-discharge 4% (5/131) 3-4 weeks post-discharge 2% (2/131)	<b>CBC abnormal</b> 1-2 weeks post-discharge 14% (2/14) 3-4 weeks post-discharge 8% (4/50) Outcomes did not differ by severity of COVID-19



Author, Year Country Study Design	Readmission	Discharge Disposition	Post-discharge Treatment	Other
		1-2 weeks post-discharge 4% (5/131) 3-4 weeks post-discharge 2% (3/131) <b>Return to Work</b> 1-2 weeks post-discharge 0% (0/131) 3-4 weeks post-discharge 2% (3/131) Outcomes did not differ by severity of COVID-19	Outcomes did not differ by severity of COVID-19	
Xu, 2020 <sup>72</sup> China Retrospective case series	NR	NR	<b>Oxygen therapy</b> 6% (5/85) (nasal cannula)	<b>60-day mortality (overall)</b> 62% (147/239) NOTE: Predictors included age >65, lymphocyte and platelet count, ARDS, acute cardiac injury, AKI, liver dysfunction, and coagulopathy

Abbreviations: ARDS=acute respiratory distress syndrome; ED=emergency department; ICU=intensive care unit; IQR=interquartile range

## APPENDIX D. PRISMA CHECKLIST

Section/topic	#	Checklist item	Reported on page #
<b>TITLE</b>			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Title page
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	Separate document
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known.	1
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	1
<b>METHODS</b>			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	3
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	4 and Table 1
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	3
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Appendix A
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	3-4
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	4
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	4 and Table 1
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	5
Summary	13	State the principal summary measures (e.g., risk ratio, difference in	5

measures		means).	
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I <sup>2</sup> ) for each meta-analysis.	5
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	N/A
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	N/A
<b>RESULTS</b>			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	6 (Figure 2)
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	7 and Appendix C
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	7
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	9-46 and Appendix C
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	N/A
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	N/A
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	N/A
<b>DISCUSSION</b>			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	46-48
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	48-49
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	52
<b>FUNDING</b>			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	Preface

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097