

APPENDIX A. SEARCH STRATEGIES

Database: Ovid MEDLINE(R) and Ovid OLDMEDLINE(R) <1946 to September Week 3 2012>,
Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations <October 02, 2012>

Search Strategy:

-
- 1 exp Brain edema/ (11605)
 - 2 exp cerebrovascular trauma/ (4960)
 - 3 exp craniocerebral trauma/ (113847)
 - 4 exp coma/ (17150)
 - 5 exp glasgow outcome scale/ (1042)
 - 6 exp glasgow coma scale/ (6068)
 - 7 ((brain* or capitis or cerebr* or crani* or hemispher* or inter-crani* or intra-crani* or skull*) adj4 (contusion* or damag* or fractur* or injur* or trauma* or wound*)).ab,ti. (77042)
 - 8 ((brain or crani* or cerebr* or head or inter-cran* or intra-cran*) adj4 (bleed* or haematoma* or haemorrhag* or hematoma* or hemorrhag* or pressure)).ti,ab. (23995)
 - 9 (Glasgow adj (coma or outcome) adj (scale* or score*)).ab,ti. (7439)
 - 10 'Rancho Los Amigos Scale'.ti,ab. (31)
 - 11 diffuse axonal injur*.ti,ab. (755)
 - 12 ((brain or cerebral or intracranial) adj3 (edema or oedema or swell*)).ab,ti. (11389)
 - 13 ((coma* or concuss* or unconscious* or 'persistent vegetative state') adj2 (damag* or fractur* or injur* or trauma* or wound*)).ti,ab. (1686)
 - 14 (mtbi or "mild trauma* injur*").tw. or "minor trauma* injur*".mp. (639)
 - 15 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 (209513)
 - 16 exp cohort studies/ (1212058)
 - 17 exp prognosis/ (967036)
 - 18 exp morbidity/ (328065)
 - 19 exp mortality/ (255922)
 - 20 exp survival analysis/ (158174)
 - 21 exp models, statistical/ (230839)
 - 22 prognos*.tw. (324957)
 - 23 course*.tw. (428896)
 - 24 diagnosed.tw. (303319)
 - 25 cohort*.tw. (221535)
 - 26 death.tw. (417341)
 - 27 predict*.tw. (808663)
 - 28 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 (3743025)
 - 29 diagnosed.tw. (303319)
 - 30 cohort:.mp. (281196)
 - 31 (predictor: or death).tw. (578864)
 - 32 exp models, statistical/ (230839)
 - 33 prognosis/ (327999)

- 34 29 or 30 or 31 or 32 or 33 (1494750)
35 28 or 34 (3743144)
36 15 and 35 (67135)
37 exp rehabilitation, vocational/ (8852)
38 exp employment/ (49891)
39 exp work/ (12406)
40 sick leave/ (3365)
41 absenteeism/ (6861)
42 exp occupational health/ (23070)
43 exp occupational medicine/ (21574)
44 exp disabled persons/ (43220)
45 “recovery of function”/ (25824)
46 exp human activities/ (291441)
47 exp self care/ (36358)
48 activities of daily living.tw. (13206)
49 (dressing or feeding or eating or toilet\$ or bathing or mobil\$ or driving or public
transport\$).tw. (409253)
50 ((daily or domestic or house or home) adj5 (activit\$ or task\$ or skill\$ or chore\$)).tw.
(34321)
51 (“work status” or “work capacity”).tw. (4948)
52 (unemployment or re-employment or underemployment or “job retention”).ti,ab. (6217)
53 (return* adj2 school).tw. (428)
54 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52
or 53 (884984)
55 15 and 54 (12268)
56 exp dementia/ (109281)
57 Delirium/ or exp Delirium, Dementia, Amnestic, Cognitive Disorders/ (165121)
58 dement*.mp. or alzheimer*.tw. (137451)
59 exp Parkinsonian Disorders/ (53442)
60 parkinson*.tw. (67897)
61 56 or 57 or 58 or 59 or 60 (263117)
62 15 and 61 (10666)
63 36 or 55 or 62 (80547)
64 animals/ not (humans/ and animals/) (3693774)
65 63 not 64 (72473)
66 limit 65 to (danish or english or french or norwegian or spanish or swedish) (62699)
67 limit 66 to yr=”2001 -Current” (36478)
68 exp “Outcome Assessment (Health Care)”/ (605282)
69 (intervention* adj3 stud*).tw. (23915)
70 68 or 69 (626005)
71 15 and 70 (12781)
72 71 not 64 (12055)
73 limit 72 to (yr=”2001 -Current” and (danish or english or french or norwegian or swedish))
(8537)
74 67 or 73 (36658)

- 75 randomized controlled trial.pt. (337763)
- 76 Randomized controlled trial/ (337763)
- 77 Randomized Controlled Trials as Topic/ (83241)
- 78 Double-Blind Method/ (117191)
- 79 clinical trial.pt. (474276)
- 80 “double blind:”.mp. (143695)
- 81 placebos/ (31353)
- 82 placebo:.mp. (158293)
- 83 random:.mp. (788646)
- 84 75 or 76 or 77 or 78 or 79 or 80 or 81 or 82 or 83 (1069355)
- 85 15 and 84 (10248)
- 86 review/ (1739065)
- 87 (medline or medlars or pubmed or grateful med or CINAHL or scisearch or psychinfo or psycinfo or psychlit or psyclit or handsearch* or hand search* or manual* search* or electronic database* or bibliographic database* or embase or lilacs or scopus or web of science).mp. (74569)
- 88 86 and 87 (48085)
- 89 meta-analysis.mp. (59484)
- 90 meta-analysis as topic/ (12450)
- 91 meta-analysis/ (36480)
- 92 systematic review*.tw. (38103)
- 93 cochrane database*.jn. (9039)
- 94 88 or 89 or 90 or 91 or 92 or 93 (112607)
- 95 15 and 94 (1435)
- 96 exp brain neoplasms/ (111650)
- 97 (cancer* or neoplasm* or tumor* or malign*).mp. and brain.tw. (75605)
- 98 exp Glioma/ (55756)
- 99 96 or 97 or 98 (164154)
- 100 15 and 99 (9656)
- 101 exp pain/ (283152)
- 102 exp chronic disease/ (210066)
- 103 101 and 102 (20005)
- 104 (chronic* adj3 pain*).mp. (32890)
- 105 103 or 104 (40897)
- 106 15 and 105 (366)
- 107 exp sports/ (100542)
- 108 exp recreation/ (116143)
- 109 (return* adj3 play*).tw. (803)
- 110 107 or 108 or 109 (116607)
- 111 15 and 110 (3782)
- 112 exp mental disorders/ (880032)
- 113 15 and 112 (15929)
- 114 exp disability evaluation/ (35932)
- 115 exp “Outcome Assessment (Health Care)”/ (605282)
- 116 disab:.tw. (119042)

- 117 114 or 115 or 116 (728782)
- 118 15 and 117 (17459)
- 119 74 or 85 or 95 or 100 or 106 or 111 or 113 or 118 (68980)
- 120 limit 119 to (english language and yr="2001 -Current" and (danish or english or french or norwegian or swedish)) (43016)
- 121 animals/ not (humans/ and animals/) (3693774)
- 122 120 not 121 (39793)
- 123 exp "United States Department of Veterans Affairs"/ or exp Veterans Health/ or exp Hospitals, Veterans/ or exp Veterans Disability Claims/ or exp Veterans/ (15311)
- 124 veteran.mp. (2331)
- 125 veterans.mp. (23349)
- 126 VA.mp. (16805)
- 127 VA.in. (58986)
- 128 VAMC.mp. (285)
- 129 VAMC.in. (2086)
- 130 exp Military Medicine/ or exp "United States Department of Defense"/ or exp Naval Medicine/ (30814)
- 131 exp Hospitals, Military/ (3861)
- 132 exp Military Facilities/ (3901)
- 133 (army or navy or air force or marines or coast guard).mp. [mp=title, abstract, original title, name of substance word, subject heading word, protocol supplementary concept, rare disease supplementary concept, unique identifier] (15842)
- 134 military.mp. or exp Military Personnel/ (22947)
- 135 soldier.mp. (1480)
- 136 soldiers.mp. (5245)
- 137 123 or 124 or 125 or 126 or 127 or 128 or 129 or 130 or 131 or 132 or 133 or 134 or 135 or 136 (151164)
- 138 122 and 137 (1170)
- 139 traumatic brain injury.mp. or exp Brain Injuries/ (48119)
- 140 tbi.mp. (10831)
- 141 139 or 140 (51178)
- 142 137 and 141 (1468)
- 143 138 or 142 (1977)
- 144 from 143 keep 1-1977 (1977)

The above search strategy was applied to two additional databases on Oct. 3, 2012, with the following yield:

PsycINFO=961

Cochrane Register of Controlled Trials (OVID)=46

APPENDIX B. STUDY SELECTION FORM

1. Language: Is the full text of the article in English?
Yes.....Proceed to #2
No.....Code **X1**. STOP
2. Population: Is the population adult, human participants who are Veterans or members of the military from any country? Studies that do not differentiate between adult and child populations, or between Veteran/military and civilian populations, will be excluded.
Yes.....Proceed to #3
No.....Code **X2**. Add code **B** if retaining for background/discussion. STOP
3. Publication type: Does the article present original study data, or is it a systematic-review or meta-analysis? Narrative or non-systematic reviews, letters, editorials, and commentaries will be excluded.
Yes.....Proceed to #4
No.....Code **X3**. Add code **B** if retaining for background/discussion. STOP
4. Case definition: Does the article stratify/examine mTBI separately from moderate to severe TBI cases? Participants can consist of a mixed group of TBI severity (mild, moderate or severe) only if the results are stratified by severity and the mTBI subjects can be clearly identified. Studies that include mixed groups of TBI severity and do not differentiate between mild, moderate and severe TBI in their analysis will be excluded. Patients must be clearly described as having mTBI, Post-Concussive Syndrome, or concussion; if none of these terms are used, patients must be clearly defined as falling within the definition of mTBI from the VA/DoD Clinical Practice Guideline for Management of Concussion/Mild Traumatic Brain Injury (2009) listed below.
Yes.....Proceed to #5
No.....Code **X4**. Add code **B** if retaining for background/discussion. STOP
5. Systematic review: Is the article a systematic review or meta-analysis of primary studies?
Yes.....Code **ISR (systematic review)**. STOP
No.....Proceed to #6
6. Sample size: Is the article a primary study with a sample size of at least 30 mTBI cases?
Yes.....Proceed to #7
No.....Code **X6**. Add code **B** if retaining for background/discussion. STOP
7. Applicability: Does the study report outcomes addressed in our Key Questions (e.g., health, cognitive, etc. for KQ1; or factors *associated* with outcomes in KQ1; or cost/utilization)?
Yes.....Proceed to #8
No.....Code **X7**. Add code **B** if retaining for background/discussion. STOP
8. Intervention studies: Is the study an intervention study?
Yes.....Proceed to #9
No.....Proceed to #10

9. Intervention outcomes: Does the intervention study report outcomes not influenced by intervention participation (e.g., baseline and/or control group outcome data)?
Yes.....Proceed to #10
No.....Code **X8**. Add code **B** if retaining for background/discussion. STOP
10. VA/DoD mTBI definition: Does the study define mTBI participants as meeting the VA/DoD Clinical Practice Guideline for Management of Concussion/Mild Traumatic Brain Injury (2009) definition of mTBI (listed below) with the exception of including/excluding/not reporting positive imaging results? Studies may use a different mTBI definition, but all criteria with the exception of positive imaging must fall within the VA/DoD definition.
Yes.....Code **IPS (primary study)**. STOP
No.....Code **X10**. Add code **B** if retaining for background/discussion. STOP

APPENDIX C. DEFINITION OF MTBI FROM THE VA/DOD CLINICAL PRACTICE GUIDELINE FOR MANAGEMENT OF CONCUSSION/MILD TRAUMATIC BRAIN INJURY (2009)

1.1 Definition of Traumatic Brain Injury

A traumatically induced structural injury and/or physiological disruption of brain function as a result of an external force that is indicated by new onset or worsening of at least one of the following clinical signs, immediately following the event:

- Any period of loss of or a decreased level of consciousness (LOC)
- Any loss of memory for events immediately before or after the injury (post-traumatic amnesia⁶)
- Any alteration in mental state at the time of the injury (confusion, disorientation, slowed thinking, etc.) (Alteration of consciousness/mental state⁵)
- Neurological deficits (weakness, loss of balance, change in vision, praxis, paresis/plegia, sensory loss, aphasia, etc.) that may or may not be transient
- Intracranial lesion

External forces may include any of the following events: the head being struck by an object, the head striking an object, the brain undergoing an acceleration/deceleration movement without direct external trauma to the head, a foreign body penetrating the brain, forces generated from events such as a blast or explosion, or other forces yet to be defined.

The above criteria define the event of a TBI. Not all individuals exposed to an external force will sustain a TBI, but any person who has a history of such an event with immediate manifestation of any of the above signs and symptoms can be said to have had a TBI.

1.2 Severity of Brain Injury Stratification

TBI is further categorized as to severity into mild, moderate, or severe based on the length of LOC, AOC, or PTA (see Table A-1). Acute injury severity is determined at the time of the injury.

- The patient is classified as mild/moderate/severe if s/he meets any of the criteria in Table A-1 within a particular severity level. If a patient meets criteria in more than one category of severity, the higher severity level is assigned.
- If it is not clinically possible to determine the brain injury level of severity because of medical complications (e.g., medically induced coma), other severity markers are required to make a determination of the severity of the brain injury.
- Abnormal structural imaging (e.g., Magnetic Resonance Imaging or Computed Tomography Scanning) attributed to the injury will result in the individual being considered clinically to have greater than mild injury.

In addition to traditional imaging studies, other imaging techniques such as functional magnetic resonance imaging, diffusion tensor imaging, positron emission tomography scanning; electrophysiological testing such as electroencephalography; and neuropsychological or other standardized testing of function have been used in the evaluation of persons with TBIs, but are

not considered in the currently accepted criteria for measuring severity at the time of the acute injury outlined in Table A -1.

The severity level has prognostic value, but does not necessarily predict the patient’s ultimate level of functioning. There is substantial evidence that the epidemiology, pathophysiology, natural history, and prognosis for concussion/mTBI are different than for moderate and severe TBI. For example, moderate and severe TBI are often associated with objective evidence of brain injury on brain scan or neurological examination (e.g., neurological deficits) and objective deficits on neuropsychological testing, whereas these evaluations are frequently not definitive in persons with concussion/mTBI. The natural history and prognosis of moderate and severe TBI are much more directly related to the nature and severity of the injury in moderate and severe TBI, whereas factors unrelated to the injury (such as co-existing mental disorders) have been shown to be the strong predictors of symptom persistence after a concussion/mTBI.

Table A-1. Classification of TBI Severity

Criteria	Mild	Moderate	Severe
Structural imaging	Normal	Normal or abnormal	Normal or abnormal
Loss of Consciousness (LOC)	0–30 min	> 30 min and < 24 hrs	> 24 hrs
Alteration of consciousness/mental state (AOC)	a moment up to 24 hrs	> 24 hours. Severity based on other criteria	
Post-traumatic amnesia (PTA)	0-1 day	> 1 and < 7 days	> 7 days
Glasgow Coma Scale (best available score in first 24 hours)	13-15	9-12	< 9

APPENDIX D. EXCLUDED STUDIES THAT DID NOT MEET MTBI DEFINITION CRITERIA

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Adams, Larson, Corrigan, et al., 2012 ¹	Length of LOC classified as < 1 minute, 1-20 minutes, or > 20 minutes. Used the Health Related Behaviors Among Active Duty Military Personnel Survey response categories: “The HRB Survey symptom response groups permit recoding LOC as up to 20 minutes and greater than 20 minutes. This provides insufficient information to code LOC using the American Congress of Rehabilitation Medicine’s definition of mild TBI.”	NR	Kay, Harrington, Adams, et al., 1993	Self-report survey
Arbisi, Polusny, Erbes, et al., 2011 ²	Adapted from the Defense and Veterans Brain Injury Center screening tool: yes to last item “dazed, confused, see stars, get knocked out or lose consciousness” classified as mTBI. No participants reported “receiving treatment while in Iraq for a TBI or were removed from assigned duties as a result of exposure to blast or other form of head trauma.”	NR	Schwab et al., 2007	Self-report mailed questionnaire
Armistead-Jehle, 2010 ³	“Screened positive on the VHA TBI screens” for “possible mTBI.” “All patients suffered at most a mild TBI..., as none reported loss of consciousness of more than 30 minutes or posttraumatic amnesia of 24 hours or more.”	NR	US Department of Veterans Affairs, 2007	Self-report
Bazarian, Donnelly, Peterson, et al., 2012 ⁴	“Mild TBI diagnosis was determined by in-person interview using a 22-item questionnaire developed to establish the nature, probability, and severity of deployment-related TBI among OEF/OIF veterans. The interview followed previously published TBI diagnostic criteria, which include confirmation of a possible TBI event, confirmation of alteration of consciousness, and confirmation of postconcussion symptoms. On the basis of the standardized clinical interview, interviewers rated the likelihood of mild TBI according to a 6-point scale: “not at all likely,” “very unlikely,” “somewhat unlikely,” “somewhat likely,” “very likely,” and “almost certainly.” These likelihood categories were used in all analyses. However, for descriptive purposes, subjects were defined as having mild TBI if interviewers rated them “very likely” or “almost certainly.””	NR	Lew, Poole, & Vanderploeg, 2007	Clinical interview

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Booth-Kewley, Highfill-McRoy, Larson, et al., 2012 ⁵	“Mild TBI symptoms were assessed using a set of questions that asked participants whether they had received an injury to the head during their most recent deployment that involved ‘being dazed, confused, or ‘seeing stars’ or ‘not remembering the injury, or losing consciousness (knocked out).’ A participant was classified as having a positive TBI screen if any of the three questions elicited a positive response.”	NR	Centers for Disease Control and Prevention & World Health Organization definitions adapted by the Defense and Veterans Brain Injury Center working group for military use	Self-report survey
Brenner, Terrio, Homaifar, et al., 2010 ⁶	Warrior Administered Retrospective Casualty Assessment Tool (WARCAT) and Brief Trauma Brain Injury Screen. “All 45 participants whose test scores were included in analyses had a history of blast exposure with alteration of or loss of consciousness (LOC).” “The nature of the most serious mTBIs reported were: <i>n</i> = 30, altered consciousness only; <i>n</i> = 12, up to 1-min LOC; and <i>n</i> = 3, one to 20-min LOC.”	NR	Soldier Readiness Process, 2007; Schwab et al., 2007	Chart review, self-report survey, and clinical interview
Brenner, Ivins, Schwab, Warden, Nelson, Jaffee, & Terrio, 2010 ⁷	As described in Terrio et al., 2009: Warrior Administered Retrospective Casualty Assessment Tool (WARCAT) and Brief Trauma Brain Injury Screen	NR	Soldier Readiness Process, 2007; Schwab et al., 2007	Chart review, self-report survey, and clinical interview
Cameron, Marshall, Sturdivant, & Lincoln, 2011 ⁸	“Incident cases of mTBI were operationally defined according to the administrative case definition proposed by the Centers for Disease Control and Prevention (CDC) for research purposes... and include ICD-9-CM codes for skull fracture (800.00, 800.5, 801.0, 801.5, 803.0, 803.5, 804.0, and 804.5), concussion (850.0, 850.1, 850.5, and 850.9), intracranial injury of unspecified nature (854.0), and head injury unspecified (959.01). In addition to the four digit codes listed, all subordinate five-digit codes were also included.”	NR	Centers for Disease Control and Prevention, 2003	Chart review
Carlson, Kehle, Meis, et al., 2011 ⁹	“Included studies must have assessed participants for a “probable” TBI (identified using self-report screening instruments) or diagnosed TBI history.”	NR	NR	NA
Clement & Kennedy, 2003 ¹⁰	LOC<60 minutes with no neurological findings	Excluded	NR	Chart review

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Dougherty, MacGregor, Han, et al., 2011 ¹¹	International Classification of Diseases, 9 th Revision (ICD-9) codes 800.0-801.9, 803.0-804.9, 850.0-854.1. “The Abbreviated Injury Scale (AIS) was used to describe the severity of brain injury... Due to a small number of TBI observations with scores of 4 (severe injury) and 5 (critical injury) in the present study, TBI severity was classified as follows: 0 = No TBI, 1 = minor, 2 = moderate and 3-5 = serious to critical. Service members with AIS scores of 6 were not eligible for inclusion in the study.”	NR	Gennarelli & Wodzin, 2005	Chart review
Drake, Gray, Yoder, et al., 2000 ¹²	“Subjects were consecutive MTBI patients... meeting specific inclusion criteria... a documented TBI classified by accepted criteria as a mild TBI.” No specific mTBI definition noted.	NR	NR	Self-report and chart review
Eskridge, 2011 ¹³	“Clinical diagnosis codes from the International Classification of Diseases, 9 th Revision (ICD-9) were assigned to each injury... In addition to the assigning of diagnosis codes, severity of each injury is assessed with two different standardized measures of injury severity; the Abbreviated Injury Scale (AIS) and the Injury Severity Scale (ISS)... The ISS for each blast episode was documented and categorized into one of four severity levels; mild (ISS 1-3), moderate (ISS 4-8), serious (ISS 9-15) and severe (ISS 16 and higher).”	NR	NR	Chart review
Fear, Jones, Groom, et al., 2009 ¹⁴	“Criteria for identification include confusion or disorientation, loss of consciousness lasting less than 30 min or post-traumatic amnesia lasting less than 24 h.” However, description of population studied only states, “we have examined the prevalence of symptoms thought to be a consequence of mTBI,” with unclear description of how mTBI was determined.	NR	Holm, et al., 2005	Chart review
Ferrier-Auerbach, Erbes, Polusny, et al., 2009 ¹⁵	“Three items adapted from the Defense and Veterans Brain Injury Center (DVBIC) Blast Exposure Screening Questionnaire... (1) Were you ever so close to a blast that you could feel the blast wave (such as heat or pressure) or afterward had trouble hearing or problems with attention or memory?... (2) Did you have any injuries from a blast, vehicle crash, bullet/shrapnel or fall?... (3) Did any injury cause you to be dazed/confused, ‘see stars,’ get knocked out, or lose consciousness?”	NR	Schwab et al., 2007	Self-report survey

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
French, Lange, Iverson, Ivins, et al., 2012 ¹⁶	LOC < 15 mins; PTA < 24 hours; absence of intracranial abnormality on computed tomography or magnetic resonance imaging scan	Excluded	Used VA/DoD criteria, but data limited to LOC < 15 minute categories.	Chart review
Gottshall, Gray, Drake, et al., 2007 ¹⁷	Definition: GCS 13-15. One patient identified as having an open head injury. LOC categorized as 31-60 (1 patient), and > 60 minutes (one patient), and up to 20 minutes (all remaining patients).	NR	American Academy of Neurology, 1997	Clinical presentation (to ED or Battalion Aid Station)
Helfer, Jordan, Lee, et al., 2011 ¹⁸	ICD-9CM codes 850.0, 850.11, 850.12, 850.2, 850.3, 850.4, 850.5, 850.9, 959.01, V15.52	NR	US Dept of Health and Human Services, 2008	Chart review
Heltemes, Dougherty, MacGregor, & Galameau, 2011 ¹⁹	International Classification of Diseases, 9 th Revision, Clinical Modification (ICD-9-CM) codes 800.0-801.9, 803.0-804.9, 850.0-854.1.	NR	Centers for Disease Control and Prevention, 2003	Chart review
Hoffer, Balaban, Gottshall, et al., 2010 ²⁰	“Definitive diagnosis of mild traumatic brain as defined by the 2007 Joint Service Surgeon General’s Definition”	NR	Assistant Secretary of Defense: Memorandum on Traumatic Brain Injury: Definition and Reporting. Available at http://www.pdhealth.mil/TBI.asp	In theater clinical evaluation
Hoffer, Donaldson, Gottshall, et al., 2009 ²¹	No specific mTBI definition listed.	NR	NR	Clinical interview
Hoge, McGurk, Thomas, et al., 2008 ²²	Positive response to three questions: LOC, being dazed and confused, seeing stars or not remembering the injury. However, “four Soldiers reported LOC lasting longer than 30 minutes. Although technically they were considered to have a moderate TBI they were not excluded because the number was low and it was not possible to verify the self-report data on any of the subjects”.	NR	DVBIC, 2006; CDC, 2003	Self-report
Ivins, Schwab, Baker, & Warden, 2003 ²³	“Head injury for which any LOC or any alteration of mental state without LOC was reported.” Reports rates of concussion grades separately, though the categories are divided by LOC < 20 minutes and LOC between 20-59 minutes.	NR	Kay et al., 1993	Chart review
Ivins, Schwab, Baker, & Warden, 2006 ²⁴	ICD-9CM codes 800.00-801.99, 803.00-804.99, and 850.0-854.19 and AIS severity codes of minor or moderate.	NR	Kay et al., 1993; Thurman & Guerrero, 1999	Chart review

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Ivins, Kane, & Schwab, 2009 ²⁵	Brief Traumatic Brain Injury Screen (BTBIS); LOC criteria < 20 mins. Additional computerized survey administered; however, it appears that only those with LOC < 20 minutes were included in the further assessment. Brief Traumatic Brain Injury Screen (BTBIS); Patients who screened positive for TBI and had LOC ≤ 20 minutes and/or possible PTA ≤ 24 hours were identified as having MTBI with LOC or possible PTA	NR	Kay et al., 1993; Thurman & Guerrero, 1999	Chart review
Ivins, 2010 ²⁶	Mapped ICD-9 CM diagnoses to the Abbreviated Injury Scale (AIS): ICD-9CM codes 800.00-801.99, 803.00-804.99, and 850.0-854.19 and AIS severity codes of minor or moderate. Combines mild and moderate into one category.	NR	Kay et al., 1993; Thurman & Guerrero, 1999	Chart review
Lange, Pancholi, Bhagwat, et al., 2012 ²⁷	“PTA < 24 hours and LOC < 15 minutes.” “It was our preference to use a LOC criterion of < 30 minutes, consistent with commonly used diagnostic criteria. . . However, the available information regarding LOC was limited to categorical data that did not allow us to differentiate between LOC greater or lower than 30 min.”	Included	Carroll et al., 2004; Management of Concussion/mTBI Working Group, 2009; ACRM, 1993	Chart review
Lange, Pancholi, Brickell, et al., 2012 ²⁸	LOC < 15 mins; PTA < 24 hours	Included	Used VA/DoD criteria, but data limited to LOC < 15 minute categories.	Chart review and clinical interview
Lange, Brickell, French, et al., 2012 ²⁹	Uncomplicated: PTA<24 hours, LOC < 15 mins, negative imaging; complicated: positive imaging. “It was our preference to use an LOC criterion of 30 min to classify MTBI consistent with commonly used military and civilian diagnostic criteria. However, the available information regarding LOC was limited to categorical data that did not allow us to differentiate between LOC greater or less than 30 min (i.e., available data = LOC < 15 min and LOC 16–60 min).”	Excluded	Used VA/DoD criteria, but data limited to LOC < 15 minute categories.	Routine comprehensive clinical evaluation
Lew, Garvert, Pogoda, et al., 2009 ³⁰	Mild TBI was defined as an initial GCS score of 13 to 15, PTA duration of <1 day, or LOC duration of <1 hour	NR	NR	Chart review and clinical evaluation

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Luis, Venderploeg, & Curtiss, 2003 ³¹	“During the interview, participants were asked, among many others, the following three questions: 1) since your discharge from active duty, have you been injured in a MVA? 2) Since your discharge from active duty have you injured your head (HI)? and (3) Did you lose consciousness as a result of the head injury?” Analysis by LOC group but time of LOC not specified.	NR	NR	Self-report
MacGregor, 2007 ³²	“An ICD-9 code in the following ranges was defined as a TBI (n = 124): 800.0-801.9 (fractures of the vault or base of the skull); 803.0-804.9 (other and unqualified and multiple fractures of the skull); and 850.0-854.1 (intracranial injury, including concussion, contusion, laceration, and hemorrhage)... Severity of TBI was indicated with the Abbreviated Injury Scale (AIS). The AIS ranges from 1 (relatively minor) to 6 (currently untreatable), and is determined separately for each different body region. Severity of TBI was determined by maximum AIS score for the head region – head AIS 1-2 indicated mild TBI, head AIS 3-5 indicated moderate-severe TBI. A majority of TBI identified via CHAMPS did not have a head AIS score present; in this case the TBI was assumed to be of mild severity due to a closed head injury.”	NR	ICD-9-CM, 2005	Chart review
MacGregor, Dougherty, & Galareau, 2011 ³³	International Classification of Diseases, 9 th Revision (ICD-9) codes 800.0-801.9, 803.0-804.9, 850.0-854.1. “The AIS was used to describe the severity of these injuries and the injuries were scored according to the following scale: 0, no injury; 1 minor; 2, moderate; 3, serious; 4, severe; 5, critical; 6, fatal injury. As per previous literature, each participant was categorized by the severity of their highest (or maximum) AIS Head score as mild (AIS score = 1-2), moderate (AIS score = 3), or severe (AIS score = 4-6).	NR	Ommaya, Ommaya, Dannenberg, et al., 1996	Chart review
MacGregor, Dougherty, Morrison, et al., 2011 ³⁴	“A concussion was defined by the ICD-9-CM code of 850.0-850.9. Severity of concussion was defined using the AIS.”	NR	NR	Chart review
MacGregor, Shaffer, Dougherty, et al. 2010 ³⁵	ICD-9 codes 800-801.9, 803-804.9, and 850-854.1; Abbreviated Injury Scale (AIS) score 1-2 = “mild,” 3-5 = “moderate to severe”	NR	Ommaya et al., 1996	Chart review

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
McGuire, Marsh, Sowin, & Robinson, 2012 ³⁶	"LOC less than 30 min or amnesia less than 60 min"	NR	Annegers et al., 1998	NR
Mora, Ritenour, Wade, et al., 2009 ³⁷	"Consciousness status to determine mTBI was queried using both codes from ICD and Abbreviated Injury Scale for indications of trauma to the head, concussive injuries, and indications of consciousness at the time of injury. A loss of consciousness served as the definition for mTBI.	NR	NR	Chart review
Morgan, Lockwood, Steinke, et al., 2012 ³⁸	No clear mTBI definition.	NR	NR	Self-report screening and clinical interview
Nelson, Weiser, Giford, et al., 2011 ³⁹	"The Brief Traumatic Brain Injury Screen was used to identify cases (probable mTBI) and controls (no mTBI). Individuals were placed in the "probable mTBI" group if they endorsed an injury (reported at least one injury on the BTBIS) and indicated that they had lost consciousness for a defined period of time (ranging from less than 1 minute to longer than 20 minutes) following the injury. Those who did not report an injury or a loss of consciousness (LOC) for any amount of time following an injury acted as the control group."	NR	Defense and Veterans Brain Injury Center (DVBIC)	Chart review
Olson-Madden, Forster, Huggins, & Schneider, 2012 ⁴⁰	"Injury severity data were coded as either mild (alteration in consciousness or loss of consciousness <= 30 minutes) or moderate/severe (loss of consciousness > 30 minutes)."	NR	Kay et al., 1993	Clinical interview
Ommaya, Ommaya, Dannenberg, et al., 1996 ⁴¹	"Head-injury-related discharge diagnosis (800.00-801.99, 803.00-804.99, and 850.0-854.19)." Injury Severity Scale and Abbreviated Injury Scale were calculated.	NR	NR	Chart review
Ommaya, Salazar, Dannenberg, et al., 1996 ⁴²	"Records with a head-injury hospital-related discharge diagnosis (800.00-801.99, 803.00-804.99, and 850.0-854.19) were identified as described in a previous study." "Maximum Abbreviated Injury Score (AIS) head and Injury Severity Score (ISS) were computed using the ICD-9 map. Mild TBI was defined as maximum AIS head equal to 1 or 2"	NR	NR	Clinical records
Pietrzak, Johnson, Goldstein, Malley, Southwick 2009 ⁴³	"Positive mTBI screen" based on endorsement of all 4 items on the DVBIC questionnaire	NR	GAO, 2008 & DVBIC, 2006	Self-report survey

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Plassman, Havlik, Steffens, et al., 2000 ⁴⁴	“1) mild injury = loss of consciousness or posttraumatic amnesia for less than 30 minutes, with no skull fracture; 2) moderate injury = loss of consciousness or post-traumatic amnesia for more than 30 minutes but less than 24 hours, and/or a skull fracture; and 3) severe injury = loss of consciousness or post-traumatic amnesia for more than 24 hours.”	NR	Frankowski, Annegers, & Whitman, 1985	Chart review
Polusny, Kehle, Nelson, Erbes, Arbisi, Thuras, 2011 ⁴⁵	Injury with altered mental status or LOC (items 1-3 on DVBIC screen)	NR	Schwab et al., 2007	Self-report in theater, 1 month prior to return home
Roebuck-Spencer, Vincent, Twille, et al., 2012 ⁴⁶	“Although data on recency and severity of injury were not available in this dataset, TBIs reported in this study are presumed to be mild in nature given that Service Members were still on active duty. Mild TBI was defined as such when individuals reported an injury event accompanied by an alteration of consciousness. This included endorsement of at least one of the following: feeling dazed or confused, experiencing loss of consciousness (LOC), or experiencing loss of memory for the injury or posttraumatic amnesia (PTA) for the event.”	NR	NR	Self-report screening tool
Romesser, Shen, Reblin, Kircher, Allen, Roberts, & Marchand, 2011 ⁴⁷	“History of military-related concussion during a VHA TBI secondary evaluation.” Excluded if “self-report suggested a history of a moderate or severe TBI (i.e., if they endorsed loss of consciousness more than 30 minutes).”	NR	NR	Clinical interview
Rona, Jones, Fear, et al., 2012 ⁴⁸	“Possible mTBI was assessed using a modified version of the BTBIS... A second item asked about possible symptoms associated with the injury. These were losing consciousness; being dazed or confused; not remembering the injury; concussion (e.g., headache, dizziness); head injury, and none of these. Participants were asked to tick all that applied. Self-report of the duration of any loss of consciousness was also obtained, we eliminated one participant from the analysis who reported prolonged loss of consciousness... Participants who endorsed at least one of these symptoms were classified as having mTBI.”	NR	Iverson, Langlois, McCrea, & Kelly, 2009	Self-report questionnaire

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Rona, Jones, Fear, et al., 2012 ⁴⁹	<p>“Possible mTBI was assessed using a modified version of the BTBIS, which included an item exploring possible causes of injury (blast, shrapnel fragments, bullet, fall, and vehicle accident and other). Participants could state that they had not suffered an injury during deployment. A second item asked about possible symptoms associated with the injury. These were losing consciousness; being dazed or confused; not remembering the injury; concussion (e.g., headache, dizziness); head injury; and none of these. Participants were asked to tick all that applied. Self-report of the duration of any loss of consciousness was also obtained, we eliminated 1 participant from the analysis who reported prolonged loss of consciousness (which would be classified traumatic brain injury, not mTBI). Participants who endorsed at least one of these symptoms were classified as having mTBI.”</p>	NR	Iverson, Langlois, McCrea, & Kelly, 2009	Self-report questionnaire
Schneiderman, Braver & Kang, 2008 ⁵⁰	Three-item Brief Traumatic Brain Injury Screen	NR	Defense and Veterans Brain Injury Center	Self-report survey
Skopp, Trofimovich, Grimes, et al., 2012 ⁵¹	<p>mild TBI (subset of all TBI codes): 310.2, 800.00-800.02, 800.06, 800.09, 800.50, 800.52, 801.00, 801.01,801.02, 801.06, 801.09, 801.50, 801.51, 801.52, 803.00-803.02, 803.06, 803.09, 803.50, 803.51, 803.52, 804.00, 804.01, 804.02, 804.06, 804.09, 804.50, 804.51, 804.52, 850.0, 805.1, 850.11, 850.9, 959.01, V15.52, V15.5_7, V15.5_C, V15.52_2, V15.52_2,V15.52_7, V15.52_C</p>	NR	DoD standard TBI surveillance case definition to ascertain TBI status and severity	Chart review

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Theeler, Flynn, & Erickson, 2010 ⁵²	“2-question screen followed by a 10-question screen... The first question asks if, while deployed, the soldier was exposed to or near a blast, improvised explosive device (IED) explosion, car bomb, suicide explosion, or exposed to any other combat event or resulted in a blow or jolt to the head. The second question asks if the soldier was involved in a motor vehicle accident, a fall, a sports accident, or any other event that caused a blow to the head or resulted in a neck whiplash. If the soldier answers yes to either question 1 or 2, 10 more questions asking about loss of consciousness, dazed sensation after the event, and neurologic or behavioral sequelae following the event are administered. This questionnaire is scored in a standardized manner from 0 to 39. The case definition for concussion constituted a score of 5 or greater on the 2-plus-10 questionnaire.”	NR	NR	Self-report survey
Theeler, Flynn, & Erickson, 2012 ⁵³	No specific criteria: 2-question followed by a 10-question screen if the soldier answers yes to either of the first 2 questions. This questionnaire is scored in a standardized manner from 0 to 39; the score from the 2-plus-10 questionnaire will herein be called the TBI Score	NR	Theeler, Flynn, & Erickson, 2010	Self-report survey
Vanderploeg, Belanger, & Curtiss, 2009 ⁵⁴	“Have you injured your head?... Did you lose consciousness as a result of the head injury?” “those individuals who required hospitalization after their head injury (n = 40) were excluded.” “head injury with altered consciousness” were classified as mild TBI.	NR	NR	Clinical interview
Vanderploeg, Curtiss & Belanger, 2005 ⁵⁵	“During the interview, participants were asked, among many others, the following three questions: 1) Since your discharge from active duty, have you been injured in a MVA? 2) Since your discharge from active duty have you injured your head (HI)? and 3) Did you lose consciousness as a result of the head injury?”	NR	NR	Self-report
Vanderploeg, Curtiss, Duchnick, Luis, 2003 ⁵⁶	Positive responses to following questions: “Since discharge from active duty, have you been injured in a MVA?”; “Since discharge from active duty, have you injured your head (from any cause)?”; and “Did you lose consciousness as a result of the head injury?”	NR	NR	Self-report survey

Author, Year	Definition	Patients with Abnormal Imaging	Citation	How Assessed
Vasterling, Brailey, Proctor, et al., 2012 ⁵⁷	“Congruent with reports showing stronger associations between clinical outcomes and TBI following loss of consciousness v. altered consciousness, only those pre- to post-deployment interval injuries resulting in loss of consciousness were queried.”	NR	NR	Self-report survey
Wilk, Herrell, Wynn, Riviere, & Hoge, 2012 ⁵⁸	“Injury resulted in being dazed, confused, or seeing stars, not remembering the injury, or losing consciousness (knocked out).”	NR	DoD/VA Brain Injury Center Brief Traumatic Brain Injury Screen	Self-report survey
Wilk, Thomas, McGurk, et al., 2010 ⁵⁹	“Injury resulted in being dazed, confused, or seeing stars, not remembering the injury, or losing consciousness (knocked out).”	NR	DoD/VA Brain Injury Center Brief Traumatic Brain Injury Screen	Self-report survey
Yurkiewicz, Lappan, Neely, et al., 2012 ⁶⁰	<p>“Defense and Veterans Brain Injury Center (DVBIC) assesses TBI severity after a blast event or motor vehicle accident, answers questions about evacuation and in-theater clinical care and medication, and provides recommendations on headache management and vision, hearing, vestibular, and neurologic issues. Successes of the program include a stratified headache protocol for primary care providers and an early management protocol for mild TBI and posttraumatic headache stratified headache protocol.”</p> <p>“We reviewed consults sent to the neurology group from October 2006 to December 2010 and consults sent to the TBI group from March 2008 to December 2010. Microsoft Excel spreadsheets were constructed to organize and analyze data regarding number of consultations, response times, location of origin, branch of service, clinical images transmitted, anatomic location of complaint, type of injury, workup recommended, and treatment and evacuation recommendations.”</p>	NR	DVBIC Brief Traumatic Brain Injury Screen	Chart review

Note. None of the studies reported information on assessor blinding of study hypotheses when assessing mTBI.

Bibliography of studies excluded for not meeting mTBI definition criteria

1. Adams RS, Larson MJ, Corrigan JD, Horgan CM, Williams TV. Frequent binge drinking after combat-acquired traumatic brain injury among active duty military personnel with a past year combat deployment. *J Head Trauma Rehabil.* Sep 2012;27(5):349-360.
2. Arbisi PA, Polusny MA, Erbes CR, Thurax P, Reddy MK. The Minnesota Multiphasic Personality Inventory-2 Restructured Form in National Guard soldiers screening positive for posttraumatic stress disorder and mild traumatic brain injury. *Psychol Assess.* Mar 2011;23(1):203-214.
3. Armistead-Jehle P. Symptom validity test performance in U.S. veterans referred for evaluation of mild TBI. *Appl Neuropsychol.* Jan 2010;17(1):52-59.
4. Bazarian JJ, Zhu T, Blyth B, Borrino A, Zhong J. Subject-specific changes in brain white matter on diffusion tensor imaging after sports-related concussion. *Magn Reson Imaging.* Feb 2012;30(2):171-180.
5. Booth-Kewley S, Highfill-McRoy RM, Larson GE, Garland CF, Gaskin TA. Anxiety and depression in marines sent to war in iraq and afghanistan. *J Nerv Ment Dis.* Sep 2012;200(9):749-757.
6. Brenner LA, Terrio H, Homaifar BY, et al. Neuropsychological test performance in soldiers with blast-related mild TBI. *Neuropsychology.* Mar 2010;24(2):160-167.
7. Brenner LA, Ivins BJ, Schwab K, et al. Traumatic brain injury, posttraumatic stress disorder, and postconcussive symptom reporting among troops returning from iraq. *J Head Trauma Rehabil.* Sep-Oct 2010;25(5):307-312.
8. Cameron KL, Marshall SW, Sturdivant RX, Lincoln AE. Trends in the incidence of physician-diagnosed mild traumatic brain injury among active duty U.S. military personnel between 1997 and 2007. *J Neurotrauma.* May 2012;29(7):1313-1321.
9. Carlson KF, Kehle SM, Meis LA, et al. Prevalence, assessment, and treatment of mild traumatic brain injury and posttraumatic stress disorder: a systematic review of the evidence. *J Head Trauma Rehabil.* Mar-Apr 2011;26(2):103-115.
10. Clement PF, Kennedy JE. Wechsler Adult Intelligence Scale-third edition characteristics of a military traumatic brain injury sample. *Mil Med.* Dec 2003;168(12):1025-1028.
11. Dougherty AL, MacGregor AJ, Han PP, Heltemes KJ, Galarneau MR. Visual dysfunction following blast-related traumatic brain injury from the battlefield. *Brain Inj.* 2011;25(1):8-13.
12. Drake AI, Gray N, Yoder S, Pramuka M, Llewellyn M. Factors predicting return to work following mild traumatic brain injury: A discriminant analysis. *The Journal of Head Trauma Rehabilitation.* Oct 2000;15(5):1103-1112.
13. Eskridge SL. Combat-related blast injuries: Injury types and outcomes. *Dissertation Abstracts International: Section B: The Sciences and Engineering.* 2012;72(7-B):3929.

14. Fear N, Jones E, Groom M, et al. Symptoms of post-concussional syndrome are nonspecifically related to mild traumatic brain injury in UK Armed Forces personnel on return from deployment in Iraq: An analysis of self-reported data. *Psychological Medicine*. Aug 2009;39(8):1379-1387.
15. Ferrier-Auerbach AG, Erbes CR, Polusny MA, Rath CM, Sponheim SR. Predictors of emotional distress reported by soldiers in the combat zone. *J Psychiatr Res*. May 2010;44(7):470-476.
16. French LM, Lange RT, Iverson GL, Ivins B, Marshall K, Schwab K. Influence of bodily injuries on symptom reporting following uncomplicated mild traumatic brain injury in US military service members. *J Head Trauma Rehabil*. Jan-Feb 2012;27(1):63-74.
17. Gottshall KR, Gray NL, Drake AI, Tejidor R, Hoffer ME, McDonald EC. To investigate the influence of acute vestibular impairment following mild traumatic brain injury on subsequent ability to remain on activity duty 12 months later. *Mil Med*. Aug 2007;172(8):852-857.
18. Helfer TM, Jordan NN, Lee RB, Pietrusiak P, Cave K, Schairer K. Noise-induced hearing injury and comorbidities among postdeployment U.S. Army soldiers: April 2003-June 2009. *Am J Audiol*. Jun 2011;20(1):33-41.
19. Heltemes KJ, Dougherty AL, MacGregor AJ, Galarneau MR. Alcohol abuse disorders among U.S. service members with mild traumatic brain injury. *Mil Med*. Feb 2011;176(2):147-150.
20. Hoffer ME, Balaban C, Gottshall K, Balough BJ, Maddox MR, Penta JR. Blast exposure: vestibular consequences and associated characteristics. *Otol Neurotol*. Feb 2010;31(2):232-236.
21. Hoffer ME, Donaldson C, Gottshall KR, Balaban C, Balough BJ. Blunt and blast head trauma: different entities. *Int Tinnitus J*. 2009;15(2):115-118.
22. Hoge CW, McGurk D, Thomas JL, Cox AL, Engel CC, Castro CA. Mild traumatic brain injury in U.S. Soldiers returning from Iraq. *N Engl J Med*. Jan 31 2008;358(5):453-463.
23. Ivins BJ, Schwab KA, Warden D, et al. Traumatic brain injury in U.S. Army paratroopers: prevalence and character. *J Trauma*. Oct 2003;55(4):617-621.
24. Ivins BJ, Schwab KA, Baker G, Warden DL. Hospital admissions associated with traumatic brain injury in the US Army during peacetime: 1990s trends. *Neuroepidemiology*. 2006;27(3):154-163.
25. Ivins BJ, Kane R, Schwab KA. Performance on the Automated Neuropsychological Assessment Metrics in a nonclinical sample of soldiers screened for mild TBI after returning from Iraq and Afghanistan: a descriptive analysis. *J Head Trauma Rehabil*. Jan-Feb 2009;24(1):24-31.
26. Ivins BJ. Hospitalization associated with traumatic brain injury in the active duty US Army: 2000-2006. *NeuroRehabilitation*. 2010;26(3):199-212.

27. Lange RT, Pancholi S, Bhagwat A, Anderson-Barnes V, French LM. Influence of poor effort on neuropsychological test performance in U.S. military personnel following mild traumatic brain injury. *J Clin Exp Neuropsychol*. 2012;34(5):453-466.
28. Lange RT, Pancholi S, Brickell TA, et al. Neuropsychological Outcome from Blast versus Non-blast: Mild Traumatic Brain Injury in U.S. Military Service Members. *J Int Neuropsychol Soc*. May 2012;18(3):595-605.
29. Lange RT, Brickell TA, French LM, et al. Neuropsychological Outcome from Uncomplicated Mild, Complicated Mild, and Moderate Traumatic Brain Injury in US Military Personnel. *Arch Clin Neuropsychol*. 2012;27(5):480-494.
30. Lew HL, Garvert DW, Pogoda TK, et al. Auditory and visual impairments in patients with blast-related traumatic brain injury: Effect of dual sensory impairment on Functional Independence Measure. *J Rehabil Res Dev*. 2009;46(6):819-826.
31. Luis CA, Vanderploeg RD, Curtiss G. Predictors of postconcussion symptom complex in community dwelling male veterans. *J Int Neuropsychol Soc*. Nov 2003;9(7):1001-1015.
32. Macgregor AJ. Physical injury and psychological outcomes among United States combat veterans. *Dissertation Abstracts International: Section B: The Sciences and Engineering*. 2007;68(6-B):3665.
33. MacGregor AJ, Dougherty AL, Galarneau MR. Injury-specific correlates of combat-related traumatic brain injury in Operation Iraqi Freedom. *The Journal of Head Trauma Rehabilitation*. Jul-Aug 2011;26(4):312-318.
34. MacGregor AJ, Dougherty AL, Morrison RH, Quinn KH, Galarneau MR. Repeated concussion among U.S. military personnel during Operation Iraqi Freedom. *J Rehabil Res Dev*. 2011;48(10):1269-1278.
35. MacGregor AJ, Shaffer RA, Dougherty AL, et al. Prevalence and psychological correlates of traumatic brain injury in operation iraqi freedom. *J Head Trauma Rehabil*. Jan-Feb 2010;25(1):1-8.
36. McGuire SA, Marsh RW, Sowin TW, Robinson AY. Aeromedical decision making and seizure risk after traumatic brain injury: longitudinal outcome. *Aviat Space Environ Med*. Feb 2012;83(2):140-143.
37. Mora AG, Ritenour AE, Wade CE, Holcomb JB, Blackburne LH, Gaylord KM. Posttraumatic stress disorder in combat casualties with burns sustaining primary blast and concussive injuries. *J Trauma*. Apr 2009;66(4 Suppl):S178-185.
38. Morgan M, Lockwood A, Steinke D, Schleenbaker R, Botts S. Pharmacotherapy regimens among patients with posttraumatic stress disorder and mild traumatic brain injury. *Psychiatric Services*. Feb 2012;63(2):182-185.
39. Nelson C, St, Weiser M, Gifford S, Gallimore J, Morningstar A. Knowledge gained from the Traumatic Brain Injury Screen-Implications for treating Canadian military personnel. *Mil Med*. Feb 2011;176(2):156-160.

40. Olson-Madden JH, Forster JE, Huggins J, Schneider A. Psychiatric diagnoses, mental health utilization, high-risk behaviors, and self-directed violence among veterans with comorbid history of traumatic brain injury and substance use disorders. *J Head Trauma Rehabil.* Sep 2012;27(5):370-378.
41. Ommaya AK, Ommaya AK, Dannenberg AL, Salazar AM. Causation, incidence, and costs of traumatic brain injury in the U.S. military medical system. *J Trauma.* Feb 1996;40(2):211-217.
42. Ommaya AK, Salazar AM, Dannenberg AL, Chervinsky AB, Schwab K. Outcome after traumatic brain injury in the U.S. military medical system. *J Trauma.* Dec 1996;41(6):972-975.
43. Pietrzak RH, Johnson DC, Goldstein MB, Malley JC, Southwick SM. Posttraumatic stress disorder mediates the relationship between mild traumatic brain injury and health and psychosocial functioning in veterans of Operations Enduring Freedom and Iraqi Freedom. *J Nerv Ment Dis.* Oct 2009;197(10):748-753.
44. Plassman BL, Havlik RJ, Steffens DC, et al. Documented head injury in early adulthood and risk of Alzheimer's disease and other dementias. *Neurology.* Oct 24 2000;55(8):1158-1166.
45. Polusny MA, Kehle SM, Nelson NW, Erbes CR, Arbisi PA, Thuras P. Longitudinal effects of mild traumatic brain injury and posttraumatic stress disorder comorbidity on postdeployment outcomes in national guard soldiers deployed to Iraq. *Arch Gen Psychiatry.* Jan 2011;68(1):79-89.
46. Roebuck-Spencer TM, Vincent AS, Twillie DA, et al. Cognitive change associated with self-reported mild traumatic brain injury sustained during the OEF/OIF conflicts. *The Clinical Neuropsychologist.* Apr 2012;26(3):473-489.
47. Romesser J, Shen S, Reblin M, et al. A preliminary study of the effect of a diagnosis of concussion on PTSD symptoms and other psychiatric variables at the time of treatment seeking among veterans. *Mil Med.* Mar 2011;176(3):246-252.
48. Rona RJ, Jones M, Fear NT, et al. Mild traumatic brain injury in UK military personnel returning from Afghanistan and Iraq: cohort and cross-sectional analyses. *J Head Trauma Rehabil.* Jan-Feb 2012;27(1):33-44.
49. Rona RJ, Jones M, Fear NT, Sundin J, Hull L, Wessely S. Frequency of mild traumatic brain injury in Iraq and Afghanistan: are we measuring incidence or prevalence? *J Head Trauma Rehabil.* Jan-Feb 2012;27(1):75-82.
50. Schneiderman AI, Braver ER, Kang HK. Understanding sequelae of injury mechanisms and mild traumatic brain injury incurred during the conflicts in Iraq and Afghanistan: persistent postconcussive symptoms and posttraumatic stress disorder. *Am J Epidemiol.* Jun 15 2008;167(12):1446-1452.
51. Skopp NA, Trofimovich L, Grimes J, Oetjen-Gerdes L, Gahm GA. Relations between suicide and traumatic brain injury, psychiatric diagnoses, and relationship problems, active component, U.S. Armed Forces, 2001-2009. *Msmr.* Feb 2012;19(2):7-11.

52. Theeler BJ, Flynn FG, Erickson JC. Headaches after concussion in US soldiers returning from Iraq or Afghanistan. *Headache*. Sep 2010;50(8):1262-1272.
53. Theeler BJ, Flynn FG, Erickson JC. Chronic Daily Headache in U.S. Soldiers After Concussion. *Headache*. 2012;52(5):732-738.
54. Vanderploeg RD, Belanger HG, Curtiss G. Mild traumatic brain injury and posttraumatic stress disorder and their associations with health symptoms. *Arch Phys Med Rehabil*. Jul 2009;90(7):1084-1093.
55. Vanderploeg RD, Curtiss G, Belanger HG. Long-term neuropsychological outcomes following mild traumatic brain injury. *J Int Neuropsychol Soc*. May 2005;11(3):228-236.
56. Vanderploeg RD, Curtiss G, Duchnick JJ, Luis CA. Demographic, medical, and psychiatric factors in work and marital status after mild head injury. *J Head Trauma Rehabil*. Mar-Apr 2003;18(2):148-163.
57. Vasterling JJ, Brailey K, Proctor SP, Kane R, Heeren T, Franz M. Neuropsychological outcomes of mild traumatic brain injury, post-traumatic stress disorder and depression in Iraq-deployed US Army soldiers. *Br J Psychiatry*. Sep 2012;201:186-192.
58. Wilk JE, Herrell RK, Wynn GH, Riviere LA, Hoge CW. Mild traumatic brain injury (concussion), posttraumatic stress disorder, and depression in U.S. soldiers involved in combat deployments: association with postdeployment symptoms. *Psychosom Med*. Apr 2012;74(3):249-257.
59. Wilk JE, Thomas JL, McGurk DM, Riviere LA, Castro CA, Hoge CW. Mild traumatic brain injury (concussion) during combat: lack of association of blast mechanism with persistent postconcussive symptoms. *J Head Trauma Rehabil*. Jan-Feb 2010;25(1):9-14.
60. Yurkiewicz IR, Lappan CM, Neely ET, et al. Outcomes from a US military neurology and traumatic brain injury telemedicine program. *Neurology*. Sep 18 2012;79(12):1237-1243.

APPENDIX E. RESULTS OF INCLUDED STUDIES

Cognitive Functioning

Table 1a. Language abilities and general fund of verbal knowledge measures in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	WTAR scaled score	---	106.7 (10.1)	105.4 (5.7)	“NS”	NR
		WAIS-III	Information scaled score	12.8 (1.7)	11.8 (2.1)	“NS”	NR
	Forensic context	WAIS-III	Information scaled score	11.97 (1.92)	11.58 (1.82)	“NS”	NR
Cooper, Chau, Armistead-Jehle, et al., 2012 ¹⁶	Blast exposed	RBANS	Language	97.43 (12.42)	92.81 (14.10)	p = 0.187	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	LOC and/or PTA	Shipley Institute of Living Scale	Vocabulary Subtest	49.15 (7.50)	49.15 (6.20)	NR	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	Mental Health Diagnosis other than PTSD	WTAR	---	96.8 (10.8)	101.7 (12.3)	NR	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	PTSD Diagnosis	WTAR	---	96.8 (10.8)	99.0 (11.2)	NR	NR
Belanger, Kretzmer, Venderploeg, & French, 2010 ¹⁰	Same population with moderate/severe TBI	WTAR	Estimated Full Scale IQ (FSIQ)	98.3 (9.3)	97.6 (9.1)	p = .67	NR
Belanger, Kretzmer, Yoash-Gantz, Pickett, & Tupler, 2009 ¹¹	Same population with moderate/severe TBI	WTAR	Estimated Full Scale IQ (FSIQ)	98.1 (14.6)	96.1 (12.8)	p = .35	NR
Cooper, Mercado-Couch, Critchfield, et al., 2010 ¹⁸	Same population without mTBI	RBANS	Language	92.90 (15.586)	93.66 (11.873)	p = 0.732	NR
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Same population without mTBI; none with poor effort	WAIS-III	Information scaled score	12.21 (2.01)	12.26 (2.25)	“NS”	d = .22
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Same population without mTBI; none have Axis I	WAIS-III	Information scaled score	12.8 (1.7)	11.9 (2.6)	“NS”	NR
		WTAR	scaled score	106.7 (10.1)	106.0 (9.2)	“NS”	NR

Table 1b. Visuospatial function in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect	
Spencer, et al., 2010 ³⁶	NA	RCFT	Figure Copy	33.6 (2.9)	NA	NA	NA	
			Figure Orientation	3.6 (1.7)	NA	NA	NA	
			Time to Copy	158.7 (63.5)	NA	NA	NA	
Cooper, Mercado-Couch, Critchfield, et al., 2010 ¹⁸	Same population without mTBI	RBANS	Visuospatial/Constructional	104.06 (13.382)	109.29 (10.47)	p = 0.007	NR	
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Same population without mTBI; none have Axis I	RCFT	Figure Copy	31.3 (3.2)	32.0 (2.1)	“NS”	NR	
			WAIS-III	Block Design scaled score	12.2 (2.6)	13.0 (2.9)	“NS”	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	RCFT	Figure Copy	31.3 (3.2)	31.5 (3.0)	“NS”	NR	
			WAIS-III	Block Design scaled score	12.2 (2.6)	12.2 (3.1)	“NS”	NR
Cooper, Chau, Armistead-Jehle, et al., 2012 ¹⁶	Blast exposed	RBANS	Visuospatial/Constructional	112.11 (9.70)	108.09 (11.80)	p = 0.159	NR	
Drag, Spencer, Walker, et al., 2012 ²⁰	HADS Anxiety	Visual Organization/Processing Factor	---	NR	NR	p > .05	r = -.12	
	HADS Depression	Visual Organization/Processing Factor	---	NR	NR	p < .001	r = -.25	
	LOC and/or PTA	Visual Organization/Processing Factor	RCFT	Figure Copy	33.72 (2.64)	33.78 (2.38)	NR	NR
			Organization	3.64 (1.74)	3.99 (1.81)	NR	NR	
			Time to Copy	170.51 (63.07)	166.17 (74.91)	NR	NR	
			---	49.08 (9.98)	50.46 (10.12)	p = .30	NR	
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	Mental Health Diagnosis other than PTSD	RCFT	Figure Copy	32.6 (4.9)	32.7 (4.7)	NR	NR	
Drag, Spencer, Walker, et al., 2012 ²⁰	PCL-M score	Visual Organization/Processing Factor	---	NR	NR	p < .05	r = -.18	
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	PTSD Diagnosis	RCFT	Figure Copy	32.6 (4.9)	32.1 (6.0)	NR	NR	
Spencer et al., 2010 ³⁶	Self-reported slowed thinking/organization	RCFT	Figure Copy	NR	NR	“NS”	r = -.13	
			Figure Orientation	NR	NR	“NS”	r = -.02	
			Time to Copy	NR	NR	“NS”	r = -.14	
Drag, Spencer, Walker, et al., 2012 ²⁰	Service connected	Visual Organization/Processing Factor	---	NR	NR	p = .57	NR	

Table 1c. Memory functioning in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	CVLT-II	Long Delay Free Recall z-score	0.5 (0.9)	-0.1 (1.1)	“Significant”	NR
			Trials 1-5 z-score	0.6 (0.9)	0.1 (1.0)	“NS”	NR
			RCFT	Delayed Recall z-score	-0.1 (1.3)	-0.5 (.98)	“NS”
Belanger, Kretzmer, Yoash-Gantz, Pickett, & Tupler, 2009 ¹¹	Blast exposure	BVM-T-R	Delayed Recall t score	48.6 (12.8)	50.7 (10.7)	“NS”	NR
			Total Recall t score	45.2 (12.3)	50.1 (10.1)	“NS”	NR
		CVLT-II	Long Delay Free Recall t-score	48.3 (9.1)	50.5 (11.4)	p = .38	NR
			Total Trials 1-5 t-score	52.8 (10.5)	54.0 (8.1)	p = .38	NR
Cooper, Chau, Armistead-Jehle, et al., 2012 ¹⁶	Blast exposure	RBANS	Immediate Memory	91.70 (11.0)	91.81 (15.67)	P = 0.994	NR
			Delayed Memory	96.96 (16.59)	95.12 (16.04)	P = 0.664	NR
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Forensic context	CVLT-II	Long Delay Free Recall z-score	0.03 (0.99)	-0.73 (1.12)	“NS”	NR
			Trials 1-5 t-score	52.82 (8.61)	46.33 (9.23)	“NS”	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	HADS Anxiety	Memory Factor	---	NR	NR	p < .001	r = -.29
Drag, Spencer, Walker, et al., 2012 ²⁰	HADS Depression	Memory Factor	---	NR	NR	p > .05	r = -.15
Drag, Spencer, Walker, et al., 2012 ²⁰	LOC and/or PTA	Memory Factor	---	49.22 (9.89)	49.61 (10.36)	NR	NR
			RBANS	Story Memory Immediate Recall	44.99 (10.19)	48.10 (10.36)	NR
		RCFT	Story Memory Delayed Recall	45.41 (10.49)	44.98 (10.71)	NR	NR
			Immediate Recall	44.43 (12.72)	44.50 (13.65)	NR	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	Mental Health Diagnosis other than PTSD	CVLT-II	Long Delay Free Recall	8.9 (3.7)	8.9 (3.9)	NR	NR
			Short Delay Free Recall	8.3 (3.5)	8.6 (3.9)	NR	NR
		CVLT-II	Total Trials 1-5	43.7 (12.0)	44.6 (13.7)	NR	NR
		RCFT	Immediate Recall	16.4 (7.7)	17.4 (7.2)	NR	NR
			Delayed Recall	16.0 (7.9)	17.4 (7.2)	NR	NR

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect			
Spencer, et al., 2010 ³⁶	No comparison group	RBANS	Story Memory Immediate Recall	17.2 (3.4)	NA	NA	NA			
			Story Memory Delayed Recall	8.4 (2.3)	NA	NA	NA			
		RCFT	Immediate Recall	19.9 (6.2)	NA	NA	NA			
			MSP subtest	Baseline: 51 (43-53) ≤ 72 hours: 44 (38.5-51)	Baseline: 50 (45-55) ≤ 72 hours: 50 (44-56)	Baseline: p = .57 ≤ 72 hours: p <.001	NR			
Drag, Spencer, Walker, et al., 2012 ²⁰	PCL-M score	Memory Factor	---	NR	NR	p < .05	r = -.34			
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	PTSD Diagnosis	CVLT-II	Long Delay Free Recall	8.9 (3.7)	7.6 (3.2)	NR	NR			
			Short Delay Free Recall	8.3 (3.5)	7.0 (3.4)	NR	NR			
			Total Trials 1-5	43.7 (12.0)	38.7 (11.1)	NR	NR			
		RCFT	Immediate Recall	16.4 (7.7)	15.7 (8.5)	NR	NR			
			Delayed Recall	16.0 (7.9)	16.1 (7.1)	NR	NR			
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Same population of research participants, none with poor effort	CVLT-II	Long Delay Free Recall z-score	0.27 (0.91)	0.45 (0.81)	"NS"	d = 0.21			
			Trials 1-5 t-score	54.43 (8.95)	57.16 (8.04)	"NS"	d = 0.32			
Belanger, Kretzmer, Yoash-Gantz, Pickett, & Tupler, 2009 ¹¹	Same population with moderate/ severe TBI	BVMT-R	Delayed Recall	NR	NR	"NS"	NR			
		BVMT-R	Total Recall	NR	NR	"NS"	NR			
		CVLT-II	Long Delay Free Recall	NR	NR	p < .05	NR			
			Total Trials 1-5	NR	NR	p < .01	NR			
Schiehser, Delis, Filoteo, et al., 2011 ³⁵	Same population with moderate/ severe TBI	Memory Composite Score	---	9.0 (2.4)	7.7 (2.8)	p = .04	NR			
Coldren, Russell, Parish, et al., 2012 ¹⁵	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	ANAM	CDD subtest	Baseline: 42.5 (36.5-50) 5+ days: 49.5 (40-56) 10+ days: 49.5 (40.1-53.1)	Baseline: 44 (37-52) 5+ days: 50.3 (44.1-57.8) 10+ days: 50.3 (44.1-57.8)	Baseline: p = 0.47 5+ days: p = 0.07 10+ days: p = 0.17	NR			
				MSP subtest	Baseline: 51 (43-53) 5+ days: 50.5 (44-58) 10+ days: 51.8 (43.5-57.9)	Baseline: 52 (44.5-57.8) 5+ days: 51.5 (44.5-57.6) 10+ days: 51.5 (44.5- 57.6)	Baseline: p = 0.29 5+ days: p = 0.47 10+ days: p = 0.77	NR		
			Cooper, Mercado-Couch, Critchfield, et al., 2010 ¹⁸		Same population without TBI	RBANS	Immediate Memory	95.14 (14.181)	96.49 (14.445)	p = .589

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect
Kelly, Coldren, Parish, et al., 2012 ²⁴	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	ANAM	CDD substest	Baseline: 42.5 (35-50) ≤ 72 hours: 42 (35-49.5)	Baseline: 45 (38-52) ≤ 72 hours: 44.75 (39-51)	Baseline: p = .20 ≤ 72 hours: p = .04	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Same population without mTBI; none have Axis I	CVLT-II	Long Delay Free Recall z-score	0.5 (0.9)	0.6 (0.8)	"NS"	NR
			Trials 1-5 z-score	0.6 (0.9)	0.8 (0.8)	"NS"	NR
		RCFT	Delayed Recall z-score	-0.1 (1.3)	-0.3 (1.3)	"NS"	NR
			Delayed Memory	96.48 (12.998)	100.42 (12.854)	p = 0.072	NR
Spencer et al., 2010 ³⁶	Self-reported memory deficits	RBANS	Story Memory Immediate Recall	NR	NR	"NS"	r = -.05
			Story Memory Delayed Recall	NR	NR	p < .05	r = -.20
		RCFT	Immediate Recall	NR	NR	"NS"	r = .08
Drag, Spencer, Walker, et al., 2012 ²⁰	Service connected	Memory Factor	---	NR	NR	p = .17	NR

Table 1d. Attention/concentration measures in studies of U.S. Veterans and members of the U.S. military with mTBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect	
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	WAIS-III	Digit Span scaled score	9.9 (2.8)	9.9 (2.1)	“NS”	NR	
	Forensic context	WAIS-III	Digit Span scaled score	9.61 (2.55)	8.42 (2.77)	“NS”	NR	
Cooper, Chau, Armistead-Jehle, et al., 2012 ¹⁶	Blast exposure	RBANS	Attention	98.89 (16.64)	94.78 (14.51)	p = 0.311	NA	
Drag, Spencer, Walker, et al., 2012 ²⁰	HADS Anxiety	Verbal Attention Factor	---	NR	NR	p < .05	r = -.24	
		Visual Attention Factor	---	NR	NR	p > .05	r = -.12	
	HADS Depression	Verbal Attention Factor	---	NR	NR	p < .05	r = -.24	
		Visual Attention Factor	---	NR	NR	p > .05	r = -.13	
	LOC and/or PTA	Verbal Attention Factor	---	49.69 (10.91)	49.87 (9.69)	p = .91	NR	
		Visual Attention Factor	---	50.67 (9.61)	49.61 (10.36)	p = .50	NR	
	LOC and/or PTA	WAIS-IV	Digit Sequencing	---	9.30 (2.76)	9.41 (2.27)	NR	NR
			Digit Span Backward	---	9.23 (2.54)	8.92 (2.18)	NR	NR
Digits Forward			---	8.39 (2.63)	8.64 (2.70)	NR	NR	
Spencer et al., 2010 ³⁶	No comparison group	WAIS-IV	Digit Sequencing	---	8.0 (1.9)	NA	NA	NA
			Digits Backward	---	7.8 (2.1)	NA	NA	NA
			Digits Forward	---	9.6 (2.1)	NA	NA	NA
Drag, Spencer, Walker, et al., 2012 ²⁰	PCL-M score	Verbal Attention Factor	---	NR	NR	p < .05	r = -.21	
		Visual Attention Factor	---	NR	NR	p < .05	r = -.20	
		Digit Span scaled score	---	9.9 (2.8)	10.9 (2.4)	“NS”	NR	
Coldren, Russell, Parish, et al., 2012 ¹⁵	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	ANAM	Mathematical Processing (MTH) subtest	Baseline:	Baseline:	Baseline:	NR	
				51 (42-57.3)	50 (42-55)	p = 0.71		
				5+ days:	5+ days:	5+ days:		
50.5 (43-56)	52 (44-59.8)	p = 0.29						
10+ days:	10+ days:	10+ days:						
50.8 (46.3-55.8)	52 (44-59.8)	p = 0.51						
Cooper, Mercado-Couch, Critchfield, et al., 2010 ¹⁸	Same population without mTBI	RBANS	Attention	84.06 (15.013)	89.74 (14.898)	p = 0.026	NR	

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect
Kelly, Coldren, Parish, et al., 2012 ²⁴	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	ANAM	Mathematical Processing (MTH) subtest	Baseline: 47 (42-57)	Baseline: 52 (42-55)	Baseline: p = .57	NR
				≤ 72 hours: 46.7 (40.5-52.5)	≤ 72 hours: 50 (44-56.6)	≤ 72 hours: p = .03	
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Same population without mTBI; none with poor effort	WAIS-III	Digit Span scaled score	10.50 (2.29)	10.65 (2.37)	“NS”	d = .06
Spencer, et al., 2010 ³⁶	Self-reported attention deficits	WAIS-IV	Digit Sequencing	NR	NR	“NS”	r = -.15
			Digits Backward	NR	NR	“NS”	r = -.11
			Digits Forward	NR	NR	“NS”	r = -.15
			Digit Sequencing	NR	NR	“NS”	r = .00
			Digits Backward	NR	NR	“NS”	r = -.14
Drag, Spencer, Walker, et al., 2012 ²⁰	Service connected	Verbal Attention Factor	---	NR	NR	p = .42	NR
		Visual Attention Factor	---	NR	NR	p = .17	NR

Table 1e. Cognitive processing speed in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Stroop Color and Word Test	Color t-score	49.5 (6.9)	45.8 (7.8)	“NS”	NR
			Word t-score	49.4 (9.8)	46.2 (8.6)	“NS”	NR
		Trail Making Test	Part A t-score	50.0 (11.7)	50.5 (9.7)	“NS”	NR
			WAIS-III	Digit Symbol Coding scaled score	10.9 (2.0)	9.8 (2.4)	“NS”
Belanger, Kretzmer, Yoash-Gantz, Pickett, & Tupler, 2009 ¹¹	Blast exposure	Trail Making Test	Part A t-score	45.8 (14.8)	46.2 (11.6)	all p values > .10	NR
		WAIS-III	Digit Symbol Coding scaled score	8.3 (2.7)	8.9 (2.5)	all p values > .10	NR

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Forensic context	Stroop Color and Word Test	Color t-score	45.29 (8.60)	40.04 (7.00)	“NS”	NR
			Word t-score	45.47 (9.01)	39.12 (7.54)	“NS”	NR
		Trail Making Test	Part A t-score	48.71 (10.44)	39.96 (9.78)	“Significant”	NR
		WAIS-III	Digit Symbol Coding scaled score	9.92 (2.53)	7.12 (2.53)	“Significant”	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	LOC and/or PTA	Trail Making Test	Part A	48.93 (10.83)	47.99 (11.04)	NR	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	Mental Health Diagnosis other than PTSD	Stroop Color & Word Test	Color	58.8 (12.1)	54.1 (11.2)	NR	NR
			Word	76.2 (16.5)	77.0 (17.7)	NR	NR
		Trail Making Test	Part A	41.8 (15.8)	41.9 (19.1)	NR	NR
Spencer et al., 2010 ³⁶	No comparison group	Trail Making Test	Part A	29.6 (13.1)	NA	NA	NA
			Procedural Reaction Time (PRT) subtest	Baseline: 52 (43-53) ≤ 72 hours: 46 (34.5-54.5)	Baseline: 50 (47-58) ≤ 72 hours: 52.5 (46.5-59)	Baseline: p = .60 ≤ 72 hours: p < .001	NR
			Simple Reaction Time (SRT) subtest	Baseline: 53 (48-55) ≤ 72 hours: 44.5 (34.5-52)	Baseline: 52 (47-55) ≤ 72 hours: 52 (47-55.5)	Baseline: p = .24 ≤ 72 hours: p < .001	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	PTSD Diagnosis	Stroop Color & Word Test	Color	58.8 (12.1)	60.8 (15.3)	NR	NR
			Word	76.2 (16.5)	79.6 (18.2)	NR	NR
		Trail Making Test	Part A	41.8 (15.8)	40.1 (15.0)	NR	NR
Coldren, Russell, Parish, et al., 2012 ¹⁵	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	ANAM	Code Substitution (CDS) subtest	Baseline: 46 (39.5-48.5) 5+ days: 52.5 (46.5-58) 10+ days: 53 (47-60)	Baseline: 48 (43-55) 5+ days: 57 (50-63) 10+ days: 57 (50-63)	Baseline: p = 0.04 5+ days: p = 0.03 10+ days: p = 0.14	NR
			Procedural Reaction Time (PRT) subtest	Baseline: 52 (43-57) 5+ days: 52 (44.5-58.5) 10+ days: 48.8 (42-62.1)	Baseline: 52 (47-58) 5+ days: 54 (47-62.5) 10+ days: 54 (47-62.5)	Baseline: p = 0.77 5+ days: p = 0.13 10+ days: p = 0.20	NR
			Simple Reaction Time (SRT) subtest	Baseline: 53 (52-55) 5+ days: 54 (48-56.5) 10+ days: 54.5 (46.6-57.6)	Baseline: 52 (47-58) 5+ days: 53.5 (48.8-56.5) 10+ days: 53.5 (48.8-56.5)	Baseline: p = .23 5+ days: p = .97 10+ days: p = .71	NR
Kelly, Coldren, Parish, et al., 2012 ²⁴	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	ANAM	Code Substitution (CDS) subtest	Baseline: 44 (37-50) ≤ 72 hours: 44 (38.5-51)	Baseline: 48 (43-55) ≤ 72 hours: 52 (45-57)	Baseline: p = .02 ≤ 72 hours: p < .001	NR

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or Median (IQR)	Comparison group Mean (SD) or Median (IQR)	p value for comparison	Magnitude of effect
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Same population without mTBI; none with poor effort	Stroop Color and Word Test	Color t-score	46.00 (8.88)	47.26 (6.90)	"NS"	d = 0.16
			Word t-score	46.29 (9.10)	48.58 (8.64)	"NS"	d = 0.26
		Trail Making Test	Part A t-score	48.89 (9.89)	49.61 (11.66)	"NS"	d = 0.10
		WAIS-III	Digit Symbol Coding scaled score	10.46 (2.52)	10.35 (2.47)	"NS"	d = 0.04
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Same population without mTBI; none have Axis I	Stroop Color and Word Test	Color t-score	49.5 (6.9)	48.9 (6.7)	"NS"	NR
Swick, Honzel, Larsen, et al., 2012 ³⁷	Same population without mTBI; participants had PTSD	Reaction time on a Go/NoGo task	---	NR	NR	p > .7	NR
Belanger, Kretzmer, Yoash-Gantz, Pickett, & Tupler, 2009 ¹¹	Same population with moderate/severe TBI	Trail Making Test	Part A	NR	NR	all p values > .10	NR
		WAIS-III	Digit Symbol Coding	NR	NR		all p values > .10
			Word t-score	49.4 (9.8)	48.3 (6.6)	"NS"	NR
		Trail Making Test	Part A t-score	50.0 (11.7)	51.8 (12.1)	"NS"	NR
		WAIS-III	Digit Symbol Coding scaled score	10.9 (2.0)	11.1 (2.7)	"NS"	NR
Spencer, et al., 2010 ³⁶	Self-reported attention deficits	Trail Making Test	Part A	NR	NR	"NS"	r = -.03
Spencer, et al., 2010 ³⁶	Self-reported slowed thinking/organization	Trail Making Test	Part A	NR	NR	"NS"	r = -.09

Table 1f. Executive functioning in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Belanger, Kretzmer, Yoash-Gantz, Pickett, & Tupler, 2009 ¹¹	Blast exposure	Trail Making Test	Part B t-score	49.1 (15.0)	45.6 (9.4)	all p values > .10	NR
	Same population with moderate/severe TBI	Trail Making Test	Part B	NR	NR	all p values > .10	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	LOC and/or PTA	Trail Making Test	Part B t-score	49.96 (8.35)	48.80 (8.65)	NR	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	Mental Health Diagnosis other than PTSD	Stroop Color & Word Test	Color Word	32.9 (7.9)	30.3 (9.0)	NR	NR
		Trail Making Test	Part B	93.0 (38.8)	106.1 (73.2)	NR	NR
	PTSD Diagnosis	Stroop Color & Word Test	Color Word	32.9 (7.9)	30.5 (8.5)	NR	NR
		Trail Making Test	Part B	93.0 (38.8)	95.2 (34.5)	NR	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	COWA t-score	---	46.4 (11.0)	42.7 (9.5)	“NS”	NR
		Stroop Color and Word Test	Color-Word t-score	51.1 (9.5)	48.3 (9.5)	“NS”	NR
		Trail Making Test	Part B t-score	51.0 (12.1)	48.4 (10.8)	“NS”	NR
	Same population without mTBI; none have Axis I	COWA t-score	---	46.4 (11.0)	48.9 (9.7)	“NS”	NR
		Stroop Color and Word Test	Color-Word t-score	51.1 (9.5)	52.6 (7.0)	“NS”	NR
		Trail Making Test	Part B t-score	51.0 (12.1)	53.6 (7.5)	“NS”	NR
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Forensic context	COWA t-score	---	44.79 (10.43)	42.83 (8.52)	“NS”	NR
		Stroop Color and Word Test	Color-Word t-score	46.95 (9.09)	43.78 (8.10)	“NS”	NR
		Trail Making Test	Part B t-score	51.45 (10.25)	44.71 (10.22)	“NS”	NR
	Same population without mTBI; none with poor effort	COWA t-score	---	47.04 (8.94)	46.97 (9.69)	“NS”	d = 0.01
		Stroop Color and Word Test	Color-Word t-score	48.86 (9.35)	50.52 (9.22)	“NS”	d = 0.18
		Trail Making Test	Part B t-score	52.21 (10.20)	52.77 (5.83)	NS	d = 0.07
Schiehser, Delis, Filoteo, et al., 2011 ³⁵	Same population with moderate/severe TBI	Executive Function Composite Score	---	10.4 (2.2)	9.5 (3.0)	p = .13	NR
Spencer, et al., 2010 ³⁶	None	Trail Making Test	Part B	72.8 (34.8)	NA	NA	NA
	Self-reported attention deficits	Trail Making Test	Part B	NR	NR	“NS”	r = -.01
	Self-reported slowed thinking/organization	Trail Making Test	Part B	NR	NR	“NS”	r = -.01

Table 1g. Effort and motivation measures in a study of U.S. Veterans with mild TBI evaluated in a research versus forensic context

Author, year	Outcome measure	Portion or subscale	Research context Mean (SD)	Forensic context Mean (SD)	p value for comparison	Magnitude of effect	
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	CVLT-II	Forced Choice	15.87 (0.41)	15.50 (1.18)	“NS”	NR	
	Effort Failures raw score	---	0.32 (0.58)	1.17 (0.87)	“Significant”	NR	
	Rey FIT	Combination	28.84 (2.06)	26.13 (3.72)	“Significant”	NR	
	VSVT	Easy Items		23.90 (0.31)	22.92 (2.00)	“Significant”	NR
		Difficult Items		22.11 (2.87)	15.63 (6.25)	“Significant”	NR
		Total Items		46.00 (3.08)	38.54 (7.43)	“Significant”	NR
	WAIS-III	Reliable Digit Span		9.63 (1.98)	8.38 (1.71)	“NS”	NR

Table 1h. Total and cross-domain composite scores in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Cooper, Chau, Armistead-Jehle, et al., 2012 ¹⁶	Blast exposure	RBANS	Total Score	98.61 (9.33)	94.88 (12.92)	p = 0.211	NR
Cooper, Mercado-Couch, Critchfield, et al., 2010 ¹⁸	Same population without mTBI	RBANS	Total Score	92.16 (11.932)	96.71 (11.672)	p = 0.023	NR
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Same population without mTBI; none with poor effort	Overall Test Battery Mean z-score	---	0.00 (0.55)	0.11 (0.42)	“NS”	d = 0.22
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Same population without mTBI; none have Axis I	Overall Test Battery Mean z-score	---	0.14 (0.69)	0.26 (0.38)	“NS”	d = .23
Schiehser, Delis, Filoteo, et al., 2011 ³⁵	Same population with moderate/severe TBI	Attention/Processing Speed Composite Score	---	9.8 (1.9)	8.5 (2.0)	p < .01	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Overall Test Battery Mean z-score	---	0.14 (0.69)	-.14 (.45)	“NS”	NR
Nelson, Hoelzle, McGuire, et al., 2010 ³⁰	Forensic context	Overall Test Battery Mean z-score	---	-0.15 (0.55)	-0.75 (0.52)	“Significant”	NR
Ruff, Ruff, & Wang, 2009 ³⁴	Headache intervention involving sleep hygiene, Prazosin, headache and pain education, and group therapy	MOCA	---	24.50 (.49)	28.60 (.59)	p < .001	NR
Ruff, Riechers, Wang, et al., 2012 ³²	LOC	MOCA	---	28.9 (.32)	25.1 (.18)	p < .001	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	Mental Health Diagnosis other than PTSD	WAIS-III (73 participants administered); WAIS-IV (9 participants administered)	WAIS-III Vocabulary, Information, Matrix Reasoning, Block Design Subscales; WAIS-IV: all subtests	97.4 (11.0)	100.2 (15.4)	NR	NR
Ruff, Ruff, & Wang, 2008 ³³	Positive neurological and/or neuropsychological findings	Number of blast exposures associated with LOC or AOC	---	2.65 (.18)	4.42 (.23)	p < .001	NR
		Number of blast exposures associated with LOC only	---	1.46 (.09)	3.91 (.20)	p < .001	NR
Gordon, Fitzpatrick, Hilsabeck, 2011 ²²	PTSD Diagnosis	WAIS-III (73 participants administered); WAIS-IV: (9 participants administered)	WAIS-III Vocabulary, Information, Matrix Reasoning, Block Design Subscales; WAIS-IV: all subtests	97.4 (11.0)	95.6 (13.3)	NR	NR

Table 1i. Self-reported cognitive deficits in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI	Cognitive Cluster	10.01 (4.90)	6.88 (5.25)	p < 0.001	NR
			Concentration	2.16 (1.17)	1.40 (1.32)	p < 0.001	NR
			Decision-Making	1.74 (1.22)	1.04 (1.15)	p < 0.001	NR
			Memory	2.36 (1.16)	1.81 (1.34)	p < 0.001	NR
			Slowed Thinking/Organization	1.89 (1.18)	1.11 (1.21)	p < 0.001	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Memory	---	1 (5.6)	4 (11.8)	“NS”	NR
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	Blast exposure	NSI	Decision-Making	NR	NR	p > .002	NR
			Memory	NR	NR	p > .002	NR
			Slowed Thinking/Organization	NR	NR	p > .002	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	HADS Anxiety	NSI	Concentration	NR	NR	p < .001	r = .56
			Decision-Making	NR	NR	p < .001	r = .52
			Memory	NR	NR	p < .001	r = .49
			Slowed Thinking/Organization	NR	NR	p < .001	r = .58
	HADS Depression	NSI	Concentration	NR	NR	p < .001	r = .55
			Decision-Making	NR	NR	p < .001	r = .57
			Memory	NR	NR	p < .001	r = .51
			Slowed Thinking/Organization	NR	NR	p < .001	r = .62
	LOC and/or PTA	NSI	Concentration	2.08 (1.21)	2.71 (1.12)	NR	NR
			Decision-Making	1.59 (1.365)	2.05 (1.36)	NR	NR
			Memory	2.17 (1.16)	2.86 (1.18)	NR	NR
			Slowed Thinking/Organization	1.80 (1.32)	2.38 (1.29)	NR	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI	Concentration	2.31 (1.08)	NA	NA	NA
			Decision-Making	1.72 (1.12)	NA	NA	NA
			Memory	2.50 (1.04)	NA	NA	NA
			Slowed Thinking/Organization	1.96 (1.18)	NA	NA	NA

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	PCL score > 50	NSI	Decision-Making	NR	NR	p < .002	NR
			Memory	NR	NR	p < .002	NR
			Slowed Thinking/ Organization	NR	NR	p < .002	NR
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI	Cognitive Cluster	3.04 (3.40)	13.32 (3.90)	p < .0001	NR
			Concentration	0.47 (0.79)	2.87 (0.88)	p < .0001	NR
			Decision-Making	0.33 (0.73)	2.28 (1.13)	p < .0001	NR
			Memory	0.95 (1.14)	3.00 (0.91)	p < .0001	NR
			Slowed Thinking/ Organization	0.42 (0.81)	2.57 (0.98)	p < .0001	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	PCL-M score	NSI	Concentration	NR	NR	p < .001	r = .59
			Decision-Making	NR	NR	p < .001	r = .63
			Memory	NR	NR	p < .001	r = .53
			Slowed Thinking/ Organization	NR	NR	p < .001	r = .68
Schiehser, Delis, Filoteo, et al., 2011 ³⁵	Same population with moderate/severe TBI	FrSBe	Subjective Executive Dysfunction pre- to post-injury change	10.8 (14.2)	21.6 (18.3)	p = .01	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	Service connected	NSI	Concentration	NR	NR	p < .05	NR
			Decision-Making	NR	NR	p < .05	NR
			Memory	NR	NR	p < .001	NR
			Slowed Thinking/ Organization	NR	NR	p < .05	NR

Physical Health Outcomes

Table 2a. Headache outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD) or % of subjects	Comparison group Mean (SD) or % of subjects	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI: Headaches	2.71 (1.10)	1.45 (1.30)	p < 0.001	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Headache	6 (33.3)	12 (35.3)	“NS”	NR
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	Blast exposure	NSI: Headaches	NR	NR	p < .002	NR
Cooper, Chau, Armistead-Jehle, et al., 2012 ¹⁶	Blast exposure	HIT-6	54.32 (9.44)	56.03 (9.54)	p = 0.489	NR
Ruff, Ruff, & Wang, 2009 ³⁴	Headache intervention involving sleep hygiene, Prazosin, headache and pain education, and group therapy	Headache frequency (number per month)	12.40 (.94)	4.77 (.19)	p < .001	NR
		Headache pain level (scale 0-10)	7.28 (.27)	4.08 (.19)	p < .001	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI: Headaches	2.29 (1.04)	NA	NA	NA
Patil, St. Andre, Crisen, et al., 2011 ³¹	NA	Neurology referral for headaches	82/246 = 33.3%	NA	NA	NA
		Neurology referral for headaches, Chronic daily type	11/246 = 4.47%	NA	NA	NA
		Neurology referral for headaches, Cluster type	1/246 = 0.41%	NA	NA	NA
		Neurology referral for headaches, Migraine type	25/246 = 10.16%	NA	NA	NA
		Neurology referral for headaches, Mixed type	8/246 = 3.25%	NA	NA	NA
		Neurology referral for headaches, Other type	1/246 = 0.41%	NA	NA	NA
		Neurology referral for headaches, Post-traumatic type	4/246 = 1.63%	NA	NA	NA
		Neurology referral for headaches, Tension type	6/246 = 2.44%	NA	NA	NA
Theeler & Erickson, 2009 ³⁸	NA	Headaches started < 1 week after trauma	12/33 = 36%	NA	NA	NA
		Headaches started > 1 month after trauma	3/33 = 9%	NA	NA	NA
		Headaches started 1 week to 1 month after trauma	1/33 = 3%	NA	NA	NA
		Unspecified onset of headache after trauma	5/33 = 15%	NA	NA	NA
		Worsening of pre-existing headaches	12/33 = 36%	NA	NA	NA
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	PCL score > 50	NSI: Headaches	NR	NR	p < .002	NR
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI: Headaches	1.01 (1.15)	2.79 (1.18)	p < .0001	NR

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD) or % of subjects	Comparison group Mean (SD) or % of subjects	p value for comparison	Magnitude of effect
Ruff, Ruff, & Wang, 2008 ³³	Positive neurological and/or neuropsychological findings	Headache frequency: > 10 per month	0/0 = 0%	31/74 = 42%	p < .001	NR
		Headache frequency: > 4 per month	2/6 = 33%	71/74 = 96%	p < .001	NR
		Headache frequency: Daily	0/0 = 0%	10/14 = 14%	p < .001	NR
		Headache pain level (scale 0-10)	4.33 (.27)	7.28 (.27)	p < .001	NR
		Headache: Migraine type	0/0 = 0%	14/74 = 19%	p < .001	NR
		Headache: Mixed type	0/0 = 0%	30/74 = 41%	p < .001	NR
		Headache: Tension type	6/6 = 100%	30/74 = 41%	p < .001	NR
Patil, St. Andre, Crisen, et al., 2011 ³¹	Referral to Neurology clinic for headaches	NSI: Headache	1.97 (0.91)	2.87 (0.80)	p < .01	NR
Theeler & Erickson, 2009 ³⁸	Same population without mTBI	Headache days per month after deployment (days/month)	11.9 (10.0)	10.3 (8.0)	NR	NR
		Headache days per month during deployment (days/month)	14.5 (11.7)	9.4 (9.4)	NR	NR
		Headache duration (hours)	8.8 (7.3)	7.5 (5.2)	NR	NR
		Headache NOS	7/33 = 21%	7/48 = 14%	NR	NR
		Headache severity (0-10 scale)	7.1 (1.5)	7.1 (1.2)	NR	NR
		Medication overuse headache	4/33 = 12%	0/48 = 0%	NR	NR
		MIDAS	30.8 (44.3)	26.8 (27.5)	NR	NR
		Migraine with aura	8/33 = 24%	3/48 = 6%	NR	NR
		Migraine without aura	15/33 = 45%	30/48 = 62%	NR	NR
		Multiple headache types	10/33 = 30%	16/48 = 33%	NR	NR
		Occipital headache	5/33 = 15%	3/48 = 6%	NR	NR
		Probable migraine	3/33 = 9%	5/48 = 10%	NR	NR
Tension-type headache	5/33 = 15%	13/48 = 27%	NR	NR		

Table 2b. Pain outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group	Comparison group	p value for comparison	Magnitude of effect
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	Pain (On a scale of 0 to 10, with 0 as no pain and 10 as the worst pain possible, how would you rate your current pain?)	Median = 3.5	Median = 2.0	p = .18	Cohen's d = .30
Lew, Pogoda, Hsu, et al., 2010 ²⁷	Same population with moderate/severe TBI	Pain in the past 30-days	112/125 = 90%	6/6 = 100%	p = .53	NR

Table 2c. Vestibular outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	Blast exposure	NSI: Feeling Dizzy	NR	NR	p > .002	NR
		NSI: Loss of Balance	NR	NR	p > .002	NR
		NSI: Poor Coordination	NR	NR	p > .002	NR
	PCL score > 50	NSI: Feeling Dizzy	NR	NR	p < .002	NR
		NSI: Loss of Balance	NR	NR	p < .002	NR
		NSI: Poor Coordination	NR	NR	p < .002	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI: Feeling Dizzy	1.47 (.86)	NA	NA	NA
		NSI: Loss of Balance	1.32 (.92)	NA	NA	NA
		NSI: Poor Coordination	1.32 (.93)	NA	NA	NA
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI: Feeling Dizzy	0.52 (0.81)	1.88 (1.13)	p < .0001	NR
		NSI: Loss of Balance	0.45 (0.66)	1.89 (1.04)	p < .0001	NR
		NSI: Poor Coordination	0.30 (0.55)	1.98 (1.04)	p < .0001	NR
Gottshall, Drake, Gray, et al., 2003 ²³	Control volunteer subjects without TBI. Not explicitly stated whether controls came from the same population as cases, but the controls were evaluated at the same time and place.	DHI	NR	NR	p < .01 for weeks 1, 2, 3, & 4 following injury	NR
	Control volunteer subjects without TBI. Not explicitly stated whether controls came from the same population as cases, but the controls were evaluated at the same time and place.	DVAT	NR	NR	p < .01 for week 1 p > .01 for week 4	NR
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI: Feeling Dizzy	1.44 (1.09)	0.94 (1.05)	p < 0.001	NR
		NSI: Loss of Balance	1.30 (1.08)	0.92 (0.98)	p = 0.002	NR
		NSI: Poor Coordination	1.71 (1.05)	0.92 (1.03)	p < 0.001	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Disorientation	7 (38.9)	13 (38.2)	"NS"	NR
		Dizziness	5 (27.8)	8 (23.5)	"NS"	NR
		Imbalance	2 (11.1)	4 (11.8)	"NS"	NR

Table 2d. Vision outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group	Comparison group	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI: Sensitivity to Light	1.56 (1.26)	0.85 (1.22)	p < 0.001	NR
		NSI: Vision Problems	1.18 (1.16)	0.86 (1.16)	p = 0.022	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Photophobia	4 (22.2)	6 (17.6)	“NS”	NR
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	Blast exposure	NSI: Sensitivity to Light	NR	NR	p > .002	NR
		NSI: Vision Problems	NR	NR	p > .002	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI: Sensitivity to Light	1.72 (1.17)	NA	NA	NA
		NSI: Vision Problems	1.51 (1.07)	NA	NA	NA
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	PCL score > 50	NSI: Sensitivity to Light	NR	NR	p < .002	NR
		NSI: Vision Problems	NR	NR	p < .002	NR
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI: Sensitivity to Light	0.65 (0.99)	2.06 (1.44)	p < .0001	NR
		NSI: Vision Problems	0.51 (0.99)	1.68 (1.23)	p < .0001	NR

Table 2e. Hearing outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	Blast exposure	NSI Hearing Difficulty	NR	NR	p < .002	NR
		Sensitivity to Noise	NR	NR	p > .002	NR
	PCL score > 50	NSI Hearing Difficulty	NR	NR	p < .002	NR
		Sensitivity to Noise	NR	NR	p < .002	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI Hearing Difficulty	1.88 (1.06)	NA	NA	NA
		Sensitivity to Noise	1.85 (1.11)	NA	NA	NA
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI Sensitivity to Noise	0.48 (0.87)	2.50 (1.09)	p < .0001	NR
		Hearing Difficulty	0.70 (1.02)	2.06 (1.11)	p < .0001	NR
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI Sensitivity to Noise	1.86 (1.17)	1.29 (1.41)	p < 0.001	NR
		Hearing Difficulty	1.46 (1.12)	1.39 (1.28)	p = 0.620	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Tinnitus	8 (44.4)	13 (38.2)	“NS”	NR
		Phonophobia	4 (22.2)	4 (11.8)	“NS”	NR

Table 2f. Neurological outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD) or % of subjects	Comparison group Mean (SD) or % of subjects	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI: Numbness or Tingling	1.02 (1.08)	1.22 (1.32)	p = 0.157	NR
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	Blast exposure	NSI: Numbness or Tingling	NR	NR	p > .002	NR
Ruff, Riechers, Wang, et al., 2012 ³²	LOC	Neurological deficits based on examination	0/16 = 0%	65/125 = 52%	p < .001	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI: Numbness or Tingling	1.61 (1.19)	NA	NA	NA
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	PCL score > 50	NSI: Numbness or Tingling	NR	NR	p < .002	NR
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI: Numbness or Tingling	0.76 (0.96)	1.83 (1.26)	p < .0001	NR

Table 2g. Nausea/appetite outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	Blast exposure	NSI Change in Taste or Smell	NR	NR	p > .002	NR
		Loss of Appetite	NR	NR	p > .002	NR
		Nausea	NR	NR	p > .002	NR
	PCL score > 50	NSI Change in Taste or Smell	NR	NR	p < .002	NR
		Loss of Appetite	NR	NR	p < .002	NR
		Nausea	NR	NR	p < .002	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI Change in Taste or Smell	0.82 (1.03)	NA	NA	NA
		Loss of Appetite	1.53 (1.13)	NA	NA	NA
		Nausea	1.13 (1.01)	NA	NA	NA
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI Change in Taste or Smell	0.15 (0.60)	1.14 (1.14)	p < .0001	NR
		Loss of Appetite	0.50 (0.84)	2.05 (1.10)	p < .0001	NR
		Nausea	0.31 (0.70)	1.57 (1.12)	p < .0001	NR
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI Change in Taste or Smell	0.54 (0.92)	0.49 (0.90)	p = 0.681	NR
		Loss of Appetite	1.44 (1.11)	1.13 (1.16)	p = 0.026	NR
		Nausea	1.12 (1.07)	0.60 (0.94)	p < 0.001	NR
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	Nausea	3 (16.7)	4 (11.8)	"NS"	NR

Mental Health Outcomes

Table 3a. PTSD outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	PCL-C				
		Arousal With Reminder	3.02 (1.39)	2.17 (1.36)	p < 0.001	NR
		Avoid Activities	2.50 (1.39)	2.03 (1.28)	p = 0.004	NR
		Avoid Thoughts	2.92 (1.42)	2.31 (1.36)	p < 0.001	NR
		Avoidance Cluster	17.72 (7.54)	13.93 (6.61)	p < 0.001	NR
		Disturbing Dreams	3.10 (1.42)	2.74 (1.47)	p = 0.042	NR
		Disturbing Memories	3.32 (1.32)	3.03 (1.43)	p = 0.086	NR
		Feeling Distant	2.97 (1.43)	2.10 (1.22)	p < 0.001	NR
		Feeling Numb	2.31 (1.41)	1.85 (1.19)	p = 0.004	NR
		Future Cut Short	2.29 (1.40)	1.85 (1.24)	p = 0.006	NR
		Hyper-Arousal Cluster	17.47 (4.91)	13.65 (5.58)	p < 0.001	NR
		Irritability	3.18 (1.32)	2.51 (1.40)	p < 0.001	NR
		Jumpy, Easily Startled	3.34 (1.30)	2.72 (1.41)	p < 0.001	NR
		Loss of Interest	2.55 (1.41)	2.01 (1.19)	p = 0.001	NR
		Poor Sleep	4.11 (1.16)	3.36 (1.48)	p < 0.001	NR
		Re-experiencing Cluster	14.95 (5.74)	12.07 (5.92)	p < 0.001	NR
		Reliving Experience	2.54 (1.29)	1.99 (1.26)	p < 0.001	NR
		Super-Alert, Watchful	3.44 (1.28)	2.67 (1.39)	p < 0.001	NR
		Total Score	50.14 (16.66)	39.64 (16.72)	p < 0.001	NR
		Trouble Concentrating	3.41 (1.26)	2.46 (1.39)	p < 0.001	NR
Trouble Remembering	2.48 (1.42)	2.05 (1.32)	p = 0.009	NR		
Upset When Reminded	2.98 (1.34)	2.42 (1.36)	p = 0.001	NR		
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	CAPS	9.8 (12.1)	54.2 (52.5)	“Significant”	NR
		SCID-I: PTSD	0 (0.0)	24 (70.6)	“NS”	NR
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	Blast exposure	PCL	37.3 (17.6)	41.5 (17.4)	p = .047	NR
Cooper, Chau, Armistead-Jehle, et al., 2012 ¹⁶	Blast exposure	PCL-M	36.29 (14.72)	37.88 (16.42)	p = .696	NR

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Kennedy, Leal, Lewis, et al., 2010 ²⁶	Blast exposure	PCL-C	Avoidance All: 15.6 (7.0) AOC: 14.8 (6.9) LOC: 16.1 (7.1)	All: 15.8 (7.4) AOC: 14.9 (7.1) LOC: 16.3 (7.5)	All: p = .826 AOC: "NS" LOC: "NS"	NR
			Hyper-Arousal All: 14.5 (5.7) AOC: 14.0 (5.6) LOC: 14.8 (5.8)	All: 15.2 (5.6) AOC: 14.6 (5.4) LOC: 15.6 (5.7)	All: p = .202 AOC: "NS" LOC: "NS"	NR
			Re-experiencing All: 12.0 (5.7) AOC: 11.8 (5.8) LOC: 12.2 (5.8)	All: 13.3 (5.9) AOC: 12.6 (5.7) LOC: 13.7 (5.9)	All: p = .020 AOC: "NS" LOC: "NS"	NR
			Total Score All: 42.7 (16.9) AOC: 40.6 (17.0) LOC: 43.1 (17.0)	All: 44.3 (17.6) AOC: 42.0 (16.8) LOC: 45.7 (17.9)	All: p = .198 AOC: "NS" LOC: "NS"	NR
Lippa, Pasternik, Benge, & Thornton, 2010 ²⁸	Blast exposure	PCL: Civilian and Military versions	49.75 (15.11)	54.45 (14.98)	p = .005	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	HADS Anxiety	PCL-M	NR	NR	p < .001	r = .76
	HADS Depression	PCL-M	NR	NR	p < .001	r = .76
Ruff, Riechers, Wang, et al., 2012 ³²	LOC	PTSD diagnosis (based on PCL-M as well as clinical interview)	1/16 = 6.25%	83/125 = 66%	p < .001	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	LOC and/or PTA	PCL-M	50.61 (16.24)	56.74 (14.61)	p < .05	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	PCL-C: Total Score	53.5 (15.6)	NA	NA	NA
Spencer et al., 2010 ³⁶	NA	PCL-M	52.4 (15.0)	NA	NA	NA
Ruff, Ruff, & Wang, 2008 ³³	Positive neurological and/or neuropsychological findings	PTSD diagnosis: (PCL score > 50 and meeting DSM-IV criteria)	11/46 = 24%	72/80 = 90%	p < .001	NR
Patil, St. Andre, Crisen, et al., 2011 ³¹	Referral to neurology clinic for headaches	PTSD (clinician confirmed or self-reported symptoms)	116/164 = 70.7%	66/82 = 80.5%	p = .10	NR
Belanger, Kretzmer, Venderploeg, & French, 2010 ¹⁰	Same population with moderate/severe TBI	PCL	35.4 (16.8)	23.5 (13.7)	p < .0001	NR
Belanger, Kretzmer, Yoash-Gantz, Pickett, & Tupler, 2009 ¹¹	Same population with moderate/severe TBI	PCL	45.5 (17.2)	30.1 (15.5)	p < .0001	NR
Cooper, Nelson, Armistead-Jehle, & Bowles, 2011 ¹⁹	Same population with moderate/severe TBI	PCL-M	42 (1)	37 (3)	NR	NR
Lew, Pogoda Hsu, et al., 2010 ²⁷	Same population with moderate/severe TBI	PTSD diagnosis	91/125 = 73%	5/6 = 83%	p = .49	NR
Cooper, Nelson, Armistead-Jehle, & Bowles, 2011 ¹⁹	Same population without mTBI	PCL-M	42 (1)	31 (4)	NR	NR
Gaylord, 2008 ²¹	Same population without mTBI	PCL-M: Score <= 44	14/31 = 45%	10/45 = 22%	p = .0345	NR
Theeler & Erickson, 2009 ³⁸	Same population without mTBI	PCL-C	34.6 (13.3)	36.0 (14.0)	NR	NR

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	CAPS: B – Re-experiencing Subscale	20.02 (7.03)	17.53 (6.06)	p = .08	Cohen’s d = 0.38
		CAPS: Total Score	74.02 (16.21)	67.20 (13.21)	p = .03	Cohen’s d = 0.46
		PCL-S	61.86 (11.04)	58.19 (12.89)	p = .17	Cohen’s d = 0.30
Cooper, Nelson, Armistead-Jehle, & Bowles, 2011 ¹⁹	Same population, neurologic patients (tumor, stroke, electrical, subarachnoid hemorrhage, anoxia, encephalitis, Parkinson’s disease, fronto-temporal dementia)	PCL-M	42 (1)	35 (5)	NR	NR
Spencer et al., 2010 ³⁶	Self-reported attention deficits	PCL-M	NR	NR	p < .001	r = .60
		PCL-M	NR	NR	p < .001	r = .48
		PCL-M	NR	NR	p < .001	r = .54
	Self-reported memory deficits	PCL-M	NR	NR	p < .001	r = .48
	Self-reported slowed thinking/organization	PCL-M	NR	NR	p < .001	r = .54

Table 3b. Measures of anxiety in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group Mean (SD)	Comparison group Mean (SD)	p value for comparison	Magnitude of effect
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	Blast exposure	NSI: Feeling Anxious	NR	NR	“NS”	NR
	PCL score > 50	NSI: Feeling Anxious	NR	NR	p < .002	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI: Feeling Anxious	2.58 (1.08)	NA	NA	NA
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI: Feeling Anxious	0.48 (0.82)	2.96 (0.91)	p < .0001	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	LOC and/or PTA	HADS: Anxiety	11.56 (4.06)	13.16 (3.92)	p < .05	NR
	HADS Depression	HADS: Anxiety	NR	NR	p < .001	r = .56
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI: Feeling Anxious	1.99 (1.24)	1.49 (1.34)	p = 0.001	NR
Spencer, et al., 2010 ³⁶	NA	HADS: Anxiety	11.9 (4.5)	NA	NA	NA
	Self-reported slowed thinking/organization	HADS: Anxiety	NR	NR	p < .001	r = .39
	Self-reported attention deficits	HADS: Anxiety	NR	NR	p < .001	r = .48
	Self-reported memory deficits	HADS: Anxiety	NR	NR	p < .001	r = .33

Table 3c. Measures of depression in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD) or % of subjects	Comparison group Mean (SD) or % of subjects	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI	Feeling Depressed or Sad	1.16 (1.26)	1.08 (1.19)	p = 0.566	NR
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	Blast exposure	NSI	Feeling Depressed or Sad	NR	NR	“NS”	NR
Drag, Spencer, Walker, et al., 2012 ²⁰	LOC and/or PTA	HADS	Depression	7.95 (4.61)	10.34 (4.013)	p < .001	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI	Feeling Depressed or Sad	2.09 (1.20)	NA	NA	NA
Spencer et al., 2010 ³⁶	NA	HADS	Depression	8.7 (4.3)	NA	NA	NA
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	PCL score > 50	NSI	Feeling Depressed or Sad	NR	NR	p < .002	NR
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI	Depression	0.30 (0.64)	2.44 (1.19)	p < .0001	NR
Swick, Honzel, Larsen, et al., 2012 ³⁷	Same population without mTBI; all participants had PTSD.	BDI-II	---	20.0 (12.3)	20.8 (9.2)	NR	NR
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	BDI-II	---	31.56 (11.06)	29.17 (10.53)	p = .29	Cohen’s d = 0.23
		Hopelessness (Who or what gives you strength and hope?)	---	6/46 = 13%	6/46 = 13%	NR	NR
		SCID-I	Major Depressive Disorder	25/46 = 54%	18/46 = 39%	p = .14	φ = .15
Spencer, et al., 2010 ³⁶	Self-reported attention deficits	HADS	Depression	NR	NR	p < .001	r = .45
	Self-reported memory deficits	HADS	Depression	NR	NR	p < .001	r = .36
	Self-reported slowed thinking/organization	HADS	Depression	NR	NR	p < .001	r = .52

Table 3d. Substance use disorders in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group % of subjects	Comparison group % of subjects	p value for comparison	Magnitude of effect
Nelson, Hoelzle, Doane, et al., 2012 ²⁹	Axis I disorder	SCID-I	Alcohol Abuse/Dependence	5/18 = 27.8%	12/34 = 35.3%	“Significant”	NR
	Same population without mTBI; none have Axis I		Alcohol Abuse/Dependence	5/18 = 27.8%	5/28 = 17.9%	“NS”	NR
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	SCID-I	Alcohol Problem	13/46 = 28%	17/46 = 37%	p = .37	φ = .09
			Drug Problem	4/46 = 9%	7/46 = 15%	p = .34	φ = .10

Table 3e. Suicide-related outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group % of subjects	Comparison group % of subjects	p value for comparison	Magnitude of effect
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	Suicidal ideation (Have you had thoughts about death or about killing yourself?)	11/44 = 25%	5/44 = 11%	p = .10	φ = .18
		Suicidal Intent (Have you ever intended to commit suicide?)	3/46 = 7%	1/46 = 2%	NR	NR
		Past Suicide Attempts (Have you ever attempted suicide?)	2/46 = 4%	2/46 = 4%	NR	NR

Table 3f. Other mental health outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group	Comparison group	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI	Affective Cluster	7.21 (4.21)	5.43 (4.54)	p < 0.001	NR
			Frustration	1.86 (1.22)	1.24 (1.29)	p < 0.001	NR
			Irritability	2.19 (1.23)	1.62 (1.32)	p < 0.001	NR
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	Blast exposure	NSI	Frustration	NR	NR	“NS”	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI	Frustration	2.41 (1.17)	NA	NA	NA
			Irritability	2.76 (1.06)	NA	NA	NA
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	PCL score > 50	NSI	Frustration	NR	NR	p < .002	NR
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI	Affective Cluster	2.97 (2.93)	14.64 (3.48)	p < .0001	NR
			Frustration	0.31 (0.68)	2.82 (0.95)	p < .0001	NR
			Irritability	0.63 (0.89)	2.95 (0.90)	p < .0001	NR
Schiehser, Delis, Filoteo, et al., 2011 ³⁵	Same population with moderate/severe TBI	FrSBe	Apathy, pre- to post-injury change	14.7 (17.2)	27.8 (16.8)	p < .01	NR
			Behavioral Disinhibition, pre- to post-injury change	5.3 (7.8)	5.5 (11.2)	p = .98	NR
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	SCID-I	Any Co-Morbid Axis I Disorder	34/46 = 78%	29/46 = 63%	p = .14	φ = .12
Cooper, Mercado-Couch, Critchfield, et al., 2010 ¹⁸	Same population without mTBI	Psychiatric Diagnosis	---	25 (50.0%)	26 (22.2%)	p = 0.001	NR

Functional/Social Outcomes

Table 4a. Employment outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group % of subjects or OR (95% CI) compared to non-mTBI control	Comparison group % of subjects or OR (95% CI) compared to non-mTBI control	p value for comparison	Magnitude of effect
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	Unemployment (% who responded 'No' to 'Are you employed?')	9/45 = 20%	14/45 = 31%	p = .23	φ = .13
Toblin, Riviere, Thomas, et al., 2012 ³⁹	LOC	Missed work: ≥2 missed workdays in the past month	1.8 (95% CI 0.9-3.5)	1.4 (95% CI 0.5-3.6)	NR	NR
		Occupational Impairment, Heavy Load: Difficulty carrying a heavy load in past month	2.2 (95% CI 1.3-3.5)	3.0 (95% CI 1.5-5.7)	NR	NR
		Occupational Impairment, Physical Training: Difficulty performing physical training (PT) in past month	1.9 (95% CI 1.2-2.9)	1.6 (95% CI 0.8-3.0)	NR	NR

Table 4b. Sleep outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	Subscale or test component	mTBI group Mean (SD), Median (IQR), or % of subjects	Comparison group Mean (SD), Median (IQR), or % of subjects	p value for comparison	Magnitude of effect
Kennedy, Cullen, Amador, et al., 2010 ²⁵	At least one additional AIS code	NSI	Fatigue	1.85 (1.11)	1.51 (1.14)	p = 0.009	NR
			Sleep	2.73 (1.10)	2.15 (1.33)	p < 0.001	NR
Belanger, Proctor-Weber, Kretzmer, et al. 2011 ¹²	Blast exposure	NSI	Fatigue	NR	NR	p < .002	NR
Ruff, Ruff, & Wang, 2009 ³⁴	Headache intervention involving sleep hygiene, Prazosin, headache and pain education, and group therapy	ESS	---	16.10 (0.28)	7.28 (0.34)	p < .001	NR
Benge, Pastorek, & Thornton, 2009 ¹³	NA	NSI	Fatigue	2.10 (1.11)	NA	NA	NA
			Sleep	2.72 (1.17)	NA	NA	NA
Coldren, Russell, Parish, et al., 2012 ¹⁵	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	Sleep	hours per night < 4	6/47 = 13%	7/108 = 7%	p = 0.21 for sleep hours overall	NR
			hours per night ≥4	40/47 = 87%	99/108 = 93%	p = 0.21 for sleep hours overall	NR
		Sleep Change	< 2 hour loss	10/47 = 23%	4/108 = 6%	p = 0.02 for sleep change overall	NR
			> 2 hour loss	33/47 = 77%	62/108 = 94%	p = 0.02 for sleep change overall	NR
Kelly, Coldren, Parish, et al., 2012 ²⁴	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	Sleep	hours per night	Median = 6 (5-7)	Median = 6 (5-7)	p = .22	NR
		Sleep Change (negative value = hours lost)	---	Median = 0 (-2.5-0)	Median = 0 (0-0)	p < 0.001	NR
Belanger, Proctor-Weber, Kretzmer, et al., 2011 ¹²	PCL score > 50	NSI	Fatigue	NR	NR	p < .002	NR
Cooper, Kennedy, Cullen, et al., 2011 ¹⁷	PCL-C ≥60 (controls had PCL-C ≤30)	NSI	Fatigue	0.86 (0.91)	2.60 (1.04)	p < .0001	NR
			Sleep	1.24 (1.17)	3.45 (0.70)	p < .0001	NR
Ruff, Ruff, & Wang, 2008 ³³	Positive neurological and/or neuropsychological findings	NSI	Sleep	5/46 = 11%	66/80 = 82.5%	p < .001	NR
Patil, St. Andre, Crisen, et al., 2011 ³¹	Referral to Neurology clinic for headaches	NSI	Sleep	2.53 (1.17)	2.78 (1.12)	p = .11	NR
Lew, Pogoda Hsu, et al., 2010 ²⁷	Same population with moderate/severe TBI	Sleep disturbance in the past 30 days	---	2.72 (1.24)	3.17 (.75)	p = .38	NR

Table 4c. Social outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome measure	mTBI group % of subjects	Comparison group % of subjects	p value for comparison	Magnitude of effect
Barnes, Walter, & Chard, 2012 ⁹	Same population without mTBI	Lack of emotional support (% who responded 'No' to 'Do you have an emotional support system?')	9/35 = 26%	6/35 = 17%	p = .38	φ = .10
		Marital status (Are you married?)	NR	NR	p = .72	φ = .04

Service Utilization/Costs

Table 5. Service utilization outcomes in studies of U.S. Veterans and members of the U.S. military with mild TBI

Author, year	Comparison group description	Outcome Measure	mTBI group OR (95% CI) compared to non-mTBI control, % of subjects, or Mean (SD)	Comparison group OR (95% CI) compared to non-mTBI control, % of subjects, or Mean (SD)	p value for comparison	Magnitude of effect
Toblin, Riviere, Thomas, et al., 2012 ³⁹	LOC	Medical utilization: ≥2 "sick call" visits in past month	2.0 (95% CI 1.3-3.1)	1.9 (95% CI 1.04-3.6)	NR	NR
Coldren, Russell, Parish, et al., 2012 ¹⁵	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	Current Counseling	2/47 = 4%	4/108 = 4%	p >0.99	NR
		Current Mental Health Medication	2/47 = 4%	9/108 = 9%	p = 0.50	NR
Kelly, Coldren, Parish, et al., 2012 ²⁴	Same population without mTBI; with minor traumatic injuries not involving the head and noninjured volunteers from same population	Current Counseling	4/66 = 6%	5/146 = 3%	p = .46	NR
		Current Mental Health Medication	3/66 = 5%	8/146 = 6%	p = .99	NR
Belanger, Kretzmer, Venderploeg, & French, 2010 ¹⁰	Same population with moderate/severe TBI	Currently taking pain medications	24/44 = 55%	37/64 = 57%	p = .44	NR
Gaylord 2008 ²¹	Same population without mTBI	Length of Hospital Stay (days)	PTSD: 13.6 (9.8) No PTSD: 19.0 (20.9)	PTSD: 16.0 (26.6) No PTSD: 14.1 (22.2)	NR	NR
		Length of Intensive Care Unit Stay (days)	PTSD: 2.0 (6.7) No PTSD: 12.1 (12.1)	PTSD: 1.6 (3.9) No PTSD: 2.8 (8.9)	NR	NR
Swick, Honzel, Larsen, et al., 2012 ³⁷	Same population without mTBI; all participants had PTSD.	Number of medications	18	5	NR	NR
Cooper, Mercado-Couch, Critchfield, et al., 2010 ¹⁸	Same population without mTBI	Pain Medication (Narcotic)	32 (64.0%)	85 (72.6%)	p = 0.264	NR

APPENDIX F. LIST OF EXCLUDED STUDIES GROUPED BY REASON FOR EXCLUSION

<i>Population does not meet criteria for adult, human subjects who are Veterans or members of the military from any country</i>
Axelrod BN, Vanderploeg RD, Rawlings DB. WAIS-R prediction equations in patients with traumatic brain injury. <i>J Clin Exp Neuropsychol</i> . Jun 1999;21(3):368-374.
Belanger HG, Curtiss G, Demery JA, Lebowitz BK, Vanderploeg RD. Factors moderating neuropsychological outcomes following mild traumatic brain injury: a meta-analysis. <i>J Int Neuropsychol Soc</i> . May 2005;11(3):215-227.
Belanger HG, Spiegel E, Vanderploeg RD. Neuropsychological performance following a history of multiple self-reported concussions: a meta-analysis. <i>J Int Neuropsychol Soc</i> . Mar 2010;16(2):262-267.
Binder LM. Assessment of malingering after mild head trauma with the Portland Digit Recognition Test.[Erratum appears in <i>J Clin Exp Neuropsychol</i> 1993 Nov;15(6):852]. <i>J Clin Exp Neuropsychol</i> . Mar 1993;15(2):170-182.
Bryan C, Anestis M. Reexperiencing symptoms and the interpersonal-psychological theory of suicidal behavior among deployed service members evaluated for traumatic brain injury. <i>J Clin Psychol</i> . Sep 2011;67(9):856-865.
Bryan C, Hernandez AM. Magnitudes of decline on Automated Neuropsychological Assessment Metrics subtest scores relative to predeployment baseline performance among service members evaluated for traumatic brain injury in Iraq. <i>J Head Trauma Rehabil</i> . Jan-Feb 2012;27(1):45-54.
Bryan CJ, Clemans TA, Hernandez AM. Perceived burdensomeness, fearlessness of death, and suicidality among deployed military personnel. <i>Personality and Individual Differences</i> . Feb 2012;52(3):374-379.
Bryan CJ, Hernandez AM. Predictors of post-traumatic headache severity among deployed military personnel. <i>Headache</i> . Jun 2011;51(6):945-953.
Cho DY, Wang YC. Comparison of the APACHE III, APACHE II and Glasgow Coma Scale in acute head injury for prediction of mortality and functional outcome. <i>Intensive Care Med</i> . Jan 1997;23(1):77-84.
Crawford F, Crynen G, Reed J, et al. Identification of plasma biomarkers of TBI outcome using proteomic approaches in an APOE mouse model. <i>J Neurotrauma</i> . Jan 20 2012;29(2):246-260.
Dobscha SK, Clark ME, Morasco BJ, Freeman M, Campbell R, Helfand M. Systematic review of the literature on pain in patients with polytrauma including traumatic brain injury. <i>Pain Med</i> . Oct 2009;10(7):1200-1217.
Eagleman DM. Using time perception to measure fitness for duty. <i>Military Psychology</i> . Jan 2009;21(Suppl 1):S123-S129.
Gervasio AH, Kreutzer JS. Kinship and family members' psychological distress after traumatic brain injury: A large sample study. <i>The Journal of Head Trauma Rehabilitation</i> . Jun 1997;12(3):14-26.
Hoffer ME, Gottshall KR, Moore R, Balough BJ, Wester D. Characterizing and treating dizziness after mild head trauma. <i>Otol Neurotol</i> . Mar 2004;25(2):135-138.
Hoffman JM, Lucas S, Dikmen S, et al. Natural history of headache after traumatic brain injury. <i>J Neurotrauma</i> . Sep 2011;28(9):1719-1725.
Kane RL, Roebuck-Spencer T, Short P, Kabat M, Wilken J. Identifying and monitoring cognitive deficits in clinical populations using Automated Neuropsychological Assessment Metrics (ANAM) tests. <i>Arch Clin Neuropsychol</i> . Feb 2007;22 Suppl 1:S115-126.

Kreutzer JS, Marwitz JH, Hsu N, Williams K, Riddick A. Marital stability after brain injury: an investigation and analysis. <i>NeuroRehabilitation</i> . 2007;22(1):53-59.
Lew HL, Chen CPC, Chen MJL, Hsu THC, Tang SFT, Date ES. Comparing the effects of different speech targets on cognitive event-related potentials: theoretical implications for evaluating brain injury. <i>Am J Phys Med Rehabil</i> . Jul 2002;81(7):524-528.
Lew HL, Gray M, Poole JH. Temporal stability of auditory event-related potentials in healthy individuals and patients with traumatic brain injury. <i>J Clin Neurophysiol</i> . Oct 2007;24(5):392-397.
Lew HL, Thomander D, Gray M, Poole JH. The effects of increasing stimulus complexity in event-related potentials and reaction time testing: clinical applications in evaluating patients with traumatic brain injury. <i>J Clin Neurophysiol</i> . Oct 2007;24(5):398-404.
Lewine JD, Davis JT, Bigler ED, et al. Objective documentation of traumatic brain injury subsequent to mild head trauma: multimodal brain imaging with MEG, SPECT, and MRI. <i>J Head Trauma Rehabil</i> . May-Jun 2007;22(3):141-155.
Luethcke CA, Bryan CJ, Morrow CE, Isler WC. Comparison of Concussive Symptoms, Cognitive Performance, and Psychological Symptoms Between Acute Blast-Versus Nonblast-Induced Mild Traumatic Brain Injury. <i>J Int Neuropsychol Soc</i> . 2011;17(1):36-45.
Maas AI, Harrison-Felix CL, Menon D, et al. Common data elements for traumatic brain injury: recommendations from the interagency working group on demographics and clinical assessment. <i>Arch Phys Med Rehabil</i> . Nov 2010;91(11):1641-1649.
Montella D, Brown SH, Elkin PL, et al. Comparison of SNOMED CT versus Medcin terminology concept coverage for mild Traumatic Brain Injury. <i>AMIA Annu Symp Proc</i> . 2011;2011:969-978.
Norman SB, Trim RS, Goldsmith AA, et al. Role of risk factors proximate to time of trauma in the course of PTSD and MDD symptoms following traumatic injury. <i>J Trauma Stress</i> . Aug 2011;24(4):390-398.
Schmid KE, Tortella FC. The diagnosis of traumatic brain injury on the battlefield. <i>Front Neurol</i> . 2012;3:90.
Smith-Seemiller L, Fow NR, Kant R, Franzen MD. Presence of post-concussion syndrome symptoms in patients with chronic pain vs mild traumatic brain injury. <i>Brain Inj</i> . 2003;17(3):199-206.
Publication contains no primary data and is not a systematic review of primary studies
Aksionoff EB, Falk NS. The differential diagnosis of perceptual deficits in traumatic brain injury patients. <i>J Am Optom Assoc</i> . Aug 1992;63(8):554-558.
Anonymous. Cognitive rehabilitation therapy in traumatic brain injury. <i>The Lancet</i> . Oct 2011;378(9801):1440.
Anonymous. The neurological burden of the war in Iraq and Afghanistan. <i>Ann Neurol</i> . Oct 2006;60(4):A13-15.
Arcure J, Harrison EE. A review of the use of early hypothermia in the treatment of traumatic brain injuries. <i>J Spec Oper Med</i> . 2009;9(3):22-25.
Barr M. Post concussion syndrome hypoglycemia and Agent Orange. <i>Aust Fam Physician</i> . Apr 1983;12(4):224.
Barrett RS. Post-traumatic headache. Combat soldiers are suffering. <i>Adv NPs PAs</i> . Jan 2012;3(1):33-34.
Belanger HG, Uomoto JM, Vanderploeg RD. The Veterans Health Administration's (VHA's) Polytrauma System of Care for mild traumatic brain injury: costs, benefits, and controversies. <i>J Head Trauma Rehabil</i> . Jan-Feb 2009;24(1):4-13.
Belanger HG, Vanderploeg RD, Curtiss G, Warden DL. Recent neuroimaging techniques in mild traumatic brain injury. <i>J Neuropsychiatry Clin Neurosci</i> . 2007;19(1):5-20.

Betthausen LM, Bahraini N, Kregel MH, Brenner LA. Self-report measures to identify post traumatic stress disorder and/or mild traumatic brain injury and associated symptoms in military veterans of Operation Enduring Freedom (OEF)/Operation Iraqi Freedom (OIF). <i>Neuropsychol Rev</i> . Mar 2012;22(1):35-53.
Bhattacharjee Y. Neuroscience. Shell shock revisited: solving the puzzle of blast trauma. <i>Science</i> . Jan 25 2008;319(5862):406-408.
Bogdanova Y, Verfaellie M. Cognitive sequelae of blast-induced traumatic brain injury: recovery and rehabilitation. <i>Neuropsychol Rev</i> . Mar 2012;22(1):4-20.
Brancu M, Straits-Troster K, Kudler H. Behavioral health conditions among military personnel and veterans: prevalence and best practices for treatment. <i>N C Med J</i> . Jan-Feb 2011;72(1):54-60.
Brenner LA, Vanderploeg RD, Terrio H. Assessment and diagnosis of mild traumatic brain injury, posttraumatic stress disorder, and other polytrauma conditions: burden of adversity hypothesis. <i>Rehabil Psychol</i> . Aug 2009;54(3):239-246.
Brenner LA. Neuropsychological and neuroimaging findings in traumatic brain injury and post-traumatic stress disorder. <i>Dialogues Clin Neurosci</i> . 2011;13(3):311-323.
Bryant RA. Disentangling mild traumatic brain injury and stress reactions. <i>The New England Journal of Medicine</i> . Jan 2008;358(5):525-527.
Caldrony RD, Radike J. Experience with mild traumatic brain injuries and postconcussion syndrome at Kandahar, Afghanistan. <i>US Army Med Dep J</i> . Jul-Sep 2010:22-30.
Cancio LC, Chung KK. The role of normoventilation in improving traumatic brain injury outcomes. <i>US Army Med Dep J</i> . Apr-Jun 2011:49-54.
Cernich AN, Kurtz SM, Mordecai KL, Ryan PB. Cognitive rehabilitation in traumatic brain injury. <i>Curr Treat Options Neurol</i> . Sep 2010;12(5):412-423.
Cifu DX, Cohen SI, Lew HL, Jaffee M, Sigford B. The history and evolution of traumatic brain injury rehabilitation in military service members and veterans. <i>Am J Phys Med Rehabil</i> . Aug 2010;89(8):688-694.
Connors S, Gordon WA, Hovda DA. Care of war veterans with mild traumatic brain injury. <i>N Engl J Med</i> . Jul 30 2009;361(5):536-537; author reply 537-538.
Cooke BB, Keltner NL. Biological perspectives: traumatic brain injury-war related: part II. <i>Perspect Psychiatr Care</i> . Jan 2008;44(1):54-57.
Cote MJ, Syam SS, Vogel WB, Cowper DC. A mixed integer programming model to locate traumatic brain injury treatment units in the Department of Veterans Affairs: a case study. <i>Health Care Manag Sci</i> . Sep 2007;10(3):253-267.
Cozzarelli TA. Evaluation and treatment of persistent cognitive dysfunction following mild traumatic brain injury. <i>J Spec Oper Med</i> . 2010;10(1):39-42.
Daggett V, Bakas T, Habermann B. A review of health-related quality of life in adult traumatic brain injury survivors in the context of combat veterans. <i>J Neurosci Nurs</i> . Apr 2009;41(2):59-71.
Das RR. Mild traumatic brain injury in U.S. soldiers returning from Iraq. <i>N Engl J Med</i> . May 15 2008;358(20):2177-2178; author reply 2179.
Dempsey KE, Dorlac WC, Martin K, et al. Landstuhl Regional Medical Center: traumatic brain injury screening program. <i>J Trauma Nurs</i> . Jan-Mar 2009;16(1):6-7, 10-12.
Department of Veterans A. Schedule for rating disabilities; evaluation of residuals of traumatic brain injury (TBI). Final rule. <i>Fed Regist</i> . Sep 23 2008;73(185):54693-54708.

Dolan S, Martindale S, Robinson J, et al. Neuropsychological sequelae of PTSD and TBI following war deployment among OEF/OIF veterans. <i>Neuropsychol Rev</i> . Mar 2012;22(1):21-34.
Doster J. Traumatic brain injury: serving our returning soldiers. <i>Tenn Med</i> . Oct 2007;100(10):45, 48.
Eibner C, Schell TL, Jaycox LH. Care of war veterans with mild traumatic brain injury. <i>N Engl J Med</i> . Jul 30 2009;361(5):537; author reply 537-538.
Eldar R, Jelic M. The association of rehabilitation and war. <i>Disabil Rehabil</i> . Sep 16 2003;25(18):1019-1023.
Evans RW. Posttraumatic headaches among United States soldiers injured in Afghanistan and Iraq. <i>Headache: The Journal of Head and Face Pain</i> . Sep 2008;48(8):1216-1225.
Felber ES. Combat-related posttraumatic headache: diagnosis, mechanisms of injury, and challenges to treatment. <i>J Am Osteopath Assoc</i> . Dec 2010;110(12):737-738.
Giles GM. Cognitive versus functional approaches to rehabilitation after traumatic brain injury: Commentary on a randomized controlled trial. <i>Am J Occup Ther</i> . Jan-Feb 2010;64(1):182-185.
Graham DP, Cardon AL. An update on substance use and treatment following traumatic brain injury. <i>Ann N Y Acad Sci</i> . Oct 2008;1141:148-162.
Graver CJ. "Reference data from the Automated Neuropsychological Assessment Metrics for use in traumatic brain injury in an active duty military sample": Comment. <i>Mil Med</i> . Dec 2009;174(12):iv-v.
Helmick K, Members of Consensus C. Cognitive rehabilitation for military personnel with mild traumatic brain injury and chronic post-concussional disorder: Results of April 2009 consensus conference. <i>NeuroRehabilitation</i> . 2010;26(3):239-255.
Hesdorffer DC, Rauch SL, Tamminga CA. Long-term psychiatric outcomes following traumatic brain injury: a review of the literature. <i>J Head Trauma Rehabil</i> . Nov-Dec 2009;24(6):452-459.
Hicks RR, Fertig SJ, Desrocher RE, Koroshetz WJ, Pancrazio JJ. Neurological effects of blast injury. <i>J Trauma</i> . May 2010;68(5):1257-1263.
Hoge CW, Castro CA. Blast-related traumatic brain injury in U.S. military personnel. <i>N Engl J Med</i> . Sep 1 2011;365(9):860; author reply 860-861.
Hoge CW, Goldberg HM, Castro CA. Care of war veterans with mild traumatic brain injury--flawed perspectives. <i>N Engl J Med</i> . Apr 16 2009;360(16):1588-1591.
Hoge CW. "Traumatic brain injury screening: Preliminary findings in a US Army Brigade Combat Team": Comment. <i>The Journal of Head Trauma Rehabilitation</i> . Jul-Aug 2009;24(4):299-301.
Jackson GL, Hamilton NS, Tupler LA. Detecting traumatic brain injury among veterans of Operations Enduring and Iraqi Freedom. <i>N C Med J</i> . Jan-Feb 2008;69(1):43-47.
Jaffee MS, Helmick KM, Girard PD, Meyer KS, Dinegar K, George K. Acute clinical care and care coordination for traumatic brain injury within Department of Defense. <i>J Rehabil Res Dev</i> . 2009;46(6):655-666.
Jasiewicz J, Kearns W, Craighead J, Fozard JL, Scott S, McCarthy J, Jr. Smart rehabilitation for the 21st century: the Tampa Smart Home for veterans with traumatic brain injury. <i>J Rehabil Res Dev</i> . 2011;48(8):vii-xviii.
Kaplan GB, Vasterling JJ, Vedak PC. Brain-derived neurotrophic factor in traumatic brain injury, post-traumatic stress disorder, and their comorbid conditions: role in pathogenesis and treatment. <i>Behav Pharmacol</i> . Sep 2010;21(5-6):427-437.
Kelly JC, Amerson EH, Barth JT. Mild traumatic brain injury: lessons learned from clinical, sports, and combat concussions. <i>rehabil</i> . 2012;2012:371970.

Kirjavainen S. Neuro-otological studies on brain injured ex-servicemen. Follow-up of 256 cases. <i>Acta Otolaryngol (Stockh)</i> . 1968;Suppl 233:231-259.
Krainin BM, Forsten RD, Kotwal RS, Lutz RH, Guskiewicz KM. Mild traumatic brain injury literature review and proposed changes to classification. <i>J Spec Oper Med</i> . 2011;11(3):38-47.
Leung LY, VandeVord PJ, Dal Cengio AL, Bir C, Yang KH, King AI. Blast related neurotrauma: a review of cellular injury. <i>Mol Cell Biomech</i> . Sep 2008;5(3):155-168.
Lew HL, Vanderploeg RD, Moore DF, et al. Overlap of mild TBI and mental health conditions in returning OIF/OEF service members and veterans. <i>J Rehabil Res Dev</i> . 2008;45(3):xi-xvi.
Lew HL, Weihing J, Myerse PJ, Pogoda TK, Goodrich GL. Dual sensory impairment (DSI) in traumatic brain injury (TBI)-An emerging interdisciplinary challenge. <i>NeuroRehabilitation</i> . 2010;26(3):213-222.
Lew HL. Rehabilitation needs of an increasing population of patients: Traumatic brain injury, polytrauma, and blast-related injuries. <i>J Rehabil Res Dev</i> . Jul-Aug 2005;42(4):xiii-xvi.
Lutz RH, Kane S, Lay J. Evidence-based diagnosis and management of mTBI in forward deployed settings: the genesis of the USASOC neurocognitive testing and post-injury evaluation and treatment program. <i>J Spec Oper Med</i> . 2010;10(1):23-38.
McCrea M, Pliskin N, Barth J, et al. Official position of the military TBI task force on the role of neuropsychology and rehabilitation psychology in the evaluation, management, and research of military veterans with traumatic brain injury. <i>Clin Neuropsychol</i> . Jan 2008;22(1):10-26.
Miller G. Neuropathology. Blast injuries linked to neurodegeneration in veterans. <i>Science</i> . May 18 2012;336(6083):790-791.
Miller G. The invisible wounds of war. Healing the brain, healing the mind. <i>Science</i> . Jul 29 2011;333(6042):514-517.
Okie S. Traumatic brain injury in the war zone. <i>N Engl J Med</i> . May 19 2005;352(20):2043-2047.
Pogoda TK, Vanderploeg RD, Cifu DX, Tun CG, Lew HL. Re: separating deployment-related traumatic brain injury and posttraumatic stress disorder in veterans: preliminary findings from the VA TBI screening program. <i>Am J Phys Med Rehabil</i> . Dec 2009;88(12):1043-1044; author reply 1044-1045.
Robinson RG, Jorge R. Longitudinal course of mood disorders following traumatic brain injury. <i>Arch Gen Psychiatry</i> . Jan 2002;59(1):23-24.
Ropper A. Brain injuries from blasts. <i>N Engl J Med</i> . Jun 2 2011;364(22):2156-2157.
Ruff RL. Mild traumatic brain injury in U.S. soldiers returning from Iraq. <i>N Engl J Med</i> . May 15 2008;358(20):2178; author reply 2179.
Ruff RM, Iverson GL, Barth JT, Bush SS, Broshek DK. Recommendations for diagnosing a mild traumatic brain injury: A National Academy of Neuropsychology education paper. <i>Arch Clin Neuropsychol</i> . Feb 2009;24(1):3-10.
Sayer NA, Rettmann NA, Carlson KF, et al. Veterans with history of mild traumatic brain injury and posttraumatic stress disorder: challenges from provider perspective. <i>J Rehabil Res Dev</i> . 2009;46(6):703-716.
Shenton M, Hamoda H, Schneiderman J, et al. A review of magnetic resonance imaging and diffusion tensor imaging findings in mild traumatic brain injury. <i>Brain Imaging and Behavior</i> . Jun 2012;6(2):137-192.
Sigford B, Cifu DX, Vanderploeg R. Care of war veterans with mild traumatic brain injury. <i>N Engl J Med</i> . Jul 30 2009;361(5):536; author reply 537-538.
SoRelle R. Head injuries linked to Alzheimer's disease. <i>Circulation</i> . Oct 31 2000;102(18):E9036-9037.

Stelmack J. Measuring outcomes of neuro-optometric care in traumatic brain injury. <i>Journal of Behavioral Optometry</i> . 2007;18(3):67-71.
Stonesifer LD. Mild traumatic brain injury in U.S. soldiers returning from Iraq. <i>N Engl J Med</i> . May 15 2008;358(20):2178; author reply 2179.
Taber KH, Hurley RA. PTSD and combat-related injuries: functional neuroanatomy. <i>J Neuropsychiatry Clin Neurosci</i> . 2009;21(1):1 p preceding 1, 1-4.
Walcott BP, Kahle KT. Blast-related traumatic brain injury in U.S. military personnel. <i>N Engl J Med</i> . Sep 1 2011;365(9):860.
Walker RL, Clark ME, Nampiaparampil DE, et al. The hazards of war: blast injury headache. <i>J Pain</i> . Apr 2010;11(4):297-302.
Xydakis MS, Butman JA, Carlo P. Blast-related traumatic brain injury in U.S. military personnel. <i>The New England Journal of Medicine</i> . Sep 2011;365(9):859.
<i>Does not distinguish mild TBI from moderate or severe</i>
Alterman AI, Tarter RE. Relationship between familial alcoholism and head injury. <i>Journal of Studies on Alcohol</i> . May 1985;46(3):256-258.
Armistead-Jehle P, Hansen CL. Comparison of the Repeatable Battery for the Assessment of Neuropsychological Status Effort Index and stand-alone symptom validity tests in a military sample. <i>Arch Clin Neuropsychol</i> . Nov 2011;26(7):592-601.
Ashla PM, McMurtray AM, Licht E, Mendez MF. Retrospective posttraumatic amnesia in traumatic brain injury. <i>The Journal of Neuropsychiatry and Clinical Neurosciences</i> . Fal 2009;21(4):467-468.
Belanger HG, King-Kallimanis B, Nelson AL, Schonfeld L, Scott SG, Vanderploeg RD. Characterizing wandering behaviors in persons with traumatic brain injury residing in veterans health administration nursing homes. <i>Arch Phys Med Rehabil</i> . Feb 2008;89(2):244-250.
Belanger HG, Vanderploeg RD, Soble JR, Richardson M, Groer S. Validity of the Veterans Health Administration's traumatic brain injury screen. <i>Arch Phys Med Rehabil</i> . Jul 2012;93(7):1234-1239.
Blennow K, Jonsson M, Andreasen N, et al. No neurochemical evidence of brain injury after blast overpressure by repeated explosions or firing heavy weapons. <i>Acta Neurol Scand</i> . Apr 2011;123(4):245-251.
Braden CA, Cuthbert JP, Brenner L, et al. Health and wellness characteristics of persons with traumatic brain injury. <i>Brain Inj</i> . 2012;26(11):1315-1327.
Brahm KD, Wilgenburg HM, Kirby J, Ingalla S, Chang C-Y, Goodrich GL. Visual impairment and dysfunction in combat-injured servicemembers with traumatic brain injury. <i>Optom Vis Sci</i> . Jul 2009;86(7):817-825.
Brenner LA, Betthausen LM, Homaifar BY, et al. Posttraumatic stress disorder, traumatic brain injury, and suicide attempt history among veterans receiving mental health services. <i>Suicide Life Threat Behav</i> . Aug 2011;41(4):416-423.
Brenner LA, Homaifar BY, Adler LE, Wolfman JH, Kemp J. Suicidality and veterans with a history of traumatic brain injury: precipitants events, protective factors, and prevention strategies. <i>Rehabil Psychol</i> . Nov 2009;54(4):390-397.
Brenner LA, Ignacio RV, Blow FC. Suicide and traumatic brain injury among individuals seeking Veterans Health Administration services. <i>J Head Trauma Rehabil</i> . Jul-Aug 2011;26(4):257-264.
Breshears RE, Brenner LA, Harwood JEF, Gutierrez PM. Predicting suicidal behavior in veterans with traumatic brain injury: the utility of the personality assessment inventory. <i>J Pers Assess</i> . Jul 2010;92(4):349-355.

Brooks E. Understanding veterans' service options and utilization patterns for PTSD and TBI. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering</i> . 2011;71(8-B):4739.
Campbell TA, Nelson LA, Lumpkin R, Yoash-Gantz RE, Pickett TC, McCormick CL. Neuropsychological measures of processing speed and executive functioning in combat veterans with PTSD, TBI, and comorbid TBI/PTSD. <i>Psychiatric Annals</i> . Aug 2009;39(8):796-803.
Capaldi VF, 2nd, Guerrero ML, Killgore WDS. Sleep disruptions among returning combat veterans from Iraq and Afghanistan. <i>Mil Med</i> . Aug 2011;176(8):879-888.
Carey ME, Young HF, Mathis JL. The outcome of 89 American and 224 Vietnamese sustaining brain wounds in Vietnam. <i>Mil Med</i> . Apr 1974;139(4):281-284.
Carey ME, Young HF, Rish BL, Mathis JL. Follow-up study of 103 American soldiers who sustained a brain wound in Vietnam. <i>J Neurosurg</i> . Nov 1974;41(5):542-549.
Carlson KF, Meis LA, Jensen AC, et al. Caregiver reports of subsequent injuries among veterans with traumatic brain injury after discharge from inpatient polytrauma rehabilitation programs. <i>J Head Trauma Rehabil</i> . Jan-Feb 2012;27(1):14-25.
Carlson KF, Nelson D, Orazem RJ, Nugent S, Cifu DX, Sayer NA. Psychiatric diagnoses among Iraq and Afghanistan war veterans screened for deployment-related traumatic brain injury. <i>J Trauma Stress</i> . Feb 2010;23(1):17-24.
Caveness WF, Walker AE, Ascroft PB. Incidence of posttraumatic epilepsy in Korean veterans as compared with those from World War I and World War II. <i>J Neurosurg</i> . Feb 1962;19:122-129.
Cernich AN, Chandler L, Scherdell T, Kurtz S. Assessment of co-occurring disorders in veterans diagnosed with traumatic brain injury. <i>J Head Trauma Rehabil</i> . Jul-Aug 2012;27(4):253-260.
Chauhan NB, Gatto R. Synergistic benefits of erythropoietin and simvastatin after traumatic brain injury. <i>Brain Res</i> . Nov 11 2010;1360:177-192.
Chemtob CM, Muraoka MY, Wu-Holt P, Fairbank JA, Hamada RS, Keane TM. Head injury and combat-related posttraumatic stress disorder. <i>J Nerv Ment Dis</i> . Nov 1998;186(11):701-708.
Chen JWY, Ruff RL, Eavey R, Wasterlain CG. Posttraumatic epilepsy and treatment. <i>J Rehabil Res Dev</i> . 2009;46(6):685-696.
Cohen SP, Plunkett AR, Wilkinson I, et al. Headaches during war: analysis of presentation, treatment, and factors associated with outcome. <i>Cephalalgia</i> . Jan 2012;32(2):94-108.
Cox DJ, Davis M, Singh H, et al. Driving rehabilitation for military personnel recovering from traumatic brain injury using virtual reality driving simulation: a feasibility study. <i>Mil Med</i> . Jun 2010;175(6):411-416.
Crawford FC, Vanderploeg RD, Freeman MJ, et al. APOE genotype influences acquisition and recall following traumatic brain injury. <i>Neurology</i> . Apr 9 2002;58(7):1115-1118.
Cullen N, Chundamala J, Bayley M, Jutai J, Erabi G. The efficacy of acquired brain injury rehabilitation. <i>Brain Inj</i> . Feb 2007;21(2):113-132.
Das RR, Moorthi RN. Traumatic brain injury in the war zone. <i>N Engl J Med</i> . Aug 11 2005;353(6):633-634.
Demakis GJ, Hammond F, Knotts A, et al. The Personality Assessment Inventory in individuals with traumatic brain injury. <i>Arch Clin Neuropsychol</i> . Jan 2007;22(1):123-130.
Detweiler MB, Arif S, Candelario J, et al. Salem VAMC-U.S. Army Fort Bragg Warrior Transition Clinic telepsychiatry collaboration: 12-month operation clinical perspective. <i>Telemed J E Health</i> . Mar 2012;18(2):81-86.

Donnelly KT, Donnelly JP, Dunnam M, et al. Reliability, sensitivity, and specificity of the VA traumatic brain injury screening tool. <i>J Head Trauma Rehabil.</i> Nov-Dec 2011;26(6):439-453.
Drake AI, Meyer KS, Cessante LM, et al. Routine TBI screening following combat deployments. <i>NeuroRehabilitation.</i> 2010;26(3):183-189.
Egede LE, Dismuke C, Echols C. Racial/Ethnic disparities in mortality risk among US veterans with traumatic brain injury. <i>Am J Public Health.</i> May 2012;102 Suppl 2:S266-271.
Elbogen EB, Johnson SC, Wagner HR, Newton VM, Beckham JC. Financial well-being and postdeployment adjustment among Iraq and Afghanistan war veterans. <i>Mil Med.</i> Jun 2012;177(6):669-675.
Elbogen EB, Wagner HR, Fuller SR, et al. Correlates of anger and hostility in Iraq and Afghanistan war veterans. <i>Am J Psychiatry.</i> Sep 2010;167(9):1051-1058.
Eonta SE, Carr W, McArdle JJ, et al. Automated Neuropsychological Assessment Metrics: repeated assessment with two military samples. <i>Aviat Space Environ Med.</i> Jan 2011;82(1):34-39.
Ettinger AB, Copeland LA, Zeber JE, Van Cott AC, Pugh MJV. Are psychiatric disorders independent risk factors for new-onset epilepsy in older individuals? <i>Epilepsy Behav.</i> Jan 2010;17(1):70-74.
Evans CD. Rehabilitation of the brain-damaged survivor. <i>Injury.</i> Nov 1976;8(2):80-97.
Fabing HD. Cerebral blast syndrome in combat soldiers. <i>Arch Neurol Psychiatry.</i> Jan 1947;57(1):14-57.
Fenimore EA. Depression following traumatic brain injury and the Hamilton Depression Rating Scale. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering.</i> 2006;67(3-B):1698.
Friedemann-Sanchez G, Sayer NA, Pickett T. Provider perspectives on rehabilitation of patients with polytrauma. <i>Arch Phys Med Rehabil.</i> Jan 2008;89(1):171-178.
Galarneau MR, Woodruff SI, Dye JL, Mohrle CR, Wade AL. Traumatic brain injury during Operation Iraqi Freedom: findings from the United States Navy-Marine Corps Combat Trauma Registry. <i>J Neurosurg.</i> May 2008;108(5):950-957.
Galloway MS, Fink DS, Millikan AM, Bell MR. Factors Associated With Physical Aggression Among US Army Soldiers. <i>Aggress Behav.</i> Sep 2012;38(5):357-367.
Goldstein G, Allen DN, Caponigro JM. A retrospective study of heterogeneity in neurocognitive profiles associated with traumatic brain injury. <i>Brain Inj.</i> 2010;24(4):625-635.
Gontkovsky ST, Souheaver GT. T-score and raw-score comparisons in detecting brain dysfunction using the booklet category test and the short category test. <i>Percept Mot Skills.</i> Feb 2002;94(1):319-322.
Gordon NG. Diagnostic efficiency of the Trail Making Test as a function of cut-off score, diagnosis, and age. <i>Percept Mot Skills.</i> Aug 1978;47(1):191-195.
Griffin JM, Friedemann-Sanchez G, Jensen AC, et al. The invisible side of war: families caring for US service members with traumatic brain injuries and polytrauma. <i>J Head Trauma Rehabil.</i> Jan-Feb 2012;27(1):3-13.
Gutierrez PM, Brenner LA, Huggins JA. A preliminary investigation of suicidality in psychiatrically hospitalized veterans with traumatic brain injury. <i>Arch.</i> 2008;12(4):336-343.
Han SD, Suzuki H, Drake AI, Jak AJ, Houston WS, Bondi MW. Clinical, cognitive, and genetic predictors of change in job status following traumatic brain injury in a military population. <i>J Head Trauma Rehabil.</i> Jan-Feb 2009;24(1):57-64.
Harmon A. A descriptive study of military family needs following a polytraumatic injury. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering.</i> 2008;69(2-B):1326.
Harrow JJ, Rashka SL, Fitzgerald SG, Nelson AL. Pressure ulcers and occipital alopecia in Operation Iraqi Freedom polytrauma casualties. <i>Mil Med.</i> Nov 2008;173(11):1068-1072.

Hill JJ, 3rd, Moberg BHP, Jr., Cullen MR. Separating deployment-related traumatic brain injury and posttraumatic stress disorder in veterans: preliminary findings from the Veterans Affairs traumatic brain injury screening program. <i>Am J Phys Med Rehabil.</i> Aug 2009;88(8):605-614.
Holsinger T, Steffens DC, Phillips C, et al. Head injury in early adulthood and the lifetime risk of depression. <i>Arch Gen Psychiatry.</i> Jan 2002;59(1):17-22.
Huckans M, Pavawalla S, Demadura T, et al. A pilot study examining effects of group-based Cognitive Strategy Training treatment on self-reported cognitive problems, psychiatric symptoms, functioning, and compensatory strategy use in OIF/OEF combat veterans with persistent mild cognitive disorder and history of traumatic brain injury. <i>J Rehabil Res Dev.</i> 2010;47(1):43-60.
Huggins JM, Brown JN, Capehart B, Townsend ML, Legge J, Melnyk SD. Medication adherence in combat veterans with traumatic brain injury. <i>Am J Health-Syst Pharm.</i> Feb 1 2011;68(3):254-258.
Ingala AM. The impact of military deployment on college adjustment. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering.</i> 2012;72(11-B):7083.
Iverson KM, Hendricks AM, Kimerling R, et al. Psychiatric diagnoses and neurobehavioral symptom severity among OEF/OIF VA patients with deployment-related traumatic brain injury: a gender comparison. <i>Womens Health Issues.</i> Jul-Aug 2011;21(4 Suppl):S210-217.
Johns JS, Cifu DX, Keyser-Marcus L, Jolles PR, Fratkin MJ. Impact of clinically significant heterotopic ossification on functional outcome after traumatic brain injury. <i>J Head Trauma Rehabil.</i> Jun 1999;14(3):269-276.
Jones A, Ingram M, Ben-Porath YS. Scores on the MMPI-2-RF Scales as a function of increasing levels of failure on cognitive symptom validity tests in a military sample. <i>The Clinical Neuropsychologist.</i> Jul 2012;26(5):790-815.
Kalkonde YV, Jawaid A, Qureshi SU, et al. Medical and environmental risk factors associated with frontotemporal dementia: A case-control study in a veteran population. <i>Alzheimer's Dement.</i> May 2012;8(3):204-210.
Kapidzic A, Vidovic M, Sinanovic O. Localisation of war craniocerebral injury as risk factor for posttraumatic epilepsy. <i>Med Arh.</i> 2011;65(6):343-344.
Kelley AM, Athy JR, Cho TH, Erickson B, King M, Cruz P. Risk propensity and health risk behaviors in U.S. army soldiers with and without psychological disturbances across the deployment cycle. <i>J Psychiatr Res.</i> May 2012;46(5):582-589.
Kennedy JE, Lumpkin RJ, Grissom JR. A survey of mild traumatic brain injury treatment in the emergency room and primary care medical clinics. <i>Mil Med.</i> Jun 2006;171(6):516-521.
Kilts JD, Tupler LA, Keefe FJ, et al. Neurosteroids and self-reported pain in veterans who served in the U.S. Military after September 11, 2001. <i>Pain Med.</i> Oct 2010;11(10):1469-1476.
King PR, Donnelly KT, Donnelly JP, et al. Psychometric study of the Neurobehavioral Symptom Inventory. <i>JRRD.</i> 2012;49(6):879-888.
King PR, Jr. A psychometric study of the Neurobehavioral Symptom Inventory. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering.</i> 2012;73(1-B):650.
Koenigs M, Huey ED, Raymond V, et al. Focal brain damage protects against post-traumatic stress disorder in combat veterans. <i>Nat Neurosci.</i> Feb 2008;11(2):232-237.
Kozminski M. Combat-related posttraumatic headache: diagnosis, mechanisms of injury, and challenges to treatment. <i>J Am Osteopath Assoc.</i> Sep 2010;110(9):514-519.
Kraft JF, Schwab KA, Salazar AM, Brown HR. Occupational and educational achievements of head injured Vietnam veterans at 15-year follow-up. <i>Arch Phys Med Rehabil.</i> Jun 1993;74(6):596-601.

Levin HS, Wilde E, Troyanskaya M, et al. Diffusion tensor imaging of mild to moderate blast-related traumatic brain injury and its sequelae. <i>J Neurotrauma</i> . Apr 2010;27(4):683-694.
Lew HL, Jerger JF, Guillory SB, Henry JA. Auditory dysfunction in traumatic brain injury. <i>J Rehabil Res Dev</i> . 2007;44(7):921-928.
Lew HL, Kraft M, Pogoda TK, Amick MM, Woods P, Cifu DX. Prevalence and characteristics of driving difficulties in Operation Iraqi Freedom/Operation Enduring Freedom combat returnees. <i>J Rehabil Res Dev</i> . 2011;48(8):913-925.
Lew HL, Otis JD, Tun C, Kerns RD, Clark ME, Cifu DX. Prevalence of chronic pain, posttraumatic stress disorder, and persistent postconcussive symptoms in OIF/OEF veterans: polytrauma clinical triad. <i>J Rehabil Res Dev</i> . 2009;46(6):697-702.
Lew HL, Pogoda TK, Baker E, et al. Prevalence of dual sensory impairment and its association with traumatic brain injury and blast exposure in OEF/OIF veterans. <i>J Head Trauma Rehabil</i> . Nov-Dec 2011;26(6):489-496.
Lew HL, Poole JH, Vanderploeg RD, et al. Program development and defining characteristics of returning military in a VA Polytrauma Network Site. <i>J Rehabil Res Dev</i> . 2007;44(7):1027-1034.
Lipsky RH, Sparling MB, Ryan LM, et al. Association of COMT Val158Met genotype with executive functioning following traumatic brain injury. <i>J Neuropsychiatry Clin Neurosci</i> . 2005;17(4):465-471.
Maestas KL, Bengel JF, Pastorek NJ, Lemaire A, Darrow R. Factor structure of posttraumatic stress disorder symptoms in OEF/OIF veterans presenting to a polytrauma clinic. <i>Rehabil Psychol</i> . Nov 2011;56(4):366-373.
Maguen S, Lau KM, Madden E, Seal K. Factors associated with completing comprehensive traumatic brain injury evaluation. <i>Mil Med</i> . Jul 2012;177(7):797-803.
Maguen S, Madden E, Lau KM, Seal K. The impact of head injury mechanism on mental health symptoms in veterans: do number and type of exposures matter? <i>J Trauma Stress</i> . Feb 2012;25(1):3-9.
Mantyla M. Post-traumatic cerebral atrophy. A study on brain-injured veterans on the Finnish wars of 1939-40 and 1941-45. <i>Ann Clin Res</i> . 1981;13(32):1-47.
Maxwell TM. The investigation of mild traumatic brain injury among the Afghanistan and Iraqi war veterans. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering</i> . 2011;72(6-B):3778.
McMurtray A, Clark DG, Christine D, Mendez MF. Early-onset dementia: frequency and causes compared to late-onset dementia. <i>Dement Geriatr Cogn Disord</i> . 2006;21(2):59-64.
Meterko M, Baker E, Stolzmann KL, Hendricks AM, Cicerone KD, Lew HL. Psychometric assessment of the Neurobehavioral Symptom Inventory-22: the structure of persistent postconcussive symptoms following deployment-related mild traumatic brain injury among veterans. <i>J Head Trauma Rehabil</i> . Jan-Feb 2012;27(1):55-62.
Miller KJ, Schwab KA, Warden DL. Predictive value of an early Glasgow Outcome Scale score: 15-month score changes. <i>J Neurosurg</i> . Aug 2005;103(2):239-245.
Morissette SB, Woodward M, Kimbrel NA, et al. Deployment-related TBI, persistent postconcussive symptoms, PTSD, and depression in OEF/OIF veterans. <i>Rehabil Psychol</i> . Nov 2011;56(4):340-350.
Nelson DV, Esty ML. Neurotherapy of traumatic brain injury/posttraumatic stress symptoms in OEF/OIF veterans. <i>J Neuropsychiatry Clin Neurosci</i> . Mar 1 2012;24(2):237-240.
Nelson LA, Yoash-Gantz RE, Pickett TC, Campbell TA. Relationship between processing speed and executive functioning performance among OEF/OIF veterans: implications for postdeployment rehabilitation. <i>J Head Trauma Rehabil</i> . Jan-Feb 2009;24(1):32-40.

Nuutila A, Huusko S. Epilepsy among brain-injured veterans 26 to 31 years following the injury. <i>Scand J Rehabil Med.</i> 1972;4(2):81-84.
Olson-Madden JH, Brenner L, Harwood JEF, Emrick CD, Corrigan JD, Thompson C. Traumatic brain injury and psychiatric diagnoses in veterans seeking outpatient substance abuse treatment. <i>J Head Trauma Rehabil.</i> Nov-Dec 2010;25(6):470-479.
Parsons TD, Courtney C, Rizzo AA, Armstrong C, Edwards J, Reger G. Virtual reality paced serial assessment test for neuropsychological assessment of a military cohort. <i>Stud Health Technol Inform.</i> 2012;173:331-337.
Pietrzak RH, Goldstein MB, Malley JC, et al. Posttraumatic growth in Veterans of Operations Enduring Freedom and Iraqi Freedom. <i>J Affect Disord.</i> Oct 2010;126(1-2):230-235.
Pugh MJV, Knoefel JE, Mortensen EM, Amuan ME, Berlowitz DR, Van Cott AC. New-onset epilepsy risk factors in older veterans. <i>J Am Geriatr Soc.</i> Feb 2009;57(2):237-242.
Quigley KS, McAndrew LM, Almeida L, et al. Prevalence of environmental and other military exposure concerns in Operation Enduring Freedom and Operation Iraqi Freedom veterans. <i>J Occup Environ Med.</i> Jun 2012;54(6):659-664.
Raymont V, Salazar AM, Krueger F, Grafman J. "Studying injured minds" - the Vietnam head injury study and 40[THIN SPACE]years of brain injury research. <i>Front Neurol.</i> 2011;2:15.
Raymont V, Salazar AM, Lipsky R, Goldman D, Tasick G, Grafman J. Correlates of posttraumatic epilepsy 35 years following combat brain injury. <i>Neurology.</i> Jul 20 2010;75(3):224-229.
Reiber GE, McFarland LV, Hubbard S, et al. Servicemembers and veterans with major traumatic limb loss from Vietnam war and OIF/OEF conflicts: survey methods, participants, and summary findings. <i>J Rehabil Res Dev.</i> 2010;47(4):275-297.
Resnik L, Gray M, Borgia M. Measurement of community reintegration in sample of severely wounded servicemembers. <i>J Rehabil Res Dev.</i> 2011;48(2):89-102.
Roberts RJ, Franzen K, Varney NR. Theta bursts, closed head injury, and partial seizure-like symptoms: a retrospective study. <i>Appl Neuropsychol.</i> 2001;8(3):140-147.
Rosenbaum M, Najenson T. Changes in life patterns and symptoms of low mood as reported by wives of severely brain-injured soldiers. <i>J Consult Clin Psychol.</i> Dec 1976;44(6):881-888.
Russell WR. The development of grand mal after missile wounds of the brain. <i>Johns Hopkins Med J.</i> May 1968;122(5):250-253.
Salazar AM, Warden DL, Schwab K, et al. Cognitive rehabilitation for traumatic brain injury: A randomized trial. Defense and Veterans Head Injury Program (DVHIP) Study Group. <i>Jama.</i> Jun 21 2000;283(23):3075-3081.
Salinsky M, Spencer D, Boudreau E, Ferguson F. Psychogenic nonepileptic seizures in US veterans. <i>Neurology.</i> Sep 6 2011;77(10):945-950.
Sayer NA, Chiros CE, Sigford B, et al. Characteristics and rehabilitation outcomes among patients with blast and other injuries sustained during the Global War on Terror. <i>Arch Phys Med Rehabil.</i> Jan 2008;89(1):163-170.
Sayer NA, Cifu DX, McNamee S, et al. Rehabilitation needs of combat-injured service members admitted to the VA Polytrauma Rehabilitation Centers: the role of PM&R in the care of wounded warriors. <i>Pm R.</i> Jan 2009;1(1):23-28.
Sayer NA, Nelson D, Nugent S. Evaluation of the Veterans Health Administration traumatic brain injury screening program in the upper Midwest. <i>J Head Trauma Rehabil.</i> Nov-Dec 2011;26(6):454-467.

Scholten JD, Sayer NA, Vanderploeg RD, Bidelspach DE, Cifu DX. Analysis of US Veterans Health Administration comprehensive evaluations for traumatic brain injury in Operation Enduring Freedom and Operation Iraqi Freedom Veterans. <i>Brain Inj.</i> Sep 2012;26(10):1177-1184.
Stelmack JA, Frith T, Van Koeveering D, Rinne S, Stelmack TR. Visual function in patients followed at a Veterans Affairs polytrauma network site: an electronic medical record review. <i>Optometry.</i> Aug 2009;80(8):419-424.
Streeter CC, Van Reekum R, Shorr RI, Bachman DL. Prior head injury in male veterans with borderline personality disorder. <i>J Nerv Ment Dis.</i> Sep 1995;183(9):577-581.
Sullivan EV, Corkin S. Selective subject attrition in a longitudinal study of head-injured veterans. <i>J Gerontol.</i> Nov 1984;39(6):718-720.
Taft CT, Kachadourian LK, Suvak MK, et al. Examining impelling and disinhibiting factors for intimate partner violence in veterans. <i>J Fam Psychol.</i> Apr 2012;26(2):285-289.
Taylor BC, Hagel EM, Carlson KF, et al. Prevalence and costs of co-occurring traumatic brain injury with and without psychiatric disturbance and pain among Afghanistan and Iraq War Veteran V.A. users. <i>Med Care.</i> Apr 2012;50(4):342-346.
Terrio H, Brenner LA, Ivins BJ, et al. Traumatic brain injury screening: preliminary findings in a US Army Brigade Combat Team. <i>J Head Trauma Rehabil.</i> Jan-Feb 2009;24(1):14-23.
Terrio HP, Nelson LA, Betthausen LM, Harwood JE, Brenner LA. Postdeployment traumatic brain injury screening questions: Sensitivity, specificity, and predictive values in returning soldiers. <i>Rehabil Psychol.</i> Feb 2011;56(1):26-31.
Van Dyke SA, Axelrod BN, Schutte C. Test-retest reliability of the Traumatic Brain Injury Screening Instrument. <i>Mil Med.</i> Dec 2010;175(12):947-949.
Vanderploeg RD, Curtiss G, Schinka JA, Lanham RA, Jr. Material-specific memory in traumatic brain injury: differential effects during acquisition, recall, and retention. <i>Neuropsychology.</i> Apr 2001;15(2):174-184.
Vanderploeg RD, Schwab K, Walker WC, et al. Rehabilitation of traumatic brain injury in active duty military personnel and veterans: Defense and Veterans Brain Injury Center randomized controlled trial of two rehabilitation approaches. <i>Arch Phys Med Rehabil.</i> Dec 2008;89(12):2227-2238.
Varney NR. Prognostic significance of anosmia in patients with closed-head trauma. <i>J Clin Exp Neuropsychol.</i> Mar 1988;10(2):250-254.
Vassallo JL, Proctor-Weber Z, Lebowitz BK, Curtiss G, Vanderploeg RD. Psychiatric risk factors for traumatic brain injury. <i>Brain Inj.</i> Jun 2007;21(6):567-573.
Vasterling JJ, Constans JI, Hanna-Pladdy B. Head injury as a predictor of psychological outcome in combat veterans. <i>J Trauma Stress.</i> Jul 2000;13(3):441-451.
Walker AE, Blumer D. The fate of World War II veterans with posttraumatic seizures. <i>Arch Neurol.</i> Jan 1989;46(1):23-26.
Walker AE, Leuchs HK, Lechtape-Gruter H, Caveness WF, Kretschman C. Life expectancy of head injured men with and without epilepsy. <i>Arch Neurol.</i> Feb 1971;24(2):95-100.
Walter KH, Barnes SM, Chard KM. The influence of comorbid MDD on outcome after residential treatment for veterans with PTSD and a history of TBI. <i>J Trauma Stress.</i> Aug 2012;25(4):426-432.
Walter KH, Kiefer SL, Chard KM. Relationship between posttraumatic stress disorder and postconcussive symptom improvement after completion of a posttraumatic stress disorder/traumatic brain injury residential treatment program. <i>Rehabil Psychol.</i> Feb 2012;57(1):13-17.

Weinstein EA, Lyerly OG. Conversion hysteria following brain injury. <i>Arch Neurol</i> . Nov 1966;15(5):545-548.
Weiss GH, Caveness WF, Einsiedel-Lechtape H, McNeel ML. Life expectancy and causes of death in a group of head-injured veterans of World War I. <i>Arch Neurol</i> . Dec 1982;39(12):741-743.
Welberg L. Traumatic brain injury: Brain trauma in military veterans. <i>Nat Rev Neurosci</i> . Jul 2012;13(7):450.
Wilder Schaaf KP. Polytrauma family needs assessment. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering</i> . 2010;71(5-B):3342.
Yaffe K, Vittinghoff E, Lindquist K, et al. Posttraumatic stress disorder and risk of dementia among US veterans. <i>Arch Gen Psychiatry</i> . Jun 2010;67(6):608-613.
Sample includes fewer than 30 mTBI cases
Akin FW, Murnane OD. Head injury and blast exposure: vestibular consequences. <i>Otolaryngol Clin North Am</i> . Apr 2011;44(2):323-334, viii.
Bahraini NH, Brenner LA, Harwood JEF, et al. Utility of the trauma symptom inventory for the assessment of post-traumatic stress symptoms in veterans with a history of psychological trauma and/or brain injury. <i>Mil Med</i> . Oct 2009;174(10):1005-1009.
Ben Arzi N, Solomon Z, Dekel R. Secondary traumatization among wives of PTSD and post-concussion casualties: distress, caregiver burden and psychological separation. <i>Brain Inj</i> . Aug 2000;14(8):725-736.
Brenner LA, Harwood JEF, Homaifar BY, Cawthra E, Waldman J, Adler LE. Psychiatric hospitalization and veterans with traumatic brain injury: a retrospective study. <i>J Head Trauma Rehabil</i> . Nov-Dec 2008;23(6):401-406.
Brenner LA, Ladley-O'Brien SE, Harwood JEF, et al. An exploratory study of neuroimaging, neurologic, and neuropsychological findings in veterans with traumatic brain injury and/or posttraumatic stress disorder. <i>Mil Med</i> . Apr 2009;174(4):347-352.
Carr ME, Jr., Masullo LN, Brown JK, Lewis PC. Creatine kinase BB isoenzyme blood levels in trauma patients with suspected mild traumatic brain injury. <i>Mil Med</i> . Jun 2009;174(6):622-625.
Chapman JC, Andersen AM, Roselli LA, Meyers NM, Pincus JH. Screening for mild traumatic brain injury in the presence of psychiatric comorbidities. <i>Arch Phys Med Rehabil</i> . Jul 2010;91(7):1082-1086.
Chard KM, Schumm JA, McIlvain SM, Bailey GW, Parkinson RB. Exploring the efficacy of a residential treatment program incorporating cognitive processing therapy-cognitive for veterans with PTSD and traumatic brain injury. <i>J Trauma Stress</i> . Jun 2011;24(3):347-351.
Classen S, Levy C, Meyer DL, Bewernitz M, Lanford DN, Mann WC. Simulated driving performance of combat veterans with mild traumatic brain injury and posttraumatic stress disorder: a pilot study. <i>Am J Occup Ther</i> . Jul-Aug 2011;65(4):419-427.
Davenport ND, Lim KO, Armstrong MT, Sponheim SR. Diffuse and spatially variable white matter disruptions are associated with blast-related mild traumatic brain injury. <i>Neuroimage</i> . Feb 1 2012;59(3):2017-2024.
Finkel A. Headaches in soldiers with mild traumatic brain injury - additional data. <i>Headache</i> . Sep 2012;52(8):1320.
Finkel AG, Yerry J, Scher A, Choi YS. Headaches in soldiers with mild traumatic brain injury: findings and phenomenologic descriptions. <i>Headache</i> . Jun 2012;52(6):957-965.
Homaifar BY, Brenner LA, Gutierrez PM, et al. Sensitivity and specificity of the Beck Depression Inventory-II in persons with traumatic brain injury. <i>Arch Phys Med Rehabil</i> . Apr 2009;90(4):652-656.

Homaifar BY, Harwood JE, Wagner TH, Brenner LA. Description of outpatient utilization and costs in group of veterans with traumatic brain injury. <i>J Rehabil Res Dev.</i> 2009;46(8):1003-1010.
Huang M-X, Nichols S, Robb A, et al. An automatic MEG low-frequency source imaging approach for detecting injuries in mild and moderate TBI patients with blast and non-blast causes. <i>Neuroimage.</i> Jul 2012;61(4):1067-1082.
Huang M-X, Theilmann RJ, Robb A, et al. Integrated imaging approach with MEG and DTI to detect mild traumatic brain injury in military and civilian patients. <i>J Neurotrauma.</i> Aug 2009;26(8):1213-1226.
Lew HL, Poole JH, Alvarez S, Moore W. Soldiers with occult traumatic brain injury. <i>Am J Phys Med Rehabil.</i> Jun 2005;84(6):393-398.
Matthews SC, Strigo IA, Simmons AN, O'Connell RM, Reinhardt LE, Moseley SA. A multimodal imaging study in U.S. veterans of Operations Iraqi and Enduring Freedom with and without major depression after blast-related concussion. <i>Neuroimage.</i> Jan 2011;54 Suppl 1:S69-75.
Peskind ER, Petrie EC, Cross DJ, et al. Cerebrocerebellar hypometabolism associated with repetitive blast exposure mild traumatic brain injury in 12 Iraq war Veterans with persistent post-concussive symptoms. <i>Neuroimage.</i> Jan 2011;54 Suppl 1:S76-82.
Roy MJ, Francis J, Friedlander J, et al. Improvement in cerebral function with treatment of posttraumatic stress disorder. <i>Ann N Y Acad Sci.</i> Oct 2010;1208:142-149.
Tan G, Fink B, Dao TK, et al. Associations among pain, PTSD, mTBI, and heart rate variability in veterans of Operation Enduring and Iraqi Freedom: a pilot study. <i>Pain Med.</i> Oct 2009;10(7):1237-1245.
Trudeau DL, Anderson J, Hansen LM, et al. Findings of mild traumatic brain injury in combat veterans with PTSD and a history of blast concussion. <i>J Neuropsychiatry Clin Neurosci.</i> 1998;10(3):308-313.
Wallace DM, Shafazand S, Ramos AR, et al. Insomnia characteristics and clinical correlates in Operation Enduring Freedom/Operation Iraqi Freedom veterans with post-traumatic stress disorder and mild traumatic brain injury: an exploratory study. <i>Sleep Med.</i> Oct 2011;12(9):850-859.
Wilkinson CW, Pagulayan KF, Petrie EC, et al. High prevalence of chronic pituitary and target-organ hormone abnormalities after blast-related mild traumatic brain injury. <i>Front Neurol.</i> 2012;3:11.
Williams JL, McDevitt-Murphy ME, Murphy JG, Crouse EM. Deployment risk factors and postdeployment health profiles associated with traumatic brain injury in heavy drinking veterans. <i>Mil Med.</i> Jul 2012;177(7):789-796.
Wolf GK, Strom TQ, Kehle SM, Eftekhari A. A preliminary examination of prolonged exposure therapy with Iraq and Afghanistan veterans with a diagnosis of posttraumatic stress disorder and mild to moderate traumatic brain injury. <i>J Head Trauma Rehabil.</i> Jan-Feb 2012;27(1):26-32.
Writer BW, Schillerstrom JE, Regwan HK, Harlan BS. Executive clock drawing correlates with performance-based functional status in people with combat-related mild traumatic brain injury and comorbid posttraumatic stress disorder. <i>J Rehabil Res Dev.</i> 2010;47(9):841-850.
Yurgelun-Todd DA, Bueler CE, McGlade EC, Churchwell JC, Brenner LA, Lopez-Larson MP. Neuroimaging correlates of traumatic brain injury and suicidal behavior. <i>J Head Trauma Rehabil.</i> Jul-Aug 2011;26(4):276-289.
<i>Does not report outcomes that address key questions</i>
McCarroll JE, Gunderson C. 5-year study of incidence rates of hospitalized cases of head injuries in the US Army. <i>Neuroepidemiology.</i> 1990;9(6):296-305.
McDonald BC, Saykin AJ, McAllister TW. Functional MRI of mild traumatic brain injury (mTBI): progress and perspectives from the first decade of studies. <i>Brain imaging behav.</i> Jun 2012;6(2):193-207.

Schwab KA, Ivins B, Cramer G, et al. Screening for traumatic brain injury in troops returning from deployment in Afghanistan and Iraq: Initial investigation of the usefulness of a short screening tool for traumatic brain injury. <i>The Journal of Head Trauma Rehabilitation</i> . Nov-Dec 2007;22(6):377-389.
Does not meet VA DoD definition for mTBI
Adams RS, Larson MJ, Corrigan JD, Horgan CM, Williams TV. Frequent binge drinking after combat-acquired traumatic brain injury among active duty military personnel with a past year combat deployment. <i>J Head Trauma Rehabil</i> . Sep 2012;27(5):349-360.
Arbisi PA, Polusny MA, Erbes CR, Thuras P, Reddy MK. The Minnesota Multiphasic Personality Inventory-2 Restructured Form in National Guard soldiers screening positive for posttraumatic stress disorder and mild traumatic brain injury. <i>Psychol Assess</i> . Mar 2011;23(1):203-214.
Armistead-Jehle P. Symptom validity test performance in U.S. veterans referred for evaluation of mild TBI. <i>Appl Neuropsychol</i> . Jan 2010;17(1):52-59.
Bazarian JJ, Zhu T, Blyth B, Borrino A, Zhong J. Subject-specific changes in brain white matter on diffusion tensor imaging after sports-related concussion. <i>Magn Reson Imaging</i> . Feb 2012;30(2):171-180.
Booth-Kewley S, Highfill-McRoy RM, Larson GE, Garland CF, Gaskin TA. Anxiety and depression in marines sent to war in Iraq and Afghanistan. <i>J Nerv Ment Dis</i> . Sep 2012;200(9):749-757.
Brenner LA, Ivins BJ, Schwab K, et al. Traumatic brain injury, posttraumatic stress disorder, and postconcussive symptom reporting among troops returning from Iraq. <i>J Head Trauma Rehabil</i> . Sep-Oct 2010;25(5):307-312.
Brenner LA, Terrio H, Homaifar BY, et al. Neuropsychological test performance in soldiers with blast-related mild TBI. <i>Neuropsychology</i> . Mar 2010;24(2):160-167.
Cameron KL, Marshall SW, Sturdivant RX, Lincoln AE. Trends in the incidence of physician-diagnosed mild traumatic brain injury among active duty U.S. military personnel between 1997 and 2007. <i>J Neurotrauma</i> . May 2012;29(7):1313-1321.
Clement PF, Kennedy JE. Wechsler Adult Intelligence Scale-third edition characteristics of a military traumatic brain injury sample. <i>Mil Med</i> . Dec 2003;168(12):1025-1028.
Dougherty AL, MacGregor AJ, Han PP, Heltemes KJ, Galarneau MR. Visual dysfunction following blast-related traumatic brain injury from the battlefield. <i>Brain Inj</i> . 2011;25(1):8-13.
Drake AI, Gray N, Yoder S, Pramuka M, Llewellyn M. Factors predicting return to work following mild traumatic brain injury: A discriminant analysis. <i>The Journal of Head Trauma Rehabilitation</i> . Oct 2000;15(5):1103-1112.
Eskridge SL. Combat-related blast injuries: Injury types and outcomes. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering</i> . 2012;72(7-B):3929.
Fear N, Jones E, Groom M, et al. Symptoms of post-concussional syndrome are nonspecifically related to mild traumatic brain injury in UK Armed Forces personnel on return from deployment in Iraq: An analysis of self-reported data. <i>Psychological Medicine</i> . Aug 2009;39(8):1379-1387.
Ferrier-Auerbach AG, Erbes CR, Polusny MA, Rath CM, Sponheim SR. Predictors of emotional distress reported by soldiers in the combat zone. <i>J Psychiatr Res</i> . May 2010;44(7):470-476.
French LM, Lange RT, Iverson GL, Ivins B, Marshall K, Schwab K. Influence of bodily injuries on symptom reporting following uncomplicated mild traumatic brain injury in US military service members. <i>J Head Trauma Rehabil</i> . Jan-Feb 2012;27(1):63-74.

Gottshall KR, Gray NL, Drake AI, Tejidor R, Hoffer ME, McDonald EC. To investigate the influence of acute vestibular impairment following mild traumatic brain injury on subsequent ability to remain on activity duty 12 months later. <i>Mil Med.</i> Aug 2007;172(8):852-857.
Helfer TM, Jordan NN, Lee RB, Pietrusiak P, Cave K, Schairer K. Noise-induced hearing injury and comorbidities among postdeployment U.S. Army soldiers: April 2003-June 2009. <i>Am J Audiol.</i> Jun 2011;20(1):33-41.
Heltemes KJ, Dougherty AL, MacGregor AJ, Galarneau MR. Alcohol abuse disorders among U.S. service members with mild traumatic brain injury. <i>Mil Med.</i> Feb 2011;176(2):147-150.
Hoffer ME, Balaban C, Gottshall K, Balough BJ, Maddox MR, Penta JR. Blast exposure: vestibular consequences and associated characteristics. <i>Otol Neurotol.</i> Feb 2010;31(2):232-236.
Hoffer ME, Donaldson C, Gottshall KR, Balaban C, Balough BJ. Blunt and blast head trauma: different entities. <i>Int Tinnitus J.</i> 2009;15(2):115-118.
Hoge CW, McGurk D, Thomas JL, Cox AL, Engel CC, Castro CA. Mild traumatic brain injury in U.S. Soldiers returning from Iraq. <i>N Engl J Med.</i> Jan 31 2008;358(5):453-463.
Ivins BJ, Kane R, Schwab KA. Performance on the Automated Neuropsychological Assessment Metrics in a nonclinical sample of soldiers screened for mild TBI after returning from Iraq and Afghanistan: a descriptive analysis. <i>J Head Trauma Rehabil.</i> Jan-Feb 2009;24(1):24-31.
Ivins BJ, Schwab KA, Baker G, Warden DL. Hospital admissions associated with traumatic brain injury in the US Army during peacetime: 1990s trends. <i>Neuroepidemiology.</i> 2006;27(3):154-163.
Ivins BJ, Schwab KA, Warden D, et al. Traumatic brain injury in U.S. Army paratroopers: prevalence and character. <i>J Trauma.</i> Oct 2003;55(4):617-621.
Ivins BJ. Hospitalization associated with traumatic brain injury in the active duty US Army: 2000-2006. <i>NeuroRehabilitation.</i> 2010;26(3):199-212.
Lange RT, Brickell TA, French LM, et al. Neuropsychological Outcome from Uncomplicated Mild, Complicated Mild, and Moderate Traumatic Brain Injury in US Military Personnel. <i>Arch Clin Neuropsychol.</i> 2012;27(5):480-494.
Lange RT, Pancholi S, Bhagwat A, Anderson-Barnes V, French LM. Influence of poor effort on neuropsychological test performance in U.S. military personnel following mild traumatic brain injury. <i>J Clin Exp Neuropsychol.</i> 2012;34(5):453-466.
Lange RT, Pancholi S, Brickell TA, et al. Neuropsychological Outcome from Blast versus Non-blast: Mild Traumatic Brain Injury in U.S. Military Service Members. <i>J Int Neuropsychol Soc.</i> May 2012;18(3):595-605.
Lew HL, Garvert DW, Pogoda TK, et al. Auditory and visual impairments in patients with blast-related traumatic brain injury: Effect of dual sensory impairment on Functional Independence Measure. <i>J Rehabil Res Dev.</i> 2009;46(6):819-826.
Luis CA, Vanderploeg RD, Curtiss G. Predictors of postconcussion symptom complex in community dwelling male veterans. <i>J Int Neuropsychol Soc.</i> Nov 2003;9(7):1001-1015.
MacGregor AJ, Dougherty AL, Galarneau MR. Injury-specific correlates of combat-related traumatic brain injury in Operation Iraqi Freedom. <i>The Journal of Head Trauma Rehabilitation.</i> Jul-Aug 2011;26(4):312-318.
MacGregor AJ, Dougherty AL, Morrison RH, Quinn KH, Galarneau MR. Repeated concussion among U.S. military personnel during Operation Iraqi Freedom. <i>J Rehabil Res Dev.</i> 2011;48(10):1269-1278.
MacGregor AJ, Shaffer RA, Dougherty AL, et al. Prevalence and psychological correlates of traumatic brain injury in operation iraqi freedom. <i>J Head Trauma Rehabil.</i> Jan-Feb 2010;25(1):1-8.

Macgregor AJ. Physical injury and psychological outcomes among United States combat veterans. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering</i> . 2007;68(6-B):3665.
McGuire SA, Marsh RW, Sowin TW, Robinson AY. Aeromedical decision making and seizure risk after traumatic brain injury: longitudinal outcome. <i>Aviat Space Environ Med</i> . Feb 2012;83(2):140-143.
Mora AG, Ritenour AE, Wade CE, Holcomb JB, Blackburne LH, Gaylord KM. Posttraumatic stress disorder in combat casualties with burns sustaining primary blast and concussive injuries. <i>J Trauma</i> . Apr 2009;66(4 Suppl):S178-185.
Morgan M, Lockwood A, Steinke D, Schleenbaker R, Botts S. Pharmacotherapy regimens among patients with posttraumatic stress disorder and mild traumatic brain injury. <i>Psychiatric Services</i> . Feb 2012;63(2):182-185.
Nelson C, St, Weiser M, Gifford S, Gallimore J, Morningstar A. Knowledge gained from the Traumatic Brain Injury Screen-Implications for treating Canadian military personnel. <i>Mil Med</i> . Feb 2011;176(2):156-160.
Olson-Madden JH, Forster JE, Huggins J, Schneider A. Psychiatric diagnoses, mental health utilization, high-risk behaviors, and self-directed violence among veterans with comorbid history of traumatic brain injury and substance use disorders. <i>J Head Trauma Rehabil</i> . Sep 2012;27(5):370-378.
Ommaya AK, Ommaya AK, Dannenberg AL, Salazar AM. Causation, incidence, and costs of traumatic brain injury in the U.S. military medical system. <i>J Trauma</i> . Feb 1996;40(2):211-217.
Ommaya AK, Salazar AM, Dannenberg AL, Chervinsky AB, Schwab K. Outcome after traumatic brain injury in the U.S. military medical system. <i>J Trauma</i> . Dec 1996;41(6):972-975.
Pietrzak RH, Johnson DC, Goldstein MB, Malley JC, Southwick SM. Posttraumatic stress disorder mediates the relationship between mild traumatic brain injury and health and psychosocial functioning in veterans of Operations Enduring Freedom and Iraqi Freedom. <i>J Nerv Ment Dis</i> . Oct 2009;197(10):748-753.
Plassman BL, Havlik RJ, Steffens DC, et al. Documented head injury in early adulthood and risk of Alzheimer's disease and other dementias. <i>Neurology</i> . Oct 24 2000;55(8):1158-1166.
Polusny MA, Kehle SM, Nelson NW, Erbes CR, Arbisi PA, Thuras P. Longitudinal effects of mild traumatic brain injury and posttraumatic stress disorder comorbidity on postdeployment outcomes in national guard soldiers deployed to Iraq. <i>Arch Gen Psychiatry</i> . Jan 2011;68(1):79-89.
Roebuck-Spencer TM, Vincent AS, Twillie DA, et al. Cognitive change associated with self-reported mild traumatic brain injury sustained during the OEF/OIF conflicts. <i>The Clinical Neuropsychologist</i> . Apr 2012;26(3):473-489.
Romesser J, Shen S, Reblin M, et al. A preliminary study of the effect of a diagnosis of concussion on PTSD symptoms and other psychiatric variables at the time of treatment seeking among veterans. <i>Mil Med</i> . Mar 2011;176(3):246-252.
Rona RJ, Jones M, Fear NT, et al. Mild traumatic brain injury in UK military personnel returning from Afghanistan and Iraq: cohort and cross-sectional analyses. <i>J Head Trauma Rehabil</i> . Jan-Feb 2012;27(1):33-44.
Rona RJ, Jones M, Fear NT, Sundin J, Hull L, Wessely S. Frequency of mild traumatic brain injury in Iraq and Afghanistan: are we measuring incidence or prevalence? <i>J Head Trauma Rehabil</i> . Jan-Feb 2012;27(1):75-82.
Schneiderman AI, Braver ER, Kang HK. Understanding sequelae of injury mechanisms and mild traumatic brain injury incurred during the conflicts in Iraq and Afghanistan: persistent postconcussive symptoms and posttraumatic stress disorder. <i>Am J Epidemiol</i> . Jun 15 2008;167(12):1446-1452.

Skopp NA, Trofimovich L, Grimes J, Oetjen-Gerdes L, Gahm GA. Relations between suicide and traumatic brain injury, psychiatric diagnoses, and relationship problems, active component, U.S. Armed Forces, 2001-2009. <i>Msmr</i> . Feb 2012;19(2):7-11.
Theeler BJ, Flynn FG, Erickson JC. Chronic Daily Headache in U.S. Soldiers After Concussion. <i>Headache</i> . 2012;52(5):732-738.
Theeler BJ, Flynn FG, Erickson JC. Headaches after concussion in US soldiers returning from Iraq or Afghanistan. <i>Headache</i> . Sep 2010;50(8):1262-1272.
Vanderploeg RD, Belanger HG, Curtiss G. Mild traumatic brain injury and posttraumatic stress disorder and their associations with health symptoms. <i>Arch Phys Med Rehabil</i> . Jul 2009;90(7):1084-1093.
Vanderploeg RD, Curtiss G, Belanger HG. Long-term neuropsychological outcomes following mild traumatic brain injury. <i>J Int Neuropsychol Soc</i> . May 2005;11(3):228-236.
Vanderploeg RD, Curtiss G, Duchnick JJ, Luis CA. Demographic, medical, and psychiatric factors in work and marital status after mild head injury. <i>J Head Trauma Rehabil</i> . Mar-Apr 2003;18(2):148-163.
Vasterling JJ, Brailey K, Proctor SP, Kane R, Heeren T, Franz M. Neuropsychological outcomes of mild traumatic brain injury, post-traumatic stress disorder and depression in Iraq-deployed US Army soldiers. <i>Br J Psychiatry</i> . Sep 2012;201:186-192.
Wilk JE, Herrell RK, Wynn GH, Riviere LA, Hoge CW. Mild traumatic brain injury (concussion), posttraumatic stress disorder, and depression in U.S. soldiers involved in combat deployments: association with postdeployment symptoms. <i>Psychosom Med</i> . Apr 2012;74(3):249-257.
Wilk JE, Thomas JL, McGurk DM, Riviere LA, Castro CA, Hoge CW. Mild traumatic brain injury (concussion) during combat: lack of association of blast mechanism with persistent postconcussive symptoms. <i>J Head Trauma Rehabil</i> . Jan-Feb 2010;25(1):9-14.
Yurkiewicz IR, Lappan CM, Neely ET, et al. Outcomes from a US military neurology and traumatic brain injury telemedicine program. <i>Neurology</i> . Sep 18 2012;79(12):1237-1243.

APPENDIX G. PEER REVIEW COMMENTS AND RESPONSES

Reviewer	Comment	Response
Question 1: Are the objectives, scope, and methods for this review clearly described?		
1	No	Noted.
1	Suggest adding in the background of Executive Summary more rationale (e.g., purpose is understanding OEF/OIF cohort vis-à-vis mTBI for planning etc).	We have made this addition.
1	Also suggest adding the criteria that were used for inclusion/exclusion at the beginning of the document.	Due to space limitations, we have left reference to inclusion/exclusion criteria in the body of the report and appendices; however, this information will be presented earlier in the planned article publication of the results.
1	Suggest being more specific when mentioning comparison to controls (e.g., on page 4, were they injured controls, postdeployed controls?)	We have included this information in the data abstraction tables.
2	No	Noted.
2	Objectives of the review are clearly described. The scope of this review is clearly described.	Noted.
2	The methods lack detail. Expectations for reporting methodological detail have grown exponentially over the past few years, and although this would expand the methods section, I highly recommend that this be done.	Noted. We have expanded this discussion.
2	Specifically, I recommend that the report follow the most recent PRISMA guidelines for reporting systematic reviews. I recognize that this is not primarily an academic document. However, adherence to the PRISMA guidelines would enhance the credibility of the review.	Noted. We have expanded our reporting to be in line with PRISMA guidelines as you suggest.
2	1. I assume that no online protocol was published, but if so, that should be reported.	We have updated this information in the report.
2	2. Eligibility criteria: The study designs included were mentioned. I have a few questions. Case control studies are not listed in the inclusion criteria. Were case control studies excluded from the search? I assume that case series and case reports were excluded, but this is not explicit. Under the criteria “Timing: No limitations based on timing” – my question – timing of what? Please clarify	Noted. Case control studies were not excluded. Timing has been updated to reflect time since injury. The inclusion criteria have been updated accordingly.
2	3. Information Sources: This is well covered. However, if study authors were contacted for additional information, this information should be included. If they were not contacted, it is fine to leave the section as it is.	Correct – no authors were contacted for additional information about studies included in this review.

Reviewer	Comment	Response
2	4. Search: The electronic search is included in appendix A. This is very good. Also, it has become standard practice to clarify that the search was developed by a library scientist experienced in database searches of this sort. It has also recently become recommended practice for a search strategy to be peer reviewed by a second librarian. Was this done? What measures were taken to ensure that the search was comprehensive?	We have added this description of procedures.
2	5. Study selection: This section is nicely detailed. Including the study selection form is very useful. Other information to consider adding: What procedures were taken where the PI and the other reviewer disagreed on relevance at the point of abstract screening? What procedures were undertaken where the reviewers disagreed on relevance at the point of full article screening? A detailed definition of TBI is provided in Appendix C. This requires a Mild TBI to have normal CT/MRI. It might be noted in the methods section (or someplace else in the document) that this is not a universally agreed upon criteria – many definitions of MTBI allow for abnormalities on imaging, differentiating these by classifying them as complicated or uncomplicated mild TBI. Of course it is important to adhere to the VA/DoD criteria, but it is a possible point of discrepancy from other criteria that should be explicitly noted.	Thank you. We have included this additional information in the updated report.
2	6. Data abstraction: Good tables. It would be useful to have details about the data abstraction process. It has become the standard to have at least two independent data abstractors – was this done? Who did the data abstraction? How was this checked? Again, where data were missing or ambiguous in the report, were authors contacted? It would also be helpful to include study design in the tables.	We have updated the report with this additional information. We chose not to include study design in the tables because the designs were often inaccurately reported in the published studies and because all studies were observational in nature. Specific study design criteria related to quality/potential for bias were abstracted in the tables (e.g., sample selection, comparison group, etc.)
2	7. Quality assessment: In this section, you state that case control and case series designs were included. This should be consistent with the statement of study design inclusion criteria. More details should be provided. Who did the quality assessment? It is standard practice to have two independent quality assessments on each paper, and to report the procedures undertaken when there is no consensus on quality. How (specifically) was the assessment of quality used in the data synthesis process? How specifically was this linked to rating the body of evidence? Who performed this linkage?	This discrepancy has been corrected. We have added this information.
2	8. Synthesis: You have provided a reasonable way of grouping the studies and no meta-analysis was conducted – I see this decision as appropriate. However, details about how the synthesis was conducted would be very good to add. This includes who conducted the synthesis, was this discussed in a larger group, were the conclusions agreed to by the working group, and what processes were used to reach agreement.	We have added this information.

Reviewer	Comment	Response
2	9. Results: I see a literature flow chart. This is very good.	Noted.
2	10. You provide well-formulated summary statements, and in your detailed results sections, you link findings to your citations. But it might also be useful to link findings to the particular studies in your summary statements as well. This is optional, but I present it as worth considering.	Due to the lengthiness of the report, we decided not to link findings to particular studies in the summary section. However, we linked findings in the results section to the citations of included studies should readers be interested in these specifics.
2	11. There is no linkage between particular studies and their study quality assessment – or more importantly, their assessed risk of bias. This does not necessarily imply the need for an overall score, and the use of an overall additive score has been widely criticized. But the reader should be able to see which studies have risk of which biases. There is some reference to overall study quality in the sections reporting summary of findings, but most recent standards recommend that this be reported in a more study-specific way. When you say “low quality” – what specifically do you mean? I also wonder whether studies with high risk of bias in their methodological quality can usefully contribute to our knowledge of TBI in the military?	We have updated our description of study quality and risk of bias to specifically describe each study as being of low quality due to the high risk of bias. We have also provided additional cautionary, interpretive statements in the summary section related to the strength of evidence from a body of low quality studies.
2	12. Were cross-sectional studies included in sections related to risk or prognostic factors. There is no information provided about the restriction of study designs for particular questions. Obviously, cross-sectional studies are appropriate for questions about prevalence of symptoms, but not for making causal inferences about these symptoms. Could this be clarified in the report?	We have clarified the inclusion of various types of study designs, highlighted findings from the only prospective study, and noted limitations of interpretation based solely on cross sectional studies reporting association rather than causal inferences.
2	13. The authors have done a good job in discussing limitations of the literature in the second last section. Would it be useful to be even more specific about biases in the Limitations and Recommendations section? E.g., providing concrete examples of recall bias; of incidence-prevalence bias (where it might exist), etc.	We have added additional discussion of these points.
3	Yes	Noted.
4	No	Noted.
4	1 The methods section of the Executive Summary is missing important information about inclusion/exclusion criteria and how quality and strength of evidence is rated. This latter point is particularly important as it makes it very difficult for the reader to understand why the authors consider the strength of the evidence to be low.	We have added this information to the executive summary section of the report.
4	It is also not clear when the authors state that no clear pattern of risk and protective factors emerge whether the studies were designed to look at risk and protective factors.	We have noted this in the discussion.

Reviewer	Comment	Response
4	2. In the Executive Summary Conclusions section (and elsewhere in the report) the authors refer to “objective results” (p.4). It would be helpful if they discussed which results were “objective” and what is meant by “objective”.	We have updated this section to describe results not based on self-report.
4	3. It is not clear to me why the authors are including the headache intervention study. How is participating in a headache intervention a protective factor? What then is meant by protective factor? Do the authors mean that the sample referred for headache intervention was different than those not referred? Why is that relevant to the KQs? In general, the information presented on this particular study is confusing throughout the report and relevance is not clear.	We have clarified references to this intervention throughout the report.
4	4. It is not clear to me why studies focused on biomarkers are included. I do not see how biomarkers are “impairments” (KQ 1) according to any definition of impairment. I do not see how biomarkers address either KQ2A (pre-injury factors) or 2B (post injury factors) if they not include a focus on outcomes, given the way that KQ is worded. How does this study meet criteria for the outcomes listed on p.10?	We agree and have moved information from this section to the discussion rather than including it in the evidence synthesis.
4	5. Table 1: I find the mTBI definition column confusing. What do the authors mean by “citation”. What do the authors mean when they list “LOC, AOC and PTA” given inclusion criteria requires DOD/VA/ACRM definition of mTBI? A note to the table may help clarify.	We have clarified this column label.
4	6. I did not finding tables summarizing strength of evidence per study. This information is particularly important given the conclusions. It would also make it easier for the reader to refer to specific studies when reviewing the report. In the absence of this information, it is difficult to interpret statements like, “Strength of the evidence was low because of”. If I missed something that was included in the report, I apologize.	We have clarified this information in the text of the report. All included studies were rated as low quality due to high risk of bias, without exception.
4	7. Tables 2–6: I would have liked to see the pertinent studies referenced rather than or at least in addition to the number of studies listed in parentheses. Number may not be as important as quality, in my opinion. Question based on Table 2: Did <u>4</u> studies find that mean scores for processing speed were within normal limits with possible exceptions of those getting C&Ps and <10 days since injury (p.24)? This is what the Table suggests, unless I am not reading it correctly.	We have referenced individual citations in the text, and due to the low quality rating of all included studies, have not identified individual studies or study quality in the table for reasons of space. You are correct in your interpretation of the presented information in the table.

Reviewer	Comment	Response
4	8. When the authors make statements like “most research reported no significant risk or protective factors,” it would be helpful to know the number/percent of studies were looking at risk/protective factors.	We have noted this information in the discussion as an overall commentary on the body of literature given that only one included study was prospective in nature, and the rest were not designed to assess risk/protective factors. We have also clarified these sentences in the report to reflect studies of association rather than implying that studies were designed to assess risk/protective factors.
4	9. I do not know that readers who are not neuropsychologists will understand how the authors are distinguishing between cognitive and physical health problems. For example, it may not be clear why visuospatial abilities falls in the cognitive domain but vision is in the physical health results section. It is not clear why the authors review “effort and motivation” at all. It is not clear why studies focused on sleep are in the functional/social outcomes section. I suggest that the authors explicitly describe the rationale for their groupings.	We have provided additional description of these decisions in the report.
4	10. In the Executive Summary Limitation section, the authors state that the studies included in the review relied on well-validated assessment tools. Are they classifying the NSI as well-validated? Perhaps this statement can be more precise.	We have made this statement more accurate and non-specific to individual tools. We refer to well-validated assessment tools being a strength of the overall body of literature, and report specific tools in the data abstraction. However, we did not to an individual literature search or other method of assessment to determine validation of each tool used in the included studies. We do also note limitations of some of the tools (e.g., the NSI) used to assess single-item, self-report outcomes.
4	11. Service Utilization and Costs: For this section, it does not make sense to me to collapse across Veteran and active duty samples given differences in the healthcare systems. I recommend that the authors clearly state which findings are specific to Veterans using VA and which are specific to active duty.	We have added a clarifying statement noting that all included studies investigated Veterans.
5	Yes	Noted.
6	Yes	Noted.
6	The objectives mention cognitive “disability” as an outcome, but based on the variables examined in the literature, cognitive “deficits” may be a better term. The language regarding mental health outcomes could also be clarified. The objectives mention “symptoms” but the operational definition is later given as diagnoses. However, examination of the measures in the studies included suggests that “symptoms” would be the more accurate term.	We have changed the wording to “deficits.”
7	Yes	Noted.

Reviewer	Comment	Response
7	These are well described.	Noted.
8	Yes	Noted.
8	Methods and objectives clearly described, but missing some information. There was no discussion of how common features of mTBI, e.g. the presence of comorbidities and multiple TBIs, were handled in inclusion criteria.	We have clarified these criteria in the report.
8	“Study relevance” was a major factor in exclusion (p.14) but no further information provided on how this criterion was applied.	Study relevance simply implies that the study must provide information included in the KQs and meet inclusion/exclusion criteria. No additional relevance criteria were applied.
8	Overall weakness in discussion of physical and neuroimaging outcomes. The scope as defined by the key questions does not include imaging or biomarkers as outcomes. It is not clear whether these topics were adequately searched according to the search strategy presented. However, I am not familiar with the abbreviations and formatting used to detail the search strategy, so this was a bit hard to follow. An explanation or reference would be helpful.	We agree and have moved the imaging/biomarker information to the discussion section and removed it from the results/evidence synthesis.
8	As a general comment, scope is very large and each element (e.g. cognitive effects) could be the subject of a report.	Noted.
9	Yes	Noted.
9	The key objectives and methodology are sound.	Noted.
10	Yes	Noted.
10	Consider rephrasing to: Key Question # 1: For Veterans/Service Members who suffer a mTBI and develop acute and or persistent sequale of mTBI symptoms what is the prevalence of health conditions (e.g. pain, headaches, insomnia, vertigo, or seizure disorder), functional limitations, (e.g. return to work/duty, marital status/family dynamics), cognitive impairment (e.g. attention, concentration or memory) and or associated mental health conditions (e.g. PTSD, depression or anxiety disorder). Key Question # 2: What factors affect outcomes for Veterans/service members with mTBI? Key Question 2A: For Veteran/military populations, are there pre-injury (premorbid) risk factors (e.g., pre-injury mental health factors, genetic factors, or prior concussions) or protective factors ??) that affect outcomes for mTBI? Key Question 2B: For Veteran/military populations, are there post-injury risk factors (e.g., PTSD, depression or anxiety) or protective factors that affect outcomes for mTBI?	We have changed the wording of the key questions slightly.
11	Yes	Noted.

Reviewer	Comment	Response
2. Is there any indication of bias in our synthesis of the evidence?		
1	No	Noted.
2	No	Noted.
2	I don't see an indication of bias, although it is always useful to discuss possible publication bias and how this might impact on the synthesis. However, a synthesis based on highly biased studies can bias a synthesis. Where studies were highly biased (low quality), can it be estimated what direction those methodological flaws would have biased the study's findings? (toward or away from the null, for example). This may impact on the interpretation of study findings.	Noted. We have added these points to the discussion.
3	No	Noted.
4	Yes	Noted.
4	1. The report states that TBI is THE leading cause of morbidity and disability in OEF/OIF. I do not believe that this is accurate. More important, one needs to distinguish between TBI history and TBI-related disability. The prevalence of TBI-related disability is unknown and some evidence suggests that persistent problems in many individuals who suffered TBI in OEF/OIF result from mental health comorbidities	Noted. We have changed the introductory paragraphs to better reflect these distinctions.
4	2. In Executive Summary Conclusions section, the authors state that "It is likely that the prevalence....is largely influenced by factors other than deployment rather than being uniquely associated with mTBI." They do NOT however describe a rationale for this conclusion. This reviewer suggests that the authors clearly build conclusions based on the literature reviewed otherwise the sentence reads like an opinion. The statement in the Executive Summary Conclusions that reads, "... the most likely exceptions are...evaluation linked with potential compensation" is particularly troubling as it seems to be based on the one study that looked at compensation effects (#34). In the Executive Summary Limitation section the authors state "...self-reported deficits are more likely to persist for individuals with mTBI particularly when associated with compensation (p.40." Again, this statement seems too strong if only one study (#34) reviewed looked at compensation effects unless that study was of very high quality	These conclusions are based on consistent findings across Veteran/military and civilian literature, though we agree that the conclusions could have been interpreted as basing conclusions on solely the Veteran/military literature, which would not have been warranted. These concluding statements have been tempered and clarified to more accurately reflect the results available from the body of literature.

Reviewer	Comment	Response
4	3. In the Summary and Discussion section the authors conclude, “The body of research ...suggest that many health consequences resolve within the first few months following injury, if not sooner” (p.49). I am not clear about what evidence they are using to draw that conclusion given the design of the studies reviewed. To which specific studies/findings are the authors referring when they state, “Objective cognitive impairment most often resolves within a few weeks of initial injury.”? I suggest that authors be very clear about the findings and studies they are using to make that claim. That claim also seems to be in direct contradiction to their statement in the Background section that “TBI is the leading cause of morbidity and disability...” It also contradicts the statement in the Service Utilization/Costs that “The long-term resource needs of recent Iraq and Afghanistan War Veterans who sustained mTBI are likely substantial”. (p.52). Why would that be the case?	We have clarified the studies on which these conclusions are based, as sometimes the quotes were in direct reference to the Carroll et al, 2004 review findings that we were summarizing. We have also tempered the introductory statements relating to the potential effects of TBI.
4	3. The authors also state in Summary and Discussion, “This report documents that litigation or evaluation for compensation as being a risk factor...(p.49)” Is this the finding based on one study referred to in multiple sections of this report (#34) or are there other studies? The basis for this statement was not clear to me.	This statement was in reference to findings from the Carroll et al, 2004 review, and this citation has been clarified.
5	Yes	Noted.
5	The bias has to do with studies included versus excluded. Your criteria are clear and stated on the bottom of page 9, of which the DoD/VA criteria are one operational definition of those criteria.	Noted.
5	However, the manuscript <u>includes</u> studies that don’t meet that criteria because they:	Noted.
5	(a) include moderates and “unclassified” severity patients [Morrisette, Woodward, Kimbrel, et al, 2011 ⁷⁵ , Schiehser, Delis, Filoteo, et al., 2011 ³⁵] or,	Thank you for catching this error. Though this paper reports findings for the mTBI only group separately from those with moderate or unclassified TBI, the specific finding we reported in our review was a combined group finding; therefore, this study has been excluded.
5	(b) includes those where TBI was not verified (“probably TBI) [Ruff, Riechers, Wang, et al., 2012 ³²], or	Though this paper reports some findings for a combined TBI group, the findings that we reported in this review are reported separately for the mTBI only group.
5	(c) at least as currently written in the tables, state criteria at variance to the DoD/VA criteria such as the GCS > 13 was used or GCS = 13 (when it should be ≥ 13) [Gaylord, Cooper, Mercado, et al., 2008 ²¹ , Cooper, Mercado-Couch, Richfield, et al., 2010 ¹⁸].	You are correct that this was inaccurately stated in the table. It should be ≥ 13, as reported in the article, and therefore these studies remain included.

Reviewer	Comment	Response
5	Studies are <u>excluded</u> , that in my view meet the DoD/VA criteria at least as well, if not better, than some of the included studies. Of course I am biased, but I'm referring to my studies:	Noted. We have reviewed all the suggested studies and agree that they provide useful information; however, we have scoped the review to include a specific subset of papers meeting VA/DoD mTBI criteria, and the papers you suggest do not fit within those pre-specified criteria and are therefore not included in the review.
5	Luis CA, Vanderploeg RD, Curtiss G. Predictors of postconcussion symptom complex in community dwelling male veterans. <i>J Int Neuropsychol Soc.</i> Nov 2003;9(7):1001-1015.	Reviewed, not included due to not meeting inclusion criteria.
5	Vanderploeg RD, Belanger HG, Curtiss G. Mild traumatic brain injury and posttraumatic stress disorder and their associations with health symptoms. <i>Arch Phys Med Rehabil.</i> Jul 2009;90(7):1084-1093.	Reviewed, not included due to not meeting inclusion criteria.
5	Vanderploeg RD, Curtiss G, Belanger HG. Long-term neuropsychological outcomes following mild traumatic brain injury. <i>J Int Neuropsychol Soc.</i> May 2005;11(3):228-236.	Reviewed, not included due to not meeting inclusion criteria.
5	Vanderploeg RD, Curtiss G, Duchnick JJ, Luis CA. Demographic, medical, and psychiatric factors in work and marital status after mild head injury. <i>J Head Trauma Rehabil.</i> Mar-Apr 2003;18(2):148-163.	Reviewed, not included due to not meeting inclusion criteria.
5	Self-report for LOC is not reliable because individuals do not know if they actually had an LOC or simply a memory gap (i.e., PTA). Our studies used alteration of consciousness defined as "loss consciousness or 'black out'". That is, either a self-reported LOC or a self-reported Alteration of consciousness. It is possible, although unlikely, that my studies included a few folks who had moderate injuries. Unlikely because the data was collected in the 1970s at which time those with anything other than a mild TBI would be hospitalized overnight at least, and no subject was hospitalized. In addition, the bias come in because other studies were included that had moderate or "unclassified" TBI severity subjects.	See above comments; we have re-reviewed all noted studies and the inclusion/exclusion criteria have been appropriately applied. Thank you for noting the possible discrepancies as one of the studies was inappropriately included in the first draft of the report and has now been excluded (Morrisette).
5	None of this would change findings, but my studies do address things the manuscripts says have not been addressed – frequencies of different medical signs and symptoms and psychosocial outcomes, as well as frequencies of neuropsychological impairments, in addition to comparison with an injury control group (groups most other studies do not have) and controlling for comorbid or premorbid medical and mental health conditions.	See above response re: included studies.

Reviewer	Comment	Response
5	<p>In addition, core findings of these studies have been replicated in a new sample using criteria that you would likely agree does meet the DoD/VA criteria for mild TBI. This study, also not included, is: Vanderploeg, R.D., Belanger, H.G., Horner, R.D., Spehar, A.M., Powell-Cope, G., Luther, S.L., Scott, S.G., (2012). Health Outcomes Associated With Military Deployment: Mild Traumatic Brain Injury, Blast, Trauma, and Combat Associations in the Florida National Guard. <i>Archives of Physical Medicine and Rehabilitation</i>, <i>93</i>, 1887-1895.</p>	<p>We have reviewed this study and it does not quite meet VA/DoD criteria, and therefore is not included in the review.</p>
6	No	Noted.
6	<p>The selection of studies is straightforward, but given the large number excluded, it might be helpful to summarize in the table of excluded studies, the specific reasons for exclusion. Without knowing how close these studies came to being eligible for inclusion in the paper, it's difficult to assess whether the eligibility criteria themselves may have incidentally introduced a bias. For example, are war-zone samples more likely to be excluded than veteran samples due to contextual constraints that somehow limited the information gathered? I'm not suggesting altering the eligibility criteria, but instead suggesting assessing potential "sampling biases" (for inclusion in the review) based on the possible identification of variables consistently associated with failure to use the DoD/VA definition.</p>	<p>We had similar concerns, and this was the rationale for including a table of studies meeting all inclusion criteria except for VA/DoD mTBI definition so that we were transparent about exclusions and readers could examine the list for possible bias in the included/excluded studies. We have also added a table of all the full-text study citations and exclusion codes.</p>
7	No	Noted.
7	<p>The review was absent of bias, and was appropriately critical of the lack of rigorous methodology, TBI severity description, and appropriate controls that appear pervasive throughout the literature on TBI in U.S. service members and Veterans.</p>	Noted.
8	Yes	Noted.
8	<p>Inclusion criteria introduced biases that should be discussed. Use of VA/DoD criteria for mTBI could time-limit the literature to after 2007; indeed, all included studies were from within the last 3 years. These criteria may also limit to military and VA-affiliated researchers and to US researchers. It is possible that exclusion of studies based on reporting of LOC and PTA led to unnecessary loss of data, as the exact value of these is usually based on self-report and is unreliable.</p>	<p>We agree that any scoping decisions, this one included, introduce potential for bias. This decision was agreed upon by stakeholders for this review in order to obtain the most accurate description of a specific population of interest: Members of the US military/Veterans with mTBI meeting VA/DoD criteria. Therefore, though the report is limited in these ways, the stakeholders agreed that other, broader reports (e.g., the Carroll et al, 2004 WHO mTBI report) could address broader/different questions. In response to your comment, we have broadened the discussion of this point in the discussion.</p>

Reviewer	Comment	Response
8	The requirement for an mTBI sample size of at least 30 will bias the type of outcome measures that are used. Because of expense and other limitations, few neuroimaging studies will fulfill this requirement. Large studies can be limited in the ability to perform in-depth testing, and so the report may be biased toward less sensitive questionnaire data and easier-to-administer testing protocols (e.g. RBANS is a screening measure designed for dementia). Large studies also may bias away from presentation of individual results, which as you note, can be informative. Few objective evaluations of physical outcomes (e.g. audiology) were included, which severely limits interpretation of this domain.	We agree, and have moved the neuroimaging information to the discussion section because of the likelihood that it is not comprehensive since we did not design the search to focus on these outcomes. We have also now added a table of excluded studies so that authors can review studies that were excluded based on sample size to gather additional information as needed.
8	Many studies served as sources across outcome domains (e.g. Nelson et al., 2012). This could potentially perpetuate any biases or limitations present in the single study across domains (e.g. recruitment setting, inadequate power).	We agree and have noted this limitation in the discussion.
9	No	Noted.
10	No	Noted.
11	No	Noted.
3. Are there any <u>published</u> or <u>unpublished</u> studies that we may have overlooked?		
1	Yes	Noted.
1	Cohen, Suri, Amick, & Yan, 2012 (published in Work)	This study does not meet our inclusion criteria because of unclearly reported definition of mTBI and because results are not reported separately for those with mTBI versus moderate/severe TBI.
2	No	Noted.
2	Not that I can think of.	Noted.
3	Yes	Noted.
3	Scholten et al, Analysis of US Veterans Health Administration comprehensive evaluations for traumatic brain injury in Operation Enduring Freedom and Operation Iraqi Freedom Veterans. Brain Inj 2012	We have reviewed this study and it does not meet inclusion criteria due to the mTBI definition used to define the cohort of participants.
4	Yes	Noted.
4	Have the authors reviewed the CBO report: The Veterans Health Administration’s Treatment of PTSD and Traumatic Brain Injury Among Recent Combat Veterans?	We have reviewed this report and agree that though it provides important information and guidance, it does not meet our criteria for inclusion in this review.
5	Yes	Noted.

Reviewer	Comment	Response
5	Vanderploeg, R.D., Belanger, H.G., Horner, R.D., Spehar, A.M., Powell-Cope, G., Luther, S.L., Scott, S.G., (2012). Health Outcomes Associated With Military Deployment: Mild Traumatic Brain Injury, Blast, Trauma, and Combat Associations in the Florida National Guard. <i>Archives of Physical Medicine and Rehabilitation</i> , <i>93</i> , 1887-1895.	We have reviewed this study and agree that though it provides important information, it does not meet VA/DoD mTBI definitional criteria and is therefore not included in this review.
6	Yes	Noted.
6	To my knowledge, the report captures all of the military/military veteran studies. However, in the discussion of meta-analytic studies from the civilian literature, it would be important to balance the discussion with the Pertab et al., meta-analysis that re-analyzes data from previous meta-analytic studies and reveals a potential qualification of prior findings.	This review has been added to the report.
7	Yes	Noted.
7	JRRD has recently come out with a TBI sensory and communications disorders edition (Vol. 49, Issue 7, 2012). Even though this journal is published after Oct. 3, 2012, these articles cover deployment-related experiences (e.g., blast, TBI), and associations with the following senses, conditions, and patterns: vestibular, visual, auditory, pain, PTSD, and referrals. This compendium fits well with the focus of the synthesis.	Thank you for this suggestion. We have reviewed all studies from this special issue none meet criteria for inclusion in this report.
7	Other published studies that were not included, though would be excluded for not meeting mTBI criteria, are: 1. Iverson, K. M., Hendricks, A., Kimerling, R., Kregel, M., Meterko, M., Stolzmann, K., Baker, E., Pogoda, T.K., Vasterling, J., & Lew, H.L. (2011). Psychiatric diagnoses and neurobehavioral symptom severity among OEF/OIF VA patients with deployment-related TBI. <i>Women's Health Issues</i> , <i>2(45)</i> , S210-S217.	We have reviewed this study and agree that though it provides important information, it does not meet VA/DoD criteria and is therefore not included in this review.
7	2. Lew, H.L., Kraft, M., Pogoda, T.K., Amick, M.M., Woods, P., & Cifu, D.X. (2011). Prevalence and Characteristics of Driving Difficulties in Operation Enduring Freedom/Operation Iraqi Freedom Combat Returnees. <i>JRRD</i> , <i>48(8)</i> , 913-926.	We have reviewed this study and agree that though it provides important information, it does not meet VA/DoD criteria and is therefore not included in this review.
7	3. Lew, H. L., Pogoda, T.K., Baker, E., Meterko, M., Stolzmann, K.L., Cifu, D.X., Amara, J.H. & Hendricks, A.M. (2011). Prevalence of dual sensory impairment and its association with traumatic brain injury and blast exposure in OEF/OIF Veterans. <i>Journal of Head Trauma & Rehabilitation</i> , <i>26(6)</i> :489-96.	We have reviewed this study and agree that though it provides important information, it does not meet VA/DoD criteria and is therefore not included in this review.
7	4. Hendricks AM, Amara J, Baker E, Charms MP, Gardner JA, Iverson KM, et al. (in press) Screening for mild traumatic brain injury in OEF-OIF deployed US military: an empirical assessment of VHA's experience. <i>Brain Injury</i>	We were not able to obtain a copy of this study for review.
8	Yes	Noted.

Reviewer	Comment	Response
8	Luethcke et al. 2010. Comparison of Concussive Symptoms, Cognitive Performance, and Psychological Symptoms Between Acute Blast-Versus Nonblast-Induced Mild Traumatic Brain Injury. <i>Journal of the International Neuropsychological Society</i> (2011), 17, 36–45.	We reviewed this study and it did not meet inclusion criteria.
8	Terrio, et al. 2009 Traumatic Brain Injury Screening: Preliminary Findings in a US Army Brigade Combat Team. <i>J Head Trauma Rehabil</i> Vol. 24, No. 1, pp. 14–23.	We reviewed this study and it did not meet inclusion criteria.
8	Caplan et al. 2010 The Structure of Postconcussive Symptoms in 3 US Military Samples. <i>J Head Trauma Rehabil</i> Vol. 25, No. 6, pp. 447–458.	We reviewed this study and it did not meet inclusion criteria.
8	Cockerham, 2009. Eye and visual function in traumatic brain injury. <i>J Rehab Research Dev</i> Volume 46, Number 6, 2009 Pages 811–818.	We reviewed this study and it did not meet inclusion criteria.
8	Akin and Murnane, 2011. Head Injury and Blast Exposure: Vestibular Consequences. <i>Otolaryngol Clin N Am</i> 44 (2011) 323–334.	We reviewed this study and it did not meet inclusion criteria.
8	Pogoda et al., 2012. Multisensory Impairment Reported by Veterans with and without Traumatic Brain Injury History. <i>J Rehab Research Dev</i> Volume 49, Number 7 Pages 971–984.	This study has been included in the report. Thank you for the suggestion.
8	Vasterling, et al. 2012. Neuropsychological outcomes of mild traumatic brain injury, post-traumatic stress disorder and depression in Iraq-deployed US Army soldiers <i>Br J Psychiatry</i> 201, 186-192	We reviewed this study and it did not meet inclusion criteria.
8	Schneibel et al., 2012 Altered brain activation in military personnel with one or more traumatic brain injuries following blast. J Int Neuropsychol Soc. 2012 Jan;18(1):89-100	We have moved the discussion of imaging and biomarkers to the discussion section of the report since it falls outside the scope of our key questions.
8	Morey et al., 2012 Effects of chronic mild traumatic brain injury on white matter integrity in Iraq and Afghanistan war veterans. Hum Brain Mapp. 2012 Jun 15	We have moved the discussion of imaging and biomarkers to the discussion section of the report since it falls outside the scope of our key questions.
8	Yurgelon-Todd, et al.,2011. Neuroimaging Correlates of Traumatic Brain Injury and Suicidal Behavior <i>J Head Trauma Rehabil</i> Vol. 26, No. 4, pp. 276–289	We reviewed this study and it did not meet inclusion criteria.
8	Sponheim, 2011. Evidence of disrupted functional connectivity in the brain after combat-related blast injury. <i>NeuroImage</i> 54 (2011) S21–S29	We have moved the discussion of imaging and biomarkers to the discussion section of the report since it falls outside the scope of our key questions.
8	Peskind et al., 2011. Cerebrocerebellar hypometabolism associated with repetitive blast exposure mild traumatic brain injury in 12 Iraq war Veterans with persistent post-concussive symptoms. <i>NeuroImage</i> 54 (2011) S76–S82.	We reviewed this study and it did not meet inclusion criteria.
9	Not aware of any that have been excluded that meet criteria for inclusion. Please see below regarding Vision data.	Noted.
10	Yes	Noted.

Reviewer	Comment	Response
10	Cooper DB et al. Relationship between mechanism of injury and neurocognitive functioning in OEF/OIF service members with mild traumatic brain injuries. Mil Med. 2012 Oct;177(10):1157-60.	This study has been included in the report. Thank you for the suggestion.
10	Scholten JD et al. Analysis of US Veterans Health Administration comprehensive evaluations for traumatic brain injury in Operation Enduring Freedom and Operation Iraqi Freedom Veterans. Brain Inj. 2012;26(10):1177-84.	We have reviewed this study and it does not meet inclusion criteria due to the mTBI definition used to define the cohort of participants.
10	Bryan CJ , et al. Loss of Consciousness, Depression, Posttraumatic Stress Disorder, and Suicide Risk Among Deployed Military Personnel With Mild Traumatic Brain Injury. J Head Trauma Rehabil. 2012 Oct 16.	This study has been added to the included studies for this report. Thank you for the suggestion.
11	Yes	Noted.
11	During an earlier call you had mentioned that another group (I believe IOM) was conducting a review of mTBI literature. It would be helpful to note this since non-Veteran studies were not included in the literature review and could have been aggregated to be the comparison to the Veteran based studies.	Agreed. The WHO group lead by Dr. Linda Carroll is updating the 2004 mTBI prognosis review, and we have included this information in our report for reader reference.
4. Please write additional suggestions or comments below. If applicable, please indicate the page and line numbers from the draft report.		
1	On page 26, please define “old learning.” More specificity in general would be helpful (so for example, for ‘memory studies’, were delayed recall, trial-by-trial learning, recognition all examined? On page 48 in the summary of results, to what type of control groups were mTBI compared? On page 53, suggest adding the number of mTBI’s as a future area for investigation. On page 54, suggest being judicious with the use of the word “persistent” since it is unclear if symptoms persist in the longitudinal sense.	We have changed “old learning” to be “language abilities and general fund of verbal knowledge.” We have included additional information on specific tests and on comparison groups in the appendix tables. We have added a statement in the discussion relating to future research of the number of TBIs. We have clarified and limited use of the term persistent throughout the report.
2	Page 49, Where other reviews are cited, it would be useful to explicitly indicate what year they were published, since early systematic reviews might simply be outdated. The study citation is included, of course, but inclusion of the year of publication in the body of this section would highlight that point.	We have included the year in text in this section.
3	Overall, an excellent evidence review.	Noted. Thank you.
3	It should be emphasized for all domains that major shortcomings in the literature are the lack of non-mTBI comparison groups and lack of adequate pre-mTBI (premorbid) data.	Agreed, and we have added this point to the discussion.
3	In section on mental disorders, some controlled civilian studies do show higher rates of mental disorders after mTBI compared with non-TBI controls (e.g., Fann et al, Arch Gen Psychiatry 2004). Presence of prior psychiatric conditions is a major risk factor. Data from this same cohort showed higher health care utilization among those with mTBI compared with non-TBI controls (Rockhill et al, J Neurotrauma 2012).	We have clarified that some civilian literature indicates higher rates of mental disorders after mTBI, as noted in the Carroll et al 2004 review.

Reviewer	Comment	Response
4	Below I provide specific comments about organization and note some areas where the terms used were not clear to me. Please disregard if not helpful in efforts to finalize the report.	Thank you for the suggestions. Some will be included in this report, though some will be included in the planned article publication rather than in this full report.
4	1. I suggest that the authors orient the reader at the beginning to their overall approach to summarizing the findings. It took this reviewer a few reads to understand how the authors were presenting the material. It was a bit confusing to me to read the summary before the sections describing findings per domain (i.e., summary of cognitive functioning in general before description in each cognitive domain). Sometimes the summaries seem completely unnecessary because only one study is reviewed. In general, there was considerable repetition because of the structure of this report, which made the report difficult to read. It takes several reads to know what is new information versus a restating of what has already been summarized.	Because some parts of this report are purposely repetitive (e.g., the executive summary), we plan to make the suggested change for the published article.
4	2. This reviewer suggests the authors state reason time since injury is important rather than assuming the reader already has this information.	Noted, and this has been added.
4	3. To help the reader understand the “cognitive function” section, I suggest the authors tell the reader that x# studies are based on neuropsychological testing and briefly explain. It would be helpful to have an appendix in which the function each test assesses is described – otherwise, I am not sure how helpful it is to list measures/acronyms in Appendix E. I did not find it useful to have the names of the tests listed in the text – in fact, I think it makes the narrative harder to read.	We have made some of these changes, including shortening the acronyms listed in the results section of the report.
4	4. In the Summary of Findings for Cognitive Function Results the author state that “standardized scores are scores associated with impairment below a certain cutoff.” This is not quite right and I do not think the sentence expresses what they intend it to.	We have corrected this sentence.
4	5. It is confusing when the authors state that risk factors include “LOC and PTA” or the like, given that they are using the DoD/VA/ACRM definition of mTBI. As opposed to what?	We have clarified that LOC and PTA were compared to just alteration of consciousness.
4	6. It would be more accurate on p.39 to state that the PCL scores suggest clinically significant <i>symptoms</i> rather than “impairment” as stated in the report	We have chosen to use the term impairment across measures to use similar terminology describing scores above a clinically significant cutoff.
4	7. The term “TBI sequelae” as used here is confusing. For example, on p. 28 the authors state that one study reported “...based on mTBI with LOC and PTA compared to those with mTBI who did not have these immediate sequelae.” What does this mean? If these individuals did not have these “sequelae”, how was mTBI determined? This is probably a matter of simple clarification.	We have changed this sentence to clarify.

Reviewer	Comment	Response
4	8. Summary and Conclusions: This section is largely devoted to discussion of the civilian literature. The subsection on Physical Health Outcomes (p.49) does not even discuss the Veteran/military studies. I suggest switching the focus so that the authors primarily discuss the literature they reviewed for this report.	Noted. We have updated the physical health section to focus the discussion on a comparison.
4	9. It seems to me that references are used inconsistently. Why are there no references for the summaries? Why are the “one study” referred to on p. 26 and the “single studies” on .34 not referenced What are the “some studies” (p.30 that found better cognitive function for those not evaluated in forensic settings? I thought there was just one study that examined this issue (#34). In some sections, it is very difficult to identify which studies are forming the basis of the authors’ conclusions.	Noted. We have made the suggested corrections to text. The format for this report is such that summaries do not contain citations; however, citations will be used consistently throughout the published article.
4	10. Note that PTSD is an AXIS I disorder (p.35). I do not understand the statement, “Finally, though many individual studies...general association between specific mental health outcomes with other mental health diagnoses and symptoms”.	We have clarified that some studies looked at only PTSD, some examined “any Axis I disorder” which would include PTSD. We have clarified the confusing sentence.
5	1. A statement is made without any supporting reference, that I do not believe the literature supports. Page 1: “TBI is the leading cause of morbidity and disability among OEF/OIF service members.” I’m not sure about “morbidity” (but I think chronic pain and mental health are higher), but I’m quite sure there is no evidence to support the “disability” claim. This statement is repeated several times throughout the manuscript.	This has been corrected throughout the manuscript.
5	2. Page 2 bottom (and elsewhere) “One or more studies have found . . . [problems in those] experiencing loss or alteration of consciousness at the time of the injury.” It seems to me that everyone included in every study meets this criteria because that is the criteria for TBI. So, that statement it seems problematic. If all studies had all subjects with an immediate event-related “loss or alteration of consciousness” how can “loss or alteration of consciousness at the time of injury” be a unique factor?	We have clarified that this refers only to patients who have PTA but not LOC or AOC, as described by the primary study authors.
5	3. Page 3 (an elsewhere): If you include my studies (or the one you inadvertently omitted because it was recently published), the statement “Similar to objective cognitive results, prevalence of self-reported cognitive deficits was not reported in the included studies” would be inaccurate.	We have updated all summary statements to reflect the final list of articles meeting inclusion criteria.

Reviewer	Comment	Response
5	4. Page 5: The statement “. . . self-reported deficits are more likely to persist for individuals with mTBI” is inaccurate. There is a difference between “persistence of symptoms” and “symptoms reported in the chronic phase” which may come-and-go or wax-and-wane. Studies have not demonstrated persistence. What they have demonstrated is problems/symptoms reported in the post-acute or chronic phase. The studies do not document that these began at the time of the mTBI and persisted over time to the time of assessment. Other civilian literature clearly documents that symptoms and problems are not persistent, but rather that they come-and-go or wax-and-wane, but are generally higher in frequency in mild TBI subjects. We don’t know if they were also higher prior to the mild TBI but we do know that they are not persistent.	We have made this change throughout the report.
5	5. The term “Language and Old Learning” is a term no one uses, and as a result is confusing. Initially I thought you were referring to “Verbal Learning and Memory”. I would suggest using the term “Language Abilities and General Fund of Verbal Knowledge” or something like that.	Thank you for this suggestion. We have made this change.
5	6. Throughout the review you refer to “statistical significance” or lack thereof. However, effect size would seem to better capture the important issue. If sample sizes are somewhat small (or there are is lot of variability across participants) a moderate effect size could be non-significant, but a moderate effect size would be clinically important.	We agree entirely. We report effect sizes in the tables whenever available; however, authors frequently did not report effect sizes or data with which effect sizes could be calculated.
5	7. A similar point to that above, is that studies may compare a mild TBI group to a non-mild TBI control group and find differences (as some studies did). However, those two groups may differ on other important factors as well that could explain group differences (e.g., education, race, age, degree of comorbid mental health or medical health conditions). It seems to me that you would want to address this issue if you can for those studies that reported group differences. An example of this is on the bottom of page 26.	Agreed. We highlighted any statistical adjustment or other adjustment for variation across groups in the text and tables; unfortunately, most studies did not provide this information, and this lack of adjustment for potential confounders is a contributing factor to the low study quality ratings for this body of literature.
5	8. Page 36: The term “associated with mTBI” is used in talking about mental health disorders and symptoms. This implies some actual association, when it is more likely that they are simply comorbid factors both due to deployment-related (or life-related) experiences that are risk factors for both mTBI and comorbid mental health conditions. -- I would suggest using the term “comorbid” rather than “associated”.	We have made this change to the paper.

Reviewer	Comment	Response
6	<p>Studies reporting mean cognitive performance scores are criticized within the report because they cannot provide prevalence estimates of impairment. However, the implication that report of percentage of participants scoring below an impairment cut-off or below a certain standard score would yield a good prevalence estimate of cognitive impairment does not take into account the premorbid cognitive abilities of participants. Clinically relevant cognitive impairment is typically thought of as an intra-individual decline. Scores normally thought of as below average may not indicate acquired impairment but instead reflect the innate potential of the individuals; similarly, above average scores may indicate a cognitive decline (i.e., impairment) in an individual with superior cognitive potential. Thus, the suggestion in the report does not go far enough. To best estimate the prevalence of cognitive impairment following a TBI, prospective, longitudinal measurement would be necessary. This is typically not feasible, but the report should nevertheless avoid implications that use of normative data without regard to the individual’s baseline potential would yield accurate prevalence estimates of cognitive decline. The paper states this in the Summary and Discussion section, but it is also important to mention it earlier when discussion impairment cut-offs based on standardized scores.</p>	<p>Noted. We have included this discussion both within the section on cognitive outcomes, and within the discussion section.</p>
7	<p>Overall comments: The synthesis was very well written and took a comprehensive approach to examining the extant literature on mTBI in U.S. service members and military. The synthesis was inconsistent with respect to its use of citations in the text, and its repetition of acronyms and abbreviations. Other editorial and substantive comments (line numbers not included in reviewed drafts):</p>	<p>Noted. Thank you.</p>
7	<p>P. 1, Background: OEF/OIF should be defined. Also, what about OND?</p>	<p>This change has been made. No studies reported outcomes for OND Veterans and the information we describe in the background is specific to OEF/OIF Veterans.</p>
7	<p>P. 1 - Methods: Define WHO</p>	<p>This change has been made.</p>
7	<p>P. 3 - define LOC, PTA</p>	<p>This change has been made.</p>
7	<p>P. 3 – First introduction of NSI – Neurobehavioral Symptoms (no S) Inventory</p>	<p>This change has been made.</p>
7	<p>P. 3 – second to last paragraph, two instances of “reported that” – delete one</p>	<p>This change has been made.</p>
7	<p>P.3 – second to last paragraph – “mTBI).” ß no reference to an open parentheses, so delete the)</p>	<p>This change has been made.</p>
7	<p>P.4—define DTI, MRI</p>	<p>This change has been made.</p>

Reviewer	Comment	Response
7	<p>P.5 – last paragraph: the recommendation to include imaging results is noted; however, such a recommendation is not feasible in terms of equipment, manpower, participant willingness to participate (might lead to biased self-selection), timing, and costs. This recommendation also needs to be reconciled with the VA/DoD Clinical Practice Guidelines (see excerpt below). Though, the point is taken that if imaging studies, especially functional fMRI or DTI, were performed more frequently, then perhaps there would be notable distinctions between “normal” and “mild TBI” states.</p> <p>From: VA/DoD (p. 16)</p>	<p>Noted, and we have changed this recommendation slightly to be more consistent with this comment.</p>
7	<p>In addition to traditional imaging studies, other imaging techniques such as functional magnetic resonance imaging, diffusion tensor imaging, positron emission tomography scanning; electrophysiological testing such as electroencephalography; and neuropsychological or other standardized testing of function have been used in the evaluation of persons with TBIs, but are not considered in the currently accepted criteria for measuring severity at the time of the acute injury outlined in Table A -1</p>	<p>Noted, and the criteria presented are consistent with the VA/DoD criteria.</p>
7	<p>Abbreviations Table: There were some abbreviations throughout the text that were not “formally” defined before their first use, so these are included along with others: (a) AOC = Alteration of consciousness/mental state ⁵ (based on VA/DOD guidelines) p.5; (b) BAMC (last row) = Brooke Army Medical Center, p. 16; (c) C&P = Compensation & Pension , p. 21; (d) CTE = Chronic Traumatic Encephalopathy, p. 10; (e) EFP = Explosively formed penetrator (?) – used on p. 17 – define; (f) mBIAS – “symptoms” misspelled, p. 6; (g) NSI = Inventory, p. 6; (h) PI = Principal Investigator, p. 9 (or just say “Principal Investigator on p. 9); (i) VHA = Veterans Health Administration (like on p. 8), p. 7; (j) For SCID, indicate Axis I, not Axis 1, p. 7; (k) Define/include VACO and PM&R, as introduced on p. 8; (l) define WHO on p. 7 (mentioned on p. 8)</p>	<p>These changes have been made.</p>
7	<p>P. 8 – cite the studies from which the “12 to 23 percent” are derived. Also consider citing Hendricks et al. (in press, Brain Injuries), who examined VA comprehensive TBI evaluation (CTBIE) data and found: “In the study population, 21.6% screened positive for potential TBI and 54.6% of these had an electronic record of a CTBIE. Of those with CTBIE records, evaluators confirmed TBI in 57.7%, yielding a best estimate that 6.8% of all those screened were confirmed to have TBI.”</p>	<p>This citation has been added.</p>
7	<p>P. 8 – second paragraph: “fame” should be “frame”</p>	<p>This change has been made.</p>

Reviewer	Comment	Response
7	P.8 – second paragraph: “factors unique to combat deployments.” I would take out “combat,” (maybe replace with “military,” but not sure if that’s necessary), since deployment-related conditions (e.g., noise in the general military environment), separate from combat, may uniquely account for experiencing post-concussive symptoms. Plus, since women are restricted from some combat roles, the use of “combat” here may minimize what women potentially experience.	This change has been made.
7	P. 8 – misplaced semi-colon in second paragraph,	This change has been made.
7	P. 8 abbreviate “Veterans Health Administration” as VHA;	This change has been made.
7	P. 8 Define PM&R and VACO	This change has been made.
7	P. 9 Search strategy – can now call it “mTBI”	This change has been made.
7	P. 9 Last paragraph – why not include “post-traumatic amnesia” along with AOC, LOC?	This change has been made.
7	P. 9 – last paragraph, discussion about “severity of sequelae” gets a little confusing, because TBI sequelae can be defined as either AOC, LOC, or PTA (which the reader might be primed for, since there was just discussion about these in the previous sentence), or TBI residual symptoms (e.g., headache, vestibular, pain, auditory, visual, etc. impairment). I don’t know if “the Severity of sequelae” sentence needs to be here, but I understand that it makes the point that very specific criteria are used to categorize TBI severity, and it’s important to note that the severity of mental/physical symptoms following a TBI event are not used to classify severity. I would replace “sequelae” with “symptoms,” and then provide a few NSI-22 symptoms to orient the reader. It might also be helpful to briefly state severity is based on the duration of LOC, AOC, or PTA. You can then list the criteria (e.g., LOC < 30 min), or refer to the reader to a more detailed description, as you currently do.	This change has been made.
7	P. 10 Identify DoD in first paragraph	This change has been made.
7	P. 10 – identify CTE in “Outcomes” paragraph	This change has been made.
7	P. 14, Figure 1: Bullet points in the last box to the right would help the criteria stand out more.	We left out bullet points for space reasons.
7	P. 15, Table 1: Barnes et al., spell out “medical”	We have corrected this mistake.
7	P. 16-17 Cooper, Mercado-Couch, et al. – BAMC should be defined earlier (e.g., in the Cooper et al. citation above it in the table); Should American “College” be “Congress?”; Was not clear about 10 participants excluded due to PTA suggesting a more severe TBI... Is it that the duration of PTA was longer than the criteria for mTBI, as specified by ACRM?	This change has been made.

Reviewer	Comment	Response
7	P. 17 Drag et al., define “C and P,” also, in mTBI definition, why not just call it “AOC?”	This change has been made.
7	P. 17 Gaylord et al., American College or Congress?; defined “EFP,” as previously noted.	This change has been made.
7	P. 18 Gordon et al., Time since Injury – define unit of time.	This change has been made.
7	P. 18, Kennedy, Leal et al., last column, just say MVA instead of spelling it out	This change has been made.
7	P. 19 MacDonald et al., use “TBI” consistently – sometimes it’s spelled out (see first and last columns)	This is a quote so it is spelled out as in the original text.
7	P. 19 Morrisette et al., spell “through,” not “thru.”	This change has been made.
7	P. 19, Nelson et al., reference 34: define what “forensic” context is.	This change has been made.
7	P. 20, Patil et al., Based on “VA”/DoD Consensus definition (reversed as written).	This change has been made.
7	P. 20 Ruff, Riechers et al., “Department of CVAMC” needs editing; also, in description for three groups of Veterans, for the first two indicate “deployment-related” LOC? Not sure what “combat Veterans without “LOC” means – is this a control group or did they have AOC or PTA?; In the mTBI definition column, not sure what the definition is referring to – did Veterans have to have LOC or AOC following the TBI, PLUS PTA?	These changes have been made.
7	P. 21 Schiesher et al., mTBI definition column – delete ©	This change has been made.
7	P. 21 Spencer et al., C&P needs to be defined earlier	This change has been made.
7	P. 21 Swick et al., Combat Veterans diagnosed, not “diagnoses”	This change has been made.
7	P. 21 Theeler et al., mTBI definition column – use DVVIC acronym	This change has been made.
7	P. 23 Third from last paragraph – can abbreviate LOC and AOC – make clear, if this is the case, that participants demonstrated better overall cognitive functioning “post-intervention?”	This change has been made.
7	P. 23 – wasn’t clear in 2nd to last paragraph – second line: Should this be “NOT” reported, or should “NOT” be deleted?	We have clarified this sentence.
7	P. 24, processing speed row, Key Question #1 column, not sure why “possible” exceptions is there – delete that word?	This change has been made.
7	P. 26, first paragraph – could you cite the 7 studies here?	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.

Reviewer	Comment	Response
7	P. 26, first paragraph, 2nd line, instead of “with” TBI, replace “with” with “experienced” or indicate “with mTBI history.” Also make this correction in Key Question 2 paragraph, first line.	This change has been made.
7	P. 26, Visuospatial abilities, first sentence, cite the 5 studies here.	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 26, last full paragraph: Just give the abbreviations/acronyms, no need to spell out the neuropsychological test names.	This change has been made.
7	P. 27, first paragraph, 3rd line: instead of saying “significant correlations,” indicate directionality, since it seems like visuospatial abilities and MH conditions would be negatively correlated.	This change has been made.
7	P. 27, first “Memory” paragraph, mid-way through, again report direction of association.	This change has been made.
7	P. 27, Key Question 1 paragraph: just report abbreviations	This change has been made.
7	P. 27, Key Question 1 paragraph, 5th from last row, use “longer” instead of “much more.”	This change has been made.
7	P. 27, second to last row, use (C&P) for compensation and pension, since this would have been introduced previously.	This change has been made.
7	P. 28, 1st paragraph, second from last row: Use LOC and PTA instead of spelling them out.	This change has been made.
7	P. 28, Attention/Concentration paragraph, cite the 7 studies after mentioning them.	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 28, Attention/Concentration paragraph, last line can read “....OR PTSD was not consistently associated with outcomes.”	This change has been made.
7	P. 28, Key Question 1 paragraph, do not spell out neuropsychological test names.	This change has been made.
7	P. 28, last paragraph, “forensic” mentioned a couple of times – define what exactly this setting is.	This change has been made.
7	P. 29, first paragraph, indicate direction of associations; do not spell out LOC, PTA	This change has been made.
7	P. 29, Key Question 1 paragraph (here the 9 studies are cited, very helpful!); , do not spell out neuropsychological test names.	This change has been made.

Reviewer	Comment	Response
7	P. 29, 30 Key Question 2 paragraphs for each page, can say C&P instead of spelling it out; on p. 30, no need to define what a C&P evaluation is here, define it when first introduced	This change has been made.
7	P. 30 Effort/Motivation paragraph – regarding “with mTBI,” see comment #41	This change has been made.
7	P. 30, Key Question 2 paragraph, see comments #56, 58	This change has been made.
7	P. 31, Key Question 1 paragraph, do not spell out neuropsychological test names.	This change has been made.
7	P. 31, Key Question 1 paragraph: There is discussion about “non-mTBI” participants – are these participants who never experienced a TBI (non-mTBI history) or who do not currently have mTBI (symptoms resolved?)	This has been clarified
7	P. 31, Key Question 2 paragraph: compensation and pension evaluation can be abbreviated	This change has been made.
7	P. 31, Self-reported Cognitive Problems paragraph. I’m wondering if “service connection” should be considered a “risk factor,” as it’s really a proxy for a disabling injury that was sustained or aggravated during military service. The point is taken that “service connection” is a short-hand for the latter, so this might be a matter of semantics.	Agreed, and this is how it is described in the primary study.
7	P. 31 – last word – instead of saying “control,” say “control group.”	This change has been made.
7	P. 32 – summary of Physical health results first paragraph – it would be helpful to cite the 16 studies here.	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 32 – summary of Physical health results first paragraph – to be consistent, remove dashes from time-since-injury.	This change has been made.
7	P. 32 – second from last paragraph, no need to spell out NSI – already introduced.	This change has been made.
7	P. 33 – Table 3 – first row, similar to comment 63, “referral to neurology clinic for headaches” is listed as a risk factor, but is likely a proxy for headache severity/frequency. Again, this might just be a matter of semantics and readers will understand the implication. Without reviewing the article, not sure if any specifics were given for why a patient might be referred to a neurology clinic, so I understand that this might be the only unit of measurement.	Agreed—and this is how it was described in the primary study.
7	P.33 – Table 3 – some articles from JRRD 49(7) would be relevant here (if you increase the time frame of your search)	We have added these articles.
7	P. 34 – First paragraph – when describing the pain level scale, please provide anchors, 0 (no pain at all?) to 10 (very severe pain?) scale.	This change has been made.

Reviewer	Comment	Response
7	P. 34 – first paragraph for “Headaches,” last line about referral to neurology clinic for headaches, see comment #68.	Noted.
7	P. 34 – Key Question 1 paragraph – here, the 9 studies being alluded to are actually cited (good!). Can also just say “NSI” here, no need to spell it out.	This change has been made.
7	P. 34 – Key Question 2 paragraph: Provide more detail in the first sentence – describe all groups being compared	This change has been made.
7	P. 34 – Key Question 2 paragraph, last sentence – see comment #68.	Noted.
7	P. 35 – “Vision” paragraph: cite the one study being discussed	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 36 – “Nausea/Appetite” first paragraph - cite the specific studies you highlight, such as the study describing mild to moderate effects of mTBI population, PTSD studies, mixed results study	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	PP. 36-37 –Summary of Mental Health Results section – cite the studies where you describe 20 studies, two studies, PTSD, alcohol abuse, etc.	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 38, Table 4, see comment #68	Noted.
7	For the PTSD sections, 17 studies are alluded to in the first paragraph, and then alluded to again in the Key Question paragraph, and then cited. Just make sure there is consistency between the sections throughout the report in terms of when you cite.	Noted.
7	P. 39 - for Key Question 2, could you add specifics? For example, for study #27, what was the association between the PCL re-experiencing cluster and blast-exposure?	This change has been made.
7	P. 39 – For the anxiety paragraph, cite the studies that are being alluded to.	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 39—Key Question 1 – just say “NSI”	This change has been made.
7	P. 40 – Key Question 1 – abbreviations	This change has been made.

Reviewer	Comment	Response
7	P. 40 – Substance Use disorders first paragraph – cite studies alluded to	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 40 –Suicide first paragraph - cite studies alluded to	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 41 – Other Mental Health Outcomes first paragraph – cite studies alluded to	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 41 – Other Mental Health Outcomes first paragraph – aren’t “frustration” and “irritability” in the NSI affective cluster (e.g., Meterko et al., 2012, Journal of Head Trauma Rehabilitation, 27(1), 55-62 Psychometric assessment of the Neurobehavioral Symptom Inventory-22: the structure of persistent postconcussive symptoms following deployment-related mild traumatic brain injury among veterans.)	Yes, and we would consider these part of the mental health related outcomes section for this report.
7	P. 41 - Key Question 1 – abbreviations	This change has been made.
7	P. 41 - Key Question 1, second to last sentence – period before 12,20 citations.	This change has been made.
7	P. 41 – Summary of functional/social outcome results: cite studies alluded to	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 41 – Summary of functional/social outcome results – make clear throughout the paragraph whether patients with mTBI, patients without mTBI, or results collapsed across both groups are being discussed.	This change has been made.
7	P. 43 – Sleep introduction paragraph – what are “positive neurological findings?”	This change has been made.
7	P. 43 – Sleep Key Question 1 – abbreviations	This change has been made.
7	P. 46 – In the “Results” paragraph: Throughout the paragraph, the word “obtaining” is used several times. I think a better word here would be “exhibiting.”	This section has been removed and edited.
7	P. 46 –Key Question 1: Abbreviations	This section has been removed and edited.
7	P. 46 –Key Question 1: I found the first sentence difficult to follow – please add some additional punctuation and clarifying language.	This section has been removed and edited.
7	P. 46 – Key Question 2: the tense was different in this paragraph than in others. Use “exhibit” instead of “obtain.”	This section has been removed and edited.

Reviewer	Comment	Response
7	P. 46— Summary of service utilization/costs results: Please cite studies alluded to.	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 48 – Key Question 1: Table X = Table 7?	This change has been made.
7	P. 50 – For mental health outcomes, it’s acknowledged that PTSD is a focus for researchers, but I’m wondering if this section could be rounded out by also examining a few other mental health conditions that are typically of interest: depression, non-PTSD anxiety, and substance use disorders? As a starting point, Thomas W. McAllister has published on mild TBI in civilian populations and its after effects (e.g., cognitive, mental health conditions; Silver JM, McAllister TW, Arciniegas DB. Depression and cognitive complaints following mild traumatic brain injury. Am J Psychiatry. 2009; 166: 653-61.)	Agreed, and due to space limitations, we provided an expanded discussion of PTSD at the request of our stakeholders, but have not provided an expanded discussion of other outcomes, instead referring readers to other reviews and studies.
7	P. 50—3rd paragraph of the MH Outcomes section: instead of “causal factors,” maybe use the term “event-related?”	This change has been made.
7	P. 50—in the “Not Surprisingly” paragraph, cite the “aforementioned” literature base. Also, in the last sentence of this paragraph, it might be more clear to say, “...related to mTBI versus other factors, such as those that are deployment-related, are not clear.”	This change has been made.
7	P. 51—first partial paragraph at top, “When individuals experience the mTBI as traumatic,” consider replacing “experience” with “perceive,” since, by definition, the experience of mTBI is traumatic, at least physiologically/functionally.	This sentence has been clarified.
7	P. 51-first full paragraph, starting with “The results,” I think more detail could be added here, such as citations, especially when you cite specific figures. Consider rewording the last sentence as: However, these high prevalence estimates may differ from results observed in civilian populations, as they may be related to unique deployment-related factors, such as combat, rather than, specifically, to the presence of mTBI.	This change has been made.
7	P. 51 – Imaging/biomarkers paragraph: cite the one study here.	We are providing summary paragraphs without citations for each section, which are followed by the cited results sections by key question. This will be consolidated for the published article version of the report.
7	P. 51 – “Although biomarkers” sentence: Do you mean “as prognostic tools among those with... “severe” TBI or with “moderate to severe” TBI? (not sure if severe was being used as a category, or if this was meant as “not mild.”	This change has been made.

Reviewer	Comment	Response
	P. 51 – “Although biomarkers” paragraph: after introducing chronic traumatic encephalopathy, put “(CTE)”	This change has been made.
7	P. 51 last paragraph, 2nd sentence, put “imaging” after “functional”; 5th sentence, use “DTI” instead of spelling it out; instead of “demonstrated” throughout this paragraph, use “observed;” Cite	This change has been made.
7	P. 52 – first paragraph, 2nd sentence; add “patients with” mTBI.	This change has been made.
7	P. 52 – first paragraph, 6th sentence, should be an “in” between “decreases memory”	This change has been made.
7	P. 52, first paragraph, 9th sentence, reword: differences between “individuals with mTBI” and “individuals without mTBI” or “individuals in the control group.”	This change has been made.
7	P. 52, first paragraph, 9th sentence – reword, such as “fMRI studies found activation differences between individuals with mTBI and individuals in the control group during cognitive and behavioral tasks consistent with...”	This change has been made.
7	P. 52, first full paragraph, first sentence, reverse last two words so that it reads: mTBI neuroimaging; cite which studies found increased vs. decreased FA; instead of “controls,” say “control participants.”	This change has been made.
7	P. 54 – first paragraph, first sentence, say “universal” limitation (instead of across the board).	This has been re-worded.
7	P. 54 – first paragraph, what is the evidence that participants aren’t blinded to study hypotheses? It’s my sense that patients are told that the purpose of the study is to “examine differences,” “observe,” etc., and aren’t informed about specific directional hypotheses until after study completion, if at all.	No studies reported that patients were blinded, and therefore we cannot assume that any were blinded.
7	P. 54 – second paragraph, 3rd-4th sentences; “wide variety of tools used to assess each outcome of interest.”	This change has been made.
7	P. 54 –Conclusions, 2nd sentence, I would say, “The literature reviewed here,”	This change has been made.
7	P. 54 – Conclusions, 4th sentence, instead of saying “negative outcomes,” consider: “Though a significant portion of individuals who have experienced an mTBI report long-term mental and physical health symptoms””not significantly different from individuals who “did not experience mTBI” or “served as controls.”	This section has been re-worded.
7	P. 54- Conclusions, 7th sentence, I would stay away from “outcomes,” and say “self-reported symptoms,” because we don’t know whether symptoms/conditions are caused (i.e. an “outcome”) by the mTBI.	This change has been made.
7	P. 54 –Conclusions, 8th sentence from last – instead of “do not have mTBI,” I would say “who have not experienced mTBI” or “who do not have mTBI history”	This change has been made.

Reviewer	Comment	Response
7	P. 54 –Conclusions, 7th sentence from last, “instead of saying “is largely influenced by” I would say “can be accounted for” by other factors that are deployment-related, rather than....”	This change has been made.
7	P. 69 – Cameron et al., “five-difit”, should be “five-digit”	This change has been made.
7	P. 72 – Hoge et al., “seeking starts” should be “seeing stars”	This change has been made.
7	P. 73- Luis et al., “loose” should be “lose”	This change has been made.
7	P. 87 & 96-Nelson et al. – why aren’t the p-values listed?	We have listed results according to what was reported in the primary studies and have put labels used by study authors in quotes to indicate a direct quote in our tables.
8	Excellent that limitation of non-reporting of impaired subgroups is emphasized.	Noted. Thank you.
8	Text is somewhat repetitive and disorganized with regards to reporting of the conclusions reached. For example, conclusions are reported on p.13 within a paragraph on Literature Flow.	We have re-ordered the presentation of findings.
8	Presentation of results needs revision. Tables contain a great deal of text and no legends. Numbers are occasionally presented without units (e.g. p.18 Gordon et al; time since injury: 20.1 (weeks?)). Table 1 should include each study’s outcome measures and preferably study hypothesis. It was good to see that effect sizes are reported in the tables in the appendices, but effect sizes are missing for most studies (if not reported, these can be calculated). Also, in cases where significance criteria have been corrected for multiple comparisons (e.g. Table 5), this should be indicated.	We have made these corrections to the table. For space reasons, we have presented some information in appendix tables rather than in text. We report the data as reported in the studies without calculating effect sizes when the authors did not provide this information; however, we are considering providing this calculated information in the article version of the report.
8	Discussion of imaging and biomarkers, p.51: it is mentioned twice that functional imaging studies have “failed” to show differences in performance along with differences in brain function, and this point would benefit from clarification. One explanation, supported by activation patterns, is that mTBI patients are able to accomplish similar test performance to uninjured controls through greater recruitment of neural resources. (e.g. McAllister, 2001, Neuroimage . 2001 Nov;14(5):1004-12.Differential working memory load effects after mild traumatic brain injury.)	We have moved all discussion of imaging results to the discussion due to searching limitations and refer readers to more comprehensive reviews of this literature.
8	p. 26 – Section title “Verbal and Old Learning” inappropriate as the tests described in this section measure vocabulary and knowledge, not learning (implies active learning and memory).	We have changed this title.
8	p. 26; paragraph 5: “...without mTBI on the RBANS Visuospatial/Constructional subscale.” This sentence does not match information presented in table 1b for reference 33	We have made sure these results are correct and consistent with the table.

Reviewer	Comment	Response
8	Within Appendix E, some information in the “comparison group description column” does not appear to describe a comparison group, but rather a covariate (e.g. PCL score).	Yes, this indicates continuous variables rather than comparison groups, and as such, correlation results are presented for these studies.
8	Readability comments:	Noted as below.
8	p.1 paragraph 1: “...and its associated post-concussion symptoms is...” should be “are”	Refers to mTBI, not the symptoms, therefore left as “is.”
8	p. 1 paragraph 1: “...a TBI while deployed).” The parenthesis does not have a partner.	This change has been made.
8	p.1 paragraph 2 and p.8 paragraph 2: “...balance problems) beyond this time fame;” should be “frame”	This change has been made.
8	p. 1 paragraph 2: “...often require the attention from a range of health care professionals...” should remove “the”	This change has been made.
8	p.3 paragraph 4: “lengthly” should be “lengthy”	This change has been made.
8	p. 3 paragraph 5: “...Veteran/military participants without mTBI).” The parenthesis does not have a partner.	This change has been made.
8	p. 9 paragraph 6: “disruption of brain function (e.g. altered of consciousness...)” should be “alteration”	This change has been made.
8	p. 28 paragraph 4: “six studies reporting...” Should be “reported”	This change has been made.
8	p. 41 paragraph 3: “Axis 1” should be “Axis I”	This change has been made.
8	p. 54 paragraph 3: “...we excluded many studies which proported to study mTBI” should be “purported”	This change has been made.
9	The work of Drs Cockerham and Goodrich might be beneficial in the section on Vision as it addresses occult visual deficits in this patient population. This can often be conflated with self-reported complaints, as previously established mechanisms of assessment were deemed not sensitive in detecting these abnormalities. Additionally, as this data is prospectively collected at all Polytrauma sites, could this data be incorporated in the analyses.	We have reviewed this literature and though it provides important information related to vision outcomes, we did not find studies meeting our inclusion criteria for this report due to the populations examined.
9	Would also recommend additional references regarding TBI incidence—pg #8 Introduction. The point could be substantiated by and WHO data or CDC Data.	We have re-worked the introduction for the report.

Reviewer	Comment	Response
10	<p>In a significant number of veterans and service members who have incurred a blast or non-blast related mTBI have lead to persistent or chronic post-concussion syndrome (PCS). Multiple studies have reported PCS-like symptoms among Veterans many years after mTBI. Agree that published studies to date has been unable to identify all the potential risk factors and a major causative role other than mTBI. ICD-10 and DSM-4 criteria have been established for diagnosing PCS and differ somewhat. There continues to be a lack of consensus regarding PCS, ICD-10 guidelines limit the symptoms to within 4 weeks of injury, while DSM-4 criteria requires symptom onset shortly after injury, but persistence at least 3 months. Despite these diagnostic guidelines, evidence suggests that symptoms can appear immediately, or weeks to months after the initial injury (Ryan et al.2006) and recent studies have reported PCS-like symptoms among Veterans many years after mTBI. (Scholten et al.2012). While these persistent symptoms are known to complicate return to work/duty and negatively affect quality of life, their trajectories and time courses are not understood and diagnosis remains challenging and relies mostly on self-report of complex symptomatology rather than objective, quantitative or biological measures. The reasons why people recover slowly or fail to recover fully from mTBIs is not known and there are no current methodologies for diagnosis or prognosis of PCS. Identifying the cognitive, clinical, and serum biomarkers that accurately diagnose veterans or service members with persistent symptoms is critical to our understanding of long-term outcomes in this patient population and needs to explored further.</p>	<p>We have expanded the discussion section to include some of these ideas.</p>
11	<p>This was a challenging area with limited available published studies. I would suggest plainly stating in the early overview section that there are no prospective randomized RCTs.</p>	<p>We have stated this in the executive summary and the body of the report.</p>
11	<p>The overview also seems somewhat contradictory when you report that imaging findings are of low strength of evidence yet your recommendations indicate that prospective studies should be designed to report imaging findings.</p>	<p>We have rearranged and clarified the imaging findings and recommendations.</p>
<p>Optional Dissemination and Implementation Questions</p>		
<p>5. Are there any VA clinical performance measures, programs, quality improvement measures, patient care services, or conferences that will be directly affected by this report? If so, please provide detail.</p>		
5	<p>There really is no new information here. Conclusions are consistent with other reviews. However, making this available to national Polytrauma calls or conferences, and Mental Health/PTSD calls and conferences would help reinforce these findings.</p>	<p>Noted. We plan to make the report findings available in a variety of formats, including VA intranet, a published article, presentation at a national neuropsychology conference, and through a VA cyber-seminar. We will consider how to expand the audience as you recommend.</p>

Reviewer	Comment	Response
6	Yes. The results raise the question of the context in which mTBI may be best treated within VA. The results also have implications for compensation and pension decisions regarding mTBI.	Noted, though we caution our readers not to make strong inferences based on the low quality literature available for synthesis in this report. Relying on these report findings in conjunction with related research on civilian populations with mTBI will provide the strongest available foundation for such weighty decisions.
7	Because “time since injury” was cited throughout the document as being related to impairment, but also noted to be missing in many studies, I think it would be very important to capture this variable in national VA databases, such as the Comprehensive TBI Evaluation database. If this information is available in any DoD databases, like Defense Manpower Data Center (DMDC), then a data exchange between DoD and VA would benefit studies by having these data located in one central source, thus reducing the risk of error through patient self-report.	We have added this to the discussion.
8	Given the limited nature and low quality of literature, it is not possible to reach firm conclusions. There is nothing to be implemented at this time.	Noted, and we encourage readers of this report to consider the results in conjunction with findings from civilian literature in order to make conclusions based on the best available evidence.
9	Vision Screening in the Inpatient/Polytrauma Units might be affected.	Noted.
10	Yes. Polytrauma/ TBI System of Care.	Noted.
11	Polytrauma System of Care, can be disseminated on one of the national calls and emailing providers with link to report	Noted.
6. Please provide any recommendations on how this report can be revised to more directly address or assist implementation needs.		
1	You might consider adding review of any studies that looked at multiple mTBI’s. It is a common “complication” Also, on page 1, I’d suggest clearly delineating the difference between cognitive performance and symptom complaints. While it’s true that the literature suggests complete cognitive performance recovery by 3 months (or even 7 days in sports literature), there is a difference between performance and symptoms. Furthermore, the civilian literature suggests that PCS symptoms in fact do not *persist* (See Meares et al) and such the use of the term ‘persist’ may be incorrect. “Presence” may be more accurate.	We have made these changes throughout the report.
5	There is no “So What” section. However, one could potentially make the suggestion that since outcomes do not differ following mTBI, that all the DoD/VHA time, energy, and attention devoted to this matter may be a less than ideal use of resources. For example: Do we really need to continue to screen for mTBI? Are the required Comprehensive TBI Evaluations following a positive TBI screen, really needed and a good use of resources?	Noted, and we have expanded our discussion of these points in the report.

Reviewer	Comment	Response
5	Are there better ways to meet the needs of those symptomatic returning service members and veterans than focusing on mTBI, when mTBI does not appear to be the factor explaining the symptoms and problems?	Noted, and we have expanded our discussion of this point in the report.
6	No recommendations. This is a thorough report with well-reasoned conclusions.	Noted. Thank you.
7	This report indicates that many of the mTBI studies performed with service members and Veterans are methodologically limited and provide low strength evidence. Because implementation should be based on strong evidence, it doesn't appear that this report should make any health services related recommendations on implementation, but should recommend that VA leaders prioritize research funding to ensure high quality research, and develop mechanisms (e.g., databases, standard communications between DoD and VA) that assist researchers in obtaining reliable data.	We agree that ideally, recommendations should be made based on strong evidence. However, in the absence of strong evidence, then the best available evidence should be the basis on which treatment and policy decisions are made. We have made cautious recommendations consistent with the best available evidence for treatment and policy as well as making strong recommendations for further high quality research as you suggest.
8	Readability and typographical errors should be addressed. Presentation of results should be revised. Consider other data presentation modes in addition to tables. Imaging and biomarkers could possibly be removed and examined in a separate report with selection criteria that are more appropriate for these kinds of studies.	Noted, and we have incorporated your suggested edits.
10	The current focus in the TBI clinics is the CHRONIC effects of mTBI 3-10 years post injury with retained sequale of the initial injury which does not completely follow the recovery pattern of the civilian mTBI population. In the civilian population the symptoms are transient and self-limiting, with apparent full recovery occurring from minutes to several weeks following injury (Levin et al., 1997) which is distinct from our veteran/service member population who have persistent symptoms and/or functional limitations (Iverson et al., 2006; Ruff et al., 1996). There needs to be further investigation into the etiology and treatment of these chronic/persistent PCS symptoms.	We agree and have expanded our discussion of treatment implications.
7. Please provide us with contact details of any additional individuals/stakeholders who should be made aware of this report.		
5	DCoE, DVVIC, VBA?	Noted.
6	No specific recommendations other than VA polytrauma staff.	Noted.
7	HSR&D/QUERI/RR&D leaders responsible for prioritizing funding, inclusive of and addition to: David X. Cifu, MD Nina A. Sayer, PhD Joel Scholten, MD Doug Bidelspach, MPT VA TBI/Polytrauma Clinic Directors	Noted.

Reviewer	Comment	Response
8	Katherine Helmick, Deputy Director, Defense and Veterans Brain Injury Center katherine.helmick@tma.osd.mil	Noted.
10	The following stakeholders should be made aware of this report: VHA Polytrauma System of Care which include Polytrauma Rehabilitation Centers (PRC), Polytrauma Transitional Rehabilitation Programs (PTRP), Polytrauma Network Site (PNS), Polytrauma Support Clinic Team (PSCT), and Polytrauma Point of Contact (PPOC). In addition the Military Health System's TBI clinics, Defense and Veterans Brain Injury Center (DVBIC) and National Intrepid Center of Excellence (NICoE) satellite TBI clinics.	Noted.